
Test2 1D residue occlusion: frozen representative cohort
sequence_101_101 | 1D_only_correct | y=0
sequence_34_34 | 1D_only_correct | y=1
sequence_388_388 | 3D_only_correct | y=0
sequence_23_23 | 3D_only_correct | y=1
sequence_143_143 | both_correct | y=0
sequence_1_1 | both_correct | y=1
sequence_336_336 | both_incorrect | y=0
sequence_8_8 | both_incorrect | y=1
M D G E G H D I N I W G A R M S P S P A A A P V S A T R G A P W S G C E G C P S R A A D R R C V C H
V I G G D E D I N E H R S L A L M Y S W S H R F I G L I
T S S G C P N F E S L V R
A C L D I G N S C R E D
I T F S K I Y R S C K S N S V C G N Q K C V R G R C V
M L K F I L A L A L F L H L T M E A S A A C K G K K C P P T G F V G M R G
K A L Y A P F D A T F P N Q N Q T R N K A V D T A P C E W Y R R V Y
M K K T F L P I F L V I L L A S Y A L A N P Q V T F S R D W G P G K K
0
10
20
30
40
Residue position (amino-acid letters shown in cells)
0.2
0.1
0.0
−0.1
−0.2
Signed within-peptide L1-normalized attribution