

Probability and Stochastic Modelling 1

Semester 1, 2022

Dr Alexander Browning

TARANG JANAWALKAR

This work is licensed under a Creative Commons
“Attribution-NonCommercial-ShareAlike 4.0 International” license.



Contents

Contents	1
1 Events and Probability	3
1.1 Events	3
1.2 Probability	4
1.3 Circuits	5
2 Independence	5
2.1 Probability Rules with Conditional	5
2.2 Conditional Independence	6
2.3 Disjoint Events	7
2.4 Subsets	7
3 Total Probability	7
4 Combinatorics	8
4.1 Ordered Sampling with Replacement	8
4.2 Ordered Sampling without Replacement	8
4.3 Unordered Sampling without Replacement	9
4.4 Unordered Sampling with Replacement	9
5 Random Variables and Distributions	9
6 Random Variables	9
6.1 Probability Distributions	9
6.2 Quantiles	10
6.3 Summary Statistics	11
6.3.1 Transformations	11
7 Special Discrete Distributions	12
7.1 Discrete Uniform Distribution	12
7.2 Bernoulli Distribution	12
7.3 Binomial Distribution	13
7.4 Geometric Distribution	13
7.4.1 Alternate Geometric Definition	14
7.5 Negative Binomial Distribution	14
7.5.1 Alternate Negative Binomial Definition	15
7.6 Poisson Distribution	15
7.7 Modelling Count Data	15
8 Special Continuous Distributions	16
8.1 Continuous Uniform Distribution	16
8.2 Exponential Distribution	16
8.3 Memoryless Property	17
8.4 Normal Distribution	17

8.5	Standard Normal Distribution	17
-----	--	----

1 Events and Probability

1.1 Events

Definition 1.1 (Event). An event is a set of outcomes in a random experiment commonly denoted by a capital letter. Events can be simple (a single event) or compound (two or more simple events).

Definition 1.2 (Sample space). The set of all possible outcomes of an experiment is known as the sample space for that experiment and is denoted Ω .

Definition 1.3 (Intersection). An intersection between two events A and B describes the set of outcomes that occur in both A and B . The intersection can be represented using the set AND operator (\cap) — $A \cap B$ (or AB).

Definition 1.4 (Disjoint). Disjoint (mutually exclusive) events are two events that cannot occur simultaneously or have no common outcomes.

Theorem 1.1.1 (Intersection of disjoint events). *The intersection of disjoint events results in the null set (\emptyset).*

Lemma 1.1.1.1. *Disjoint events are **dependent** events as the occurrence of one means the other cannot occur.*

Definition 1.5 (Union). A union of two events A and B describes the set of outcomes in either A or B . The union is represented using the set OR operator (\cup) — $A \cup B$.

Definition 1.6 (Complement). The complement of an event E is the set of all other outcomes in Ω . The complement of E is denoted \bar{E} .

Theorem 1.1.2 (Intersection of complement set).

$$A\bar{A} = \emptyset$$

Theorem 1.1.3 (Union of complement set).

$$A \cup \bar{A} = \Omega$$

Definition 1.7 (Subset). A is a (non-strict) subset of B if all elements in A are also in B . This can be denoted as $A \subset B$.

Theorem 1.1.4. *All events E are subsets of Ω .*

Theorem 1.1.5. *Given $A \subset B$*

$$AB = A \quad \text{and} \quad A \cup B = B$$

Corollary 1.1.5.1. *Given $\emptyset \subset E$*

$$\emptyset E = \emptyset \quad \text{and} \quad \emptyset \cup E = E$$

Theorem 1.1.6 (Associative Identities).

$$A(BC) = (AB)C$$

$$A \cup (B \cup C) = (A \cup B) \cup C$$

Theorem 1.1.7 (Distributive Identities).

$$A(B \cup C) = AB \cup AC$$

$$A \cup BC = (A \cup B)(A \cup C)$$

1.2 Probability

Definition 1.8 (Probability). Probability is a measure of the likeliness of an event occurring. The probability of an event E is denoted $\Pr(E)$ (sometimes $P(E)$).

$$0 \leq \Pr(E) \leq 1$$

where a probability of 0 never happens, and 1 always happens.

Theorem 1.2.1 (Probability of Ω).

$$\Pr(\Omega) = 1$$

Theorem 1.2.2 (Complement rule). *The probability of the complement of E is given by*

$$\Pr(\bar{E}) = 1 - \Pr(E)$$

Theorem 1.2.3 (Multiplication rule for independent events). *The probability of the intersection between two independent events A and B is given by*

$$\Pr(AB) = \Pr(A)\Pr(B)$$

Theorem 1.2.4 (Addition rule for independent events). *The probability of the union between two independent events A and B is given by*

$$\Pr(A \cup B) = \Pr(A) + \Pr(B) - \Pr(AB).$$

If A and B are disjoint, then $\Pr(AB) = 0$, so that $\Pr(A \cup B) = \Pr(A) + \Pr(B)$.

Corollary 1.2.4.1 (Addition rule for 3 events). *The addition rule for 3 events is as follows*

$$\Pr(A \cup B \cup C) = \Pr(A) + \Pr(B) + \Pr(C) - \Pr(AB) - \Pr(AC) - \Pr(BC) + \Pr(ABC).$$

Proof. If we write $D = A \cup B$ and apply the addition rule twice, we have

$$\begin{aligned} \Pr(A \cup B \cup C) &= \Pr(D \cup C) \\ &= \Pr(D) + \Pr(C) - \Pr(DC) \\ &= \Pr(A \cup B) + \Pr(C) - \Pr((A \cup B)C) \\ &= \Pr(A) + \Pr(B) - \Pr(AB) + \Pr(C) - \Pr(AC \cup BC) \\ &= \Pr(A) + \Pr(B) - \Pr(AB) + \Pr(C) - (\Pr(AC) + \Pr(BC) - \Pr(ACBC)) \\ &= \Pr(A) + \Pr(B) + \Pr(C) - \Pr(AB) - \Pr(AC) - \Pr(BC) + \Pr(ABC) \end{aligned}$$

□

Theorem 1.2.5 (De Morgan's laws). *Recall De Morgan's Laws:*

$$\begin{aligned} \overline{A \cup B} &= \bar{A} \bar{B} \\ \overline{AB} &= \bar{A} \cup \bar{B}. \end{aligned}$$

Taking the negation of both sides and applying the complement rule yields

$$\begin{aligned} \Pr(A \cup B) &= 1 - \Pr(\bar{A} \bar{B}) \\ \Pr(AB) &= 1 - \Pr(\bar{A} \cup \bar{B}) \end{aligned}$$

1.3 Circuits

A signal can pass through a circuit if there is a functional path from start to finish.

We can define a circuit where each component i functions with probability p , and is independent of other components.

Then W_i to be the event in which the associated component i functions, we can determine the event S in which the system functions, and probability $\Pr(S)$ that the system functions.

As the probability that any component functions is p , in other words

$$\Pr(W_i) = p,$$

$\Pr(S)$ will be a function of p defined $f : [0, 1] \rightarrow [0, 1]$.

2 Independence

Definition 2.1 (Conditional probability). When discussing multiple events, it is possible that the occurrence of one event changes the probability that another will occur. This can be denoted using a vertical bar, and is read as “the probability of event A given B ”:

$$\Pr(A|B) = \frac{\Pr(AB)}{\Pr(B)}.$$

Definition 2.2 (Multiplication rule). For events A and B , the general multiplication rule states that

$$\Pr(AB) = \Pr(A|B)\Pr(B)$$

Theorem 2.0.1 (Independent events). *If A and B are independent events then*

$$\Pr(A|B) = \Pr(A)$$

$$\Pr(B|A) = \Pr(B)$$

Theorem 2.0.2 (Complement of independent events). *If A and B are independent, all complement pairs are also independent. Given*

$$\Pr(A|B) = \Pr(A)$$

$$\Pr(B|A) = \Pr(B)$$

the following statements are also true

$$\Pr(A|\overline{B}) = \Pr(A)$$

$$\Pr(B|\overline{A}) = \Pr(B)$$

$$\Pr(\overline{A}|B) = \Pr(\overline{A})$$

$$\Pr(\overline{B}|A) = \Pr(\overline{B})$$

$$\Pr(\overline{A}|\overline{B}) = \Pr(\overline{A})$$

$$\Pr(\overline{B}|\overline{A}) = \Pr(\overline{B})$$

2.1 Probability Rules with Conditional

All probability rules hold when conditioning on some event C .

Theorem 2.1.1 (Complement rule with condition).

$$\Pr(\bar{A} | C) = 1 - \Pr(A | C)$$

Theorem 2.1.2 (Addition rule with condition).

$$\Pr(A \cup B | C) = \Pr(A | C) + \Pr(B | C) - \Pr(AB | C)$$

Theorem 2.1.3 (Multiplication rule with condition).

$$\Pr(AB | C) = \Pr(A | BC) \Pr(B | C)$$

In the above examples, all probabilities are conditional on the sample space, hence we are effectively changing the sample space.

2.2 Conditional Independence

Definition 2.3 (Conditional independence). Suppose events A and B are not independent, i.e.,

$$\Pr(A | B) \neq \Pr(A)$$

but they become independent when conditioned with another event C , i.e.,

$$\Pr(A | BC) = \Pr(A | C)$$

Here we say that A and B are **conditionally independent** given C . Furthermore

$$\Pr(AB | C) = \Pr(A | C) \Pr(B | C)$$

Conversely, events A and B may be conditionally dependent but unconditionally independent, i.e.,

$$\begin{aligned} \Pr(A | B) &= \Pr(A) \\ \Pr(A | BC) &\neq \Pr(A | C) \\ \Pr(AB | C) &= \Pr(A | BC) \Pr(B | C) \end{aligned}$$

Theorem 2.2.1. *Given events A , B , and C . Pairwise independence does not imply mutual independence. I.e.,*

$$\begin{cases} \Pr(AB) = \Pr(A) \Pr(B) \\ \Pr(AC) = \Pr(A) \Pr(C) \\ \Pr(BC) = \Pr(B) \Pr(C) \end{cases}$$

does not imply

$$\Pr(ABC) = \Pr(A) \Pr(B) \Pr(C).$$

In summary, independence should not be assumed unless explicitly stated.

2.3 Disjoint Events

Theorem 2.3.1 (Probability of disjoint events). *The probability of disjoint events A and B is given by*

$$\begin{aligned}\Pr(AB) &= 0 \\ \Pr(\emptyset) &= 0.\end{aligned}$$

Disjoint events are highly dependent events, since the occurrence of one means the other cannot occur. This implies

$$\Pr(A|B) = 0$$

2.4 Subsets

Theorem 2.4.1 (Probability of subsets). *If $A \subset B$ then $\Pr(A) \leq \Pr(B)$. We also know that $\Pr(AB) = \Pr(A)$ and $\Pr(A \cup B) = \Pr(B)$. Here, if A happens, then B definitely happens.*

$$\Pr(B|A) = 1$$

Given $\Pr(AB) = \Pr(A)$

$$\Pr(A|B) = \frac{\Pr(A)}{\Pr(B)}$$

These events are also highly dependent.

3 Total Probability

Definition 3.1 (Marginal probability). Marginal probability is the probability of an event irrespective of the outcome of another variable.

Theorem 3.0.1 (Total probability for complements). *By writing the event A as $AB \cup A\bar{B}$, and noting that AB and $A\bar{B}$ are disjoint, the marginal probability of A is given by*

$$\Pr(A) = \Pr(AB) + \Pr(A\bar{B}).$$

By applying the multiplication rule to each joint probability:

$$\Pr(A) = \Pr(A|B)\Pr(B) + \Pr(A|\bar{B})\Pr(\bar{B})$$

Theorem 3.0.2 (Law of total probability). *The previous theorem partitioned Ω into disjoint events B and \bar{B} .*

By partitioning Ω into a collection of disjoint events B_1, B_2, \dots, B_n , such that $\bigcup_{i=1}^n B_i = \Omega$, we have

$$\Pr(A) = \sum_{i=1}^n \Pr(A|B_i)\Pr(B_i)$$

Theorem 3.0.3 (Bayes' Theorem). *Given the probability for A given B , the probability of the reverse direction is given by*

$$\Pr(A|B) = \frac{\Pr(B|A)\Pr(A)}{\Pr(B)}$$

4 Combinatorics

Definition 4.1 (Number of outcomes). Let $|A|$ denote the number of outcomes in an event A .

Theorem 4.0.1 (Addition principle). *Given a sample space S with k disjoint events $\{S_1, \dots, S_k\}$, where the i th event has n_i possible outcomes, the number of possible samples from any event is given by*

$$|\bigcup_{i=1}^k S_i| = \sum_{i=1}^k n_i$$

Theorem 4.0.2 (Multiplication principle). *Given a sample space S with k events $\{S_1, \dots, S_k\}$, where the i th event has n_i possible outcomes, the number of possible samples from every event is given by*

$$|\bigcap_{i=1}^k S_i| = \prod_{i=1}^k n_i$$

Theorem 4.0.3 (Counting probability). *Given a sample space S with equally likely outcomes, the probability of an event $S_i \subset S$ is given by*

$$\Pr(S_i) = \frac{|S_i|}{|S|}$$

4.1 Ordered Sampling with Replacement

When ordering is important and repetition is allowed, the total number of ways to choose k objects from a set with n elements is

$$n^k$$

4.2 Ordered Sampling without Replacement

When ordering is important and repetition is not allowed, the total number of ways to arrange k objects from a set of n elements is known as a k -permutation of n -elements denoted ${}^n P_k$

$$\begin{aligned} {}^n P_k &= n \times (n-1) \times \dots \times (n-k+1) \\ &= \frac{n!}{(n-k)!} \end{aligned}$$

for $0 \leq k \leq n$.

Definition 4.2 (Permutation of n elements). An n -permutation of n elements is the permutation of those elements. In this case, $k = n$, so that

$$\begin{aligned} {}^n P_n &= n \times (n-1) \times \dots \times (n-n+1) \\ &= n! \end{aligned}$$

4.3 Unordered Sampling without Replacement

When ordering is not important and repetition is not allowed, the total number of ways to choose k objects from a set of n elements is known as a k -combination of n -elements denoted nC_k or $\binom{n}{k}$

$$\begin{aligned} {}^nC_k &= \frac{{}^nP_k}{k!} \\ &= \frac{n!}{k!(n-k)!} \end{aligned}$$

for $0 \leq k \leq n$. We divide by $k!$ because any k -element subset of n -elements can be ordered in $k!$ ways.

4.4 Unordered Sampling with Replacement

When ordering is not important and repetition is allowed, the total number of ways to choose k objects from a set with n elements is

$$\binom{n+k-1}{k}$$

5 Random Variables and Distributions

6 Random Variables

Definition 6.1 (Random variable). A random variable X is a measurable variable whose value holds some uncertainty.

An event is when a random variable assumes a certain value or range of values.

Definition 6.2 (Discrete random variables). A discrete random variable takes discrete values.

Definition 6.3 (Continuous random variables). A continuous random variable can take any real value.

6.1 Probability Distributions

Definition 6.4 (Probability distribution). The probability distribution of a random variable X is a function that links all outcomes $x \in \Omega$ to the probability that they will occur $\Pr(X = x)$.

Definition 6.5 (Probability mass function). The probability distribution of a discrete random variable X is described by a Probability Mass Function (PMF) p_x .

$$\Pr(X = x) = p_x$$

p_x is a valid PMF provided,

$$\forall x \in \Omega : \Pr(X = x) \geq 0 \quad \text{and} \quad \sum_{\Omega} \Pr(X = x) = 1.$$

Definition 6.6 (Probability density function). The probability distribution of a continuous random variable X is described by a Probability Density Function (PDF) $f(x)$. The probability that X is exactly equal to a specific value is always 0. Therefore we compute probabilities over intervals:

$$\Pr(x_1 \leq X \leq x_2) = \int_{x_1}^{x_2} f(u) du$$

$f(x)$ is a valid PDF provided,

$$\forall x \in \Omega : f(x) \geq 0 \quad \text{and} \quad \int_{\Omega} f(u) du = 1.$$

Definition 6.7 (Cumulative distribution function). The Cumulative Distribution Function (CDF) computes the probability that the random variable is less than or equal to a particular realisation x . For $U = \{k \in \Omega : k \leq x\}$

$$F(x) = \Pr(X \leq x) = \begin{cases} \sum_{u \in U} p_u & \text{for discrete random variables} \\ \int_U f(u) du & \text{for continuous random variables.} \end{cases}$$

$F(x)$ is a valid CDF if:

1. F is monotonically increasing and continuous
2. $\lim_{x \rightarrow -\infty} F(x) = 0$
3. $\lim_{x \rightarrow \infty} F(x) = 1$

We can recover the PDF given the CDF, by using the Fundamental Theorem of Calculus.

$$\frac{dF(x)}{dx} = \frac{d}{dx} \int_{-\infty}^x f(u) du = f(x)$$

Definition 6.8 (Complementary CDF). For a continuous random variable X the complement function,

$$\Pr(X > x) = 1 - \Pr(X \leq x) = 1 - F(x)$$

is called the complementary CDF, or the survival function.

6.2 Quantiles

Definition 6.9 (p -Quantile). For a continuous random variable, the p -quantile, x , is defined such that

$$F(x) = \int_{-\infty}^x f(u) du = p.$$

Definition 6.10 (Median). The median, m , is a special p -quantile defined as the value such that

$$\int_{-\infty}^m f(u) du = \int_m^{\infty} f(u) du = \frac{1}{2}.$$

Definition 6.11 (Lower and upper quartile). Likewise the lower quartile and upper quartiles are two values q_1 and q_2 such that

$$\int_{-\infty}^{q_1} f(u) du = \frac{1}{4}$$

and

$$\int_{-\infty}^{q_2} f(u) du = \frac{3}{4}.$$

Definition 6.12 (Quantile function). The quantile function is the inverse of the CDF and can be used to find the x that a certain p provides. I.e.,

$$x = F^{-1}(p) = Q(p)$$

6.3 Summary Statistics

Definition 6.13 (Expectation). The expectation $E(X)$, or $\mathbb{E}(X)$ of a random variable X is its expected value given an infinite number of observations.

The expectation is also known as the mean of the X , denoted μ .

$$E(X) = \begin{cases} \sum_{x \in \Omega} x p_x & \text{for discrete variables} \\ \int_{\Omega} x f(x) dx & \text{for continuous variables} \end{cases}$$

Theorem 6.3.1. Using integration by parts, it can be proved that

$$E(X) = - \int_{-\infty}^0 F(x) dx + \int_0^{\infty} (1 - F(x)) dx$$

Definition 6.14 (Variance). The variance $\text{Var}(X)$, or $\mathbb{V}(X)$ of a random variable X is a measure of spread of the distribution (defined as the average squared distance of each value from the mean). Variance is also denoted as σ^2 .

$$\begin{aligned} \text{Var}(X) &= \begin{cases} \sum_{x \in \Omega} (x - \mu)^2 p_x & \text{for discrete variables} \\ \int_{\Omega} (x - \mu)^2 f(x) dx & \text{for continuous variables} \end{cases} \\ &= E(X^2) - E(X)^2 \end{aligned}$$

Definition 6.15 (Standard deviation). The standard deviation is defined as

$$\sigma = \sqrt{\text{Var}(X)}$$

6.3.1 Transformations

For a simple linear function of a random variable

$$\begin{aligned} E(aX \pm b) &= a E(X) \pm b \\ \text{Var}(aX \pm b) &= a^2 \text{Var}(X) \end{aligned}$$

7 Special Discrete Distributions

7.1 Discrete Uniform Distribution

A discrete uniform distribution describes the probability distribution of a single trial in a set of equally likely elements.

A discrete random variable X with a discrete uniform distribution is denoted

$$X \sim \text{Uniform}(a, b)$$

with

$$\begin{aligned}\Pr(X = x) &= \frac{1}{b - a + 1} \\ \Pr(X \leq x) &= \frac{x - a + 1}{b - a + 1}\end{aligned}$$

for outcomes $x \in \{a, a + 1, \dots, b - 1, b\}$. We can also summarise the following:

$$\begin{aligned}\mathbb{E}(X) &= \frac{a + b}{2} \\ \text{Var}(X) &= \frac{(b - a + 1)^2 - 1}{12}\end{aligned}$$

7.2 Bernoulli Distribution

A Bernoulli (or binary) distribution describes the probability distribution of a Boolean-valued outcome, i.e., success (1) or failure (0).

A discrete random variable X with a Bernoulli distribution is denoted

$$X \sim \text{Bernoulli}(p)$$

with

$$\begin{aligned}\Pr(X = x) &= \begin{cases} 1 - p & x = 0 \\ p & x = 1 \end{cases} \\ &= p^x (1 - p)^{1-x} \\ \Pr(X \leq x) &= \begin{cases} 0 & x < 0 \\ 1 - p & 0 \leq x < 1 \\ 1 & k \geq 1 \end{cases}\end{aligned}$$

for a probability $p \in [0, 1]$ and outcomes $x \in \{0, 1\}$. We can also summarise the following:

$$\begin{aligned}\mathbb{E}(X) &= p \\ \text{Var}(X) &= p(1 - p)\end{aligned}$$

where $(1 - p)$ is sometimes denoted as q .

7.3 Binomial Distribution

A binomial distribution describes the probability distribution of the number of successes for n independent trials with the same probability of success p .

A discrete random variable X with a binomial distribution is denoted

$$X \sim B(n, p)$$

with

$$\begin{aligned}\Pr(X = x) &= \binom{n}{x} p^x (1-p)^{n-x} \\ \Pr(X \leq x) &= \sum_{u=0}^x \binom{n}{u} p^u (1-p)^{n-u}\end{aligned}$$

for number of successes $x \in \{0, 1, \dots, n\}$.

Here each individual trial is a Bernoulli trial, so that X can be written as the sum of n *independent and identically distributed* (iid) Bernoulli random variables, Y_1, Y_2, \dots, Y_n .

$$X = Y_1 + Y_2 + \dots + Y_n, \quad Y_i \stackrel{\text{iid}}{\sim} \text{Bernoulli}(p) : \forall i \in \{1, 2, \dots, n\}.$$

We can then summarise the following:

$$\begin{aligned}\mathbb{E}(X) &= np \\ \text{Var}(X) &= np(1-p)\end{aligned}$$

Proof. Given n trials, the probability of x successes will be p^x . Similarly the probability of $n-x$ failures will be $(1-p)^{n-x}$.

We then consider the number of ways to choose x successes out of n trials, i.e., $\binom{n}{x}$.

The intersection of these three events gives the PMF for a binomial distribution. \square

7.4 Geometric Distribution

A geometric distribution describes the probability distribution of the number of trials up to and including the first success, where each trial is independent and has the same probability of success p .

A discrete random variable N with a geometric distribution is denoted

$$N \sim \text{Geom}(p)$$

with

$$\begin{aligned}\Pr(N = n) &= (1-p)^{n-1} p \\ \Pr(N \leq n) &= 1 - (1-p)^n\end{aligned}$$

for number of trials $n \geq 1$.

We can also summarise the following:

$$\begin{aligned}\mathbb{E}(N) &= \frac{1}{p} \\ \text{Var}(N) &= \frac{1-p}{p^2}\end{aligned}$$

7.4.1 Alternate Geometric Definition

We can alternatively consider the number of failures until a success, Y :

$$Y = N - 1$$

Therefore the PMF and CDF for Y are:

$$\Pr(Y = y) = (1 - p)^y p$$

$$\Pr(Y \leq y) = 1 - (1 - p)^{y+1}$$

for number of failures $y \geq 0$. This gives the following summary statistics using transformation rules:

$$\begin{aligned} E(Y) &= \frac{1 - p}{p} \\ \text{Var}(Y) &= \frac{1 - p}{p^2} \end{aligned}$$

7.5 Negative Binomial Distribution

A negative binomial distribution describes the probability distribution of the number of trials until $k \geq 1$ successes, where each trial is independent and has the same probability of success p .

A discrete random variable N with a negative binomial distribution is denoted

$$N \sim \text{NB}(k, p)$$

with

$$\begin{aligned} \Pr(N = n) &= \binom{n-1}{k-1} (1-p)^{n-k} p^k \\ \Pr(N \leq n) &= \sum_{u=k}^n \binom{u-1}{k-1} (1-p)^{u-k} p^k \end{aligned}$$

for number of trials $n \geq k$. Here each individual trial is a Geometric trial, so that N can be written as the sum of k *independent and identically distributed* (iid) Geometric random variables, Y_1, Y_2, \dots, Y_k .

$$N = Y_1 + Y_2 + \dots + Y_k, \quad Y_i \stackrel{\text{iid}}{\sim} \text{Geom}(p) : \forall i \in \{1, 2, \dots, k\}.$$

We can then summarise the following:

$$\begin{aligned} E(N) &= \frac{k}{p} \\ \text{Var}(N) &= \frac{k(1-p)}{p^2} \end{aligned}$$

Proof. Given n trials, the probability of k successes will be p^k . Similarly the probability of $n - k$ failures will be $(1 - p)^{n-k}$.

We then consider the number of ways to arrange $k - 1$ successes for $n - 1$ trials, because the last trial must be a success, i.e., $\binom{n-1}{k-1}$.

The intersection of these three events gives the PMF for a negative binomial distribution. \square

7.5.1 Alternate Negative Binomial Definition

We can alternatively consider the number of failures Y until k successes:

$$Y = N - k$$

The PMF and CDF for Y are given by:

$$\begin{aligned}\Pr(Y = y) &= \binom{y+k-1}{k-1} (1-p)^y p^k \\ \Pr(Y \leq y) &= \sum_{u=0}^y \binom{u+k-1}{k-1} (1-p)^u p^k\end{aligned}$$

for number of failures $y \geq 0$. This gives the following summary statistics using transformation rules:

$$\begin{aligned}\mathbb{E}(Y) &= \frac{k(1-p)}{p} \\ \text{Var}(Y) &= \frac{k(1-p)}{p^2}\end{aligned}$$

7.6 Poisson Distribution

A Poisson distribution describes the probability distribution of the number of events N which occur over a fixed interval of time λ .

A discrete random variable N with a Poisson distribution is denoted

$$N \sim \text{Pois}(\lambda)$$

with

$$\begin{aligned}\Pr(N = n) &= \frac{\lambda^n e^{-\lambda}}{n!} \\ \Pr(N \leq n) &= e^{-\lambda} \sum_{u=0}^n \frac{\lambda^u}{u!}\end{aligned}$$

for number of events $n \geq 0$. We can also summarise the following:

$$\begin{aligned}\mathbb{E}(N) &= \lambda \\ \text{Var}(N) &= \lambda\end{aligned}$$

7.7 Modelling Count Data

If we want to utilise these distributions to model data, we can use the following observations:

- Poisson (mean = variance)
- Binomial (underdispersed, mean > variance)
- Geometric/Negative Binomial (overdispersed, mean < variance)

8 Special Continuous Distributions

8.1 Continuous Uniform Distribution

A continuous uniform distribution describes the probability distribution of an outcome within some interval, where the probability of an outcome in one interval is the same as all other intervals of the same length.

A continuous random variable X with a continuous uniform distribution is denoted

$$X \sim U(a, b)$$

with

$$\begin{aligned} f(x) &= \frac{1}{b-a} \\ F(x) &= \frac{x-a}{b-a} \end{aligned}$$

for outcomes $a < x < b$. We can also summarise the following:

$$\begin{aligned} E(X) &= \frac{a+b}{2} \\ \text{Var}(X) &= \frac{(b-a)^2}{12} \\ m &= \frac{a+b}{2} \end{aligned}$$

8.2 Exponential Distribution

An exponential distribution describes the probability distribution of the time between events with rate η .

A continuous random variable T with an exponential distribution is denoted

$$T \sim \text{Exp}(\eta)$$

with

$$\begin{aligned} f(t) &= \eta e^{-\eta t} \\ F(t) &= 1 - \eta e^{-\eta t} \end{aligned}$$

for time $t > 0$. We can also summarise the following:

$$\begin{aligned} E(X) &= \frac{1}{\eta} \\ \text{Var}(X) &= \frac{1}{\eta^2} \\ m &= \frac{\ln(2)}{\eta} \end{aligned}$$

Proof. By considering an event taking longer than t seconds, we can represent this as nothing happening over the interval $[0, t]$. Using $T \sim \text{Exp}(\eta)$ and $N \sim \text{Pois}(\eta t)$, we have

$$\Pr(T > t) = \Pr(N = 0) = e^{-\eta t}$$

where $\lambda = \eta t$. The CDF for the exponential distribution is then

$$\begin{aligned}\Pr(T < t) &= 1 - \Pr(T > t) \\ &= 1 - e^{-\eta t}.\end{aligned}$$

□

8.3 Memoryless Property

In an exponential distribution with $T \sim \text{Exp}(\eta)$, the distribution of the waiting time $t + s$ until a certain event does not depend on how much time t has already passed.

$$\Pr(T > s + t \mid T > t) = \Pr(T > s).$$

The same property also applies in an Geometric distribution with $N \sim \text{Geom}(p)$.

8.4 Normal Distribution

The normal distribution is used to represent many random situations, in particular, measurements and their errors. This distribution arises in many statistical problems and can be used to approximate other distributions under certain conditions.

A continuous random variable X with a normal distribution is denoted

$$X \sim N(\mu, \sigma^2)$$

with

$$\begin{aligned}f(t) &= \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}} \\ F(t) &= \frac{1}{2} \left(1 + \text{erf}\left(\frac{x-\mu}{\sigma\sqrt{2}}\right) \right)\end{aligned}$$

for $x \in \mathbb{R}$ where $\text{erf}(z) = \frac{2}{\sqrt{\pi}} \int_0^z e^{-t^2} dt$ is the error function. We can also summarise the following:

$$\begin{aligned}\mathbb{E}(X) &= \mu \\ \text{Var}(X) &= \sigma^2\end{aligned}$$

Given the complexity of the analytic expressions for the PDF and CDF of the normal distribution, we often use software to numerically determine probabilities associated with normal distributions.

8.5 Standard Normal Distribution

Given $X \sim N(\mu, \sigma^2)$, consider the transformation

$$Z = \frac{X - \mu}{\sigma}$$

so that $Z \sim N(0, 1)$. This distribution is called the standard normal distribution. This allows us to deal with the standard normal distribution regardless of μ and σ .