

Written HW1

Logan Tanner
I use Neovim (BTW)

May 15, 2026

- Suppose that we have 10 coins, which are weighted so that when flipped the i th coin shows heads with probability $p = i/10$ ($i = 1, \dots, 10$).
Setup:

$$P(C_i) = \text{probability of the } i^{\text{th}} \text{ coin being flipped} = \frac{1}{10}$$

$$P(H) = \text{probability of a Head being flipped}$$

- If we randomly select a coin, flip it, and get heads, what is the probability that it is the 3rd coin?

Answer:

$$P(C_3|H) = \frac{P(C_3 \cap H)}{P(H)}$$

$$P(H) = \left(\sum_{i=1}^{10} \frac{i}{10}\right) \times \frac{1}{10} = \frac{55}{100}$$

$$P(C_3 \cap H) = P(C_3) \times P(H|C_3) = \frac{1}{10} \times \frac{3}{10} = \frac{3}{100}$$

$$P(C_3|H) = \frac{3}{100} \times \frac{100}{55} = \boxed{\frac{3}{55}}$$

- What is the probability that it is the i th coin?

$$P(C_i|H) = \frac{P(C_i \cap H)}{P(H)}$$

$$P(C_i \cap H) = P(C_i) \times P(H|C_i) = \frac{1}{10} \times \frac{i}{10} = \frac{i}{100}$$

$$P(C_i|H) = \frac{i}{100} \times \frac{100}{55} = \boxed{\frac{i}{55}}$$

2. It is important to consider blood type for both the recipient and donor for any blood transfusion: AB recipients can receive blood of any type, and type A or B recipients can receive blood of their own type or type O, but type O can only receive O type blood. Given that the percentages of type O, A, B, and AB in the population are 34%, 38%, 20%, and 8% respectively, find the probability that a random donor can give blood to a random recipient.

3. Let E_1 , E_2 , and F be events, with $P(F) > 0$. Use the definition of conditional probability along with other results from class to show that Property 3 also holds for conditional probabilities. That is, show that

$$P(E_1 \cup E_2|F) = P(E_1|F) + P(E_2|F) - P(E_1 \cap E_2|F)$$

4. Consider rolling two fair dice repeatedly either a roll sums to 6, or a 2 is rolled (on one or both dice). In class, we computed that the probability that a roll sums to 6 strictly before any 2s are rolled is $\frac{3}{14}$.
- (a) Find the probability that a 2 is rolled strictly before a roll sums to 6.
 - (b) What is the third possibility? What is its probability?

5. You're on a game show! There are three doors labeled A , B , and C . A new car is behind one of the three doors, but you don't know which one. You select one of the door, say, door A . The host then opens one of doors B or C , as follows: if the car is behind door B , then he opens door C ; if the car is behind door C , then he opens door B ; and if the car is behind door A , then he opens either door B or C with probability $\frac{1}{2}$ each. (In all cases, the door opened by the host will not have the car behind it.)

The host then gives you the choice of either sticking with door A , or switching to the remaining unopened door. You win the car if and only if it is behind your final door selection. Without loss of generality, we assume that the host opens door B .

- (a) If you stick with your original choice (i.e., door A), conditioned on the host having opened door B , then what is your probability of winning?
- (b) If you switched to the remaining door (i.e., door C), conditioned on the host having opened door B , then what is your probability of winning?
- (c) Supposed we change the rules so that, if you originally chose door A and the car was indeed behind door A , then the host always opens door B . Do the answers to questions 1 and 2 change?