

Superconductivity in 2D Multicomponents Materials

Time-reversal symmetry breaking in 2D multi-components material
using Monte Carlo simulation

Logan Peschieras

Department of Physics
Stockholm University

Supervisor: Johan Carlström Email: johan.carlstrom@fysik.su.se

Contact Person: Vladimir Krasnov Email: vladimir.krasnov@fysik.su.se



Stockholm
University


Table of Contents

- 1 Theory in superconductivity
 - Theory in superconductivity
 - Cooper pairs
 - Moiré materials
 - Josephson coupling in the London limit
 - Time-reversal symmetry
 - Phase stiffness(2D), $U(1)$ symmetry(3D)
- 2 Statistical Mechanics, Monte Carlo Simulation, Metropolis Algorithm
 - Statistical Mechanics
 - Monte Carlo Simulation
 - Metropolis Algorithm
- 3 Results: The program
 - The code
 - Results
 - Flaws, error and accuracy
 - Interpretation
- 4 References



Theory in superconductivity

- Ohm's law : $R = \frac{U}{I}$
- Superconductivity (1911) $\rightarrow$ no resistance.
- Duality wave-particle in metals by Bloch (1930):





Cooper pairs

- Order Parameter:


$$O_s = \int \frac{\Psi}{A} dx$$

Cooper pair formation



Moiré materials

- Twisting of two 2D graphene from one to another:



Josephson coupling in the London limit

- Ginzburg-Landau free energy:

$$f = \alpha|\Psi|^2 + \frac{\beta}{2}|\Psi|^4 + \frac{1}{2m}|(-i\hbar\nabla - 2e\vec{A})\Psi|^2 + \frac{\mu_0 B^2}{2}$$

- But in 2D:

$$f = \alpha|\Psi|^2 + \frac{\beta}{2}|\Psi|^4 + \frac{1}{2m}|\nabla\Psi|^2$$


- In our scenario:

$$f = \alpha_i|\Psi_i|^2 + \frac{\beta_i}{2}|\Psi_i|^4 + \frac{1}{2m}|\nabla\Psi_i|^2 + \epsilon_{ij}\Psi_i\Psi_j^* + \text{constant}$$



Josephson coupling in the London limit

System:



Each superconductivity component forms on a Fermi surface. p_i belongs to Fermi surface i .


Time-reversal symmetry

- Time reversal operator T : $T\Psi(x, t) = \Psi^*(x, -t)$
- $T = UK$, U = unitary operator, K = complex conjugate operator
- Kramer's theorem: $T^2 = \pm 1$
- $TpT^{-1} = -p$ (=anti-commute)
- $TJT^{-1} = -J$
- $TxT^{-1} = x$ (=commute)
- GL-theory $\rightarrow$ non vanishing order parameter $S \rightarrow$ broken time-reversal symmetry $TS = -S$




Phase stiffness(2D), $U(1)$ symmetry(3D)

- Phase stiffness $\rightarrow$ energy required to change the phase order $\rightarrow$ lattice distortion
- $U(1)$ symmetry $\rightarrow$ gauge symmetry conserving laws of physics under transformations to fields and particle
- Cooper pairs $\rightarrow U(1)$ symmetry spontaneously broken.



- Boltzmann factor:

$$P(E_i) = \sum_i e^{-\beta H_i}; \beta = \frac{1}{kT}$$



- Hamiltonian:

$$H = - \sum_{A, \langle i, j \rangle} \cos(\varphi_{A,i} - \varphi_{A,j}) + \eta \sum_{i, A \neq B} \cos(\varphi_{A,i} - \varphi_{B,i})$$



Monte Carlo Simulation

- (1)-Input values
- (2)-New example sample of values
- (3)-Mathematical model
- (4)-Output the accepted values
- (5)-Analysis of the data given



Metropolis Algorithm

- Generating/ Selecting an initial value θ_0
- The number of simulations = $n \rightarrow$ we repeat n times:
 - New value proposed θ^* from proposal probability distribution $p(\theta^*|\theta_n)$
 - Rate: $r = \frac{p(\theta^*)p(\theta_n|\theta^*)}{p(\theta_n)p(\theta^*|\theta_n)}$
 - If $r > 1$: $\theta_n = \theta^*$
 - If $1 > r > 0$: $\theta_n = \theta^*$ is r probable and $\theta_n = \theta_0$ is $1 - r$ probable.
 - $\theta_n =$ new initial values $\rightarrow$ compared with a new proposed value in next simulation.
- Probability density symmetrical $\rightarrow \frac{p(\theta_n|\theta^*)}{p(\theta^*|\theta_n)} = 1 \rightarrow r = \frac{p(\theta^*)}{p(\theta_n)}$



The code

- Boltzmann factor/ Probability of the Hamiltonian($k = 1$):

$$\text{prob}(H) = e^{-\frac{H}{T}}$$

- Ratio:



$$r = \frac{\text{prob}(H_2)}{\text{prob}(H_1)}$$

- if p_1, p_2, p_3 in one order $\rightarrow$ phase order "up".
- Else, phase order "down".

Example of phase order "up" and "down"




- Average order parameter $\langle |S| \rangle$ related to time-reversal symmetry in function of the temperature:



Results

- Order parameter S_i related to time-reversal symmetry for a lattice 5x5 at $T=0.5\text{K}$ and $T=10\text{K}$:



Flaws, error and accuracy

- Standard deviation

$$\sigma = \sqrt{\frac{\langle |S|^2 \rangle - \langle |S| \rangle^2}{N}}$$

- with

$$\langle |S| \rangle = \frac{1}{N} \sum_{n=1}^N |S_n|$$


and

$$\langle |S|^2 \rangle = \frac{1}{N} \sum_{n=1}^N |S_n|^2$$



Flaws, error and accuracy

- Standard deviation over the total number of simulation N :



Flaws, error and accuracy


- Correlation function between the simulations:

$$c(l) = \sqrt{\frac{1}{N-l} \sum_{n=1}^{N-l} (S_n - \langle S \rangle)(S_{n+l} - \langle S \rangle)}$$

- When $\langle S \rangle = 0$,


$$c(l) = \sqrt{\frac{1}{N-l} \sum_{n=1}^{N-l} S_n S_{n+l}}$$

- Correlation between n -th and $n+1$ -th value over the simulation number n :




Interpretation

- Separation of the normal time-reversal symmetry and spontaneously broken time-reversal symmetry state:



Conclusion

- Picture of a superfluid liquid helium:



References

- V. Schmidt, P. Müller, and A. Ustinov, The Physics of Superconductors, 1st ed. Tiergartenstraße 17, 69121 Heidelberg, Germany: Springer-Verlag Berlin Heidelberg GmbH, 1997.
- J. Bobroff, "Quelle est la tête d'un electron dans un metal," Instagram- Julien Bobroff, 2023. W. Bao-Tian, L. Peng-Fei, B. Tao, Y. Wen, E. Olle, Z. Jijun, and W. Fangwei, "Superconductivity in two-dimensional phosphorus carbide (-pc)," Physical Chemistry Chemical Physics, Royal Society of Chemistry, 2018.
- D. Griffiths and D. Schroeter, Introduction to Quantum Mechanics, 3rd ed. The Old Schools, Trinity Ln, Cambridge CB2 1TN, Great Britain: Cambridge University Press, 2018.
- J. Anandan, "Causality, symmetries and quantum mechanics," Foundations of Physics Letters, 2002.
- "T-symmetry," Wikipedia, 2023. . Zaki and P. Johnson, "Time-reversal symmetry breaking in a superconductor," SciTechDaily, 2021.
- "Spin correlation in cooper pair demonstrated," Quantum Grad, 2022. I. Mangel, I. Kapon, N. Blau, K. Golubkov, N. Gavish, and A. Keren, "Stiffnessometer: A magnetic-field-free superconducting stiffness meter and its application," Physical Review covering condensed matter and materials physics, 2020.
- IvanaG, "Superconductor symmetry breaking," StackExchange, 2017. D. V. Schroeder, An Introduction to Thermal Physics, 1st ed. Addison Wesley Longman: Addison Wesley, 1999.
- Amazon-Web-services, "What is monte carlo simulation," AWS HPC Blog, 2023.
- B. Lambert, A Student's Guide to Bayesian Statistics, 1st ed. Thousand Oaks, California, United-States: SAGE Publications Ltd, 2007.
- M. Allen and D. Tildesley, Computer Simulation of Liquids, 1st ed. Oxford, Great- Britain: Oxford Science Publications, 2007.
- Unknown, "Error analysis in monte carlo," UC Davis, California's College Town.
- L. Burrows, "Taking the guesswork out of twistrionics," Harvard John A. Paulson School of Engineering and Applied Sciences, 2020.
- "Helium liquid," Wikipedia.

