
Row-normalised |interaction| dot + outline = FDR q ≤ 0·10
0·0
0·5
1·0
a
|SHAP interaction| per biomarker × modifier category
Diastolic BP
Heart rate
Creatinine
Haematocrit
Haemoglobin
CD4 count
HIV viral load
LDL-cholesterol
Body fat %
Waist-hip ratio
Age
Sex
HIV status
SES
Temp×Temp
b
FDR-significant interactions per biomarker
FDR-significant interactions (n)
c
Modifier-category composition of FDR-significant interactions
Age 2 (8%)
Sex 8 (32%)
HIV status 0 (0%)
SES 2 (8%)
Temp×Temp 13 (52%)
FDR-significant interactions (n)
d
Ten strongest FDR-significant interactions
Creatinine · Male (vs F) × 3d
|SHAP|=0·127, q=0·032
Haematocrit · Male (vs F) × 3d
|SHAP|=0·039, q=0·026
LDL-cholesterol · temp lag 30d × 14d
|SHAP|=0·008, q=0·072
Body fat % · temp lag 30d × 14d
|SHAP|=0·086, q=0·033
Waist-hip ratio · Male (vs F) × 21d
|SHAP|=0·002, q=0·008
Haematocrit · Male (vs F) × 21d
|SHAP|=0·038, q=0·026
LDL-cholesterol · year × 21d
|SHAP|=0·007, q=0·072
Waist-hip ratio · age years × 21d
|SHAP|=0·002, q=0·008
Creatinine · Male (vs F) × 30d
|SHAP|=0·108, q=0·032
Body fat % · temp lag 3d × 1d
|SHAP|=0·070, q=0·066
Scaled interaction magnitude
(within-biomarker)