
* ρ ≥ 0·70 counted as concordant (bold outline in panel a).




a

Spearman ρ of SHAP lag profile vs primary specification

	SA4 p95	SA4 p90	SA5 apparent T	SA5 heat index	SA5 wet bulb
Diastolic BP	0·11	0·68	0·21	0·11	0·54
Heart rate	0·82	0·82	0·75	0·89	0·86
Creatinine	0·29	0·39	0·21	0·79	0·57
Haematocrit	0·68	0·82	0·50	0·79	0·21
Haemoglobin	0·96	0·89	0·75	0·79	0·86
CD4 count	0·75	0·96	0·36	−0·43	0·36
HIV viral load	0·32	0·54	0·14	0·04	−0·11
LDL-cholesterol	0·89	0·86	0·75	0·50	0·68
Body fat %	1·00	0·89	0·61	0·36	0·54
Waist-hip ratio	0·96	0·96	0·14	−0·14	0·21


b

Sensitivity ranking



c

Exposure-metric sensitivity (SA5) – dry-bulb reference vs humidity-aware indices



d

SA3 | Lag-profile agreement by HIV status

7 of 8

comparable biomarkers had ρ < 0·70 between HIV-positive and HIV-negative lag profiles

Lower agreement (ρ < 0·70):

- Body fat %
- Waist-hip ratio
- Creatinine
- Diastolic BP
- Haematocrit
- Haemoglobin
- LDL-cholesterol

Concordant (ρ ≥ 0·70):

- Heart rate

No HIV-negative comparator: CD4 count, HIV viral load