

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/314870040>

Prices and Asymptotics for Discrete Variance Swaps

Article in SSRN Electronic Journal · January 2012

DOI: 10.2139/ssrn.2115881

CITATIONS

0

READS

47

2 authors:



Carole Bernard

University of Waterloo

142 PUBLICATIONS 2,307 CITATIONS

[SEE PROFILE](#)



Zhenyu Cui

Stevens Institute of Technology

212 PUBLICATIONS 1,454 CITATIONS

[SEE PROFILE](#)

Prices and Asymptotics for Discrete Variance Swaps

CAROLE BERNARD ^{*}and ZHENYU CUI ^{†‡}

Draft: May 30, 2013

Abstract

We study the fair strike of a discrete variance swap for a general time-homogeneous stochastic volatility model. In the special cases of Heston, Hull-White and Schöbel-Zhu stochastic volatility models we give simple explicit expressions (improving Broadie and Jain (2008a) in the case of the Heston model). We give conditions on parameters under which the fair strike of a discrete variance swap is higher or lower than that of the continuous variance swap. The interest rate and the correlation between the underlying price and its volatility are key elements in this analysis. We derive asymptotics for the discrete variance swaps and compare our results with those of Broadie and Jain (2008a), Jarrow et al. (2013) and Keller-Ressel and Griessler (2012).

Key-words: Discrete Variance swap, Heston model, Hull-White model, Schöbel-Zhu model.

^{*}C. Bernard is with the department of Statistics and Actuarial Science at the University of Waterloo, Email: c3bernar@uwaterloo.ca. C. Bernard acknowledges support from the Natural Sciences and Engineering Research Council of Canada.

[†]Corresponding author. Zhenyu Cui is a Ph.D. candidate in Statistics at the University of Waterloo, Email: cuiizhyu@gmail.com. Z. Cui acknowledges support from the Bank of Montreal Capital Markets Advanced Research Scholarship.

[‡]Both authors thank Jinyuan Zhang for her help as research assistant, as well as seminar participants Christa Cuchiero, Olympia Hadjiladis, Antoine Jacquier, Adam Kolkiewicz, Roger Lee, Don McLeish, Johannes Ruf, and David Saunders for helpful suggestions. We are particularly grateful to an anonymous referee for his/her constructive and helpful comments on an earlier draft of this paper.

Prices and Asymptotics for Discrete Variance Swaps

1 Introduction

A variance swap is a derivative contract that pays at a fixed maturity T the difference between a given level (fixed leg) and a realized level of variance over the swap's life (floating leg). Nowadays, variance swaps on stock indices are broadly used and highly liquid. Less standardized variance swaps could be linked to other types of underlying stocks such as currencies or commodities. They can be useful for hedging volatility risk exposure or for taking positions on future realized volatility. For example, Carr and Lee (2007) price options on realized variance and realized volatility by using variance swaps as pricing and hedging instruments. See Carr and Lee (2009) for a history of volatility derivatives. As noted by Jarrow et al. (2013), most academic studies¹ focus on continuously sampled variance and volatility swaps. However, existing volatility derivatives tend to be based on the realized variance computed from the discretely sampled log stock price and continuously sampled derivatives prices may only be used as approximations. As pointed out in Sepp (2012), some care is needed to replace the discrete realized variance by the continuous quadratic variation. By standard probability arguments, the discretely sampled realized variance converges to the quadratic variation of the log stock price in probability. However, this does not guarantee that it converges in expectation. Jarrow et al. (2013) provide sufficient conditions such that the convergence in expectation happens when the stock is modeled by a general semi-martingale, and concrete examples where this convergence fails.

In this paper we study discretely sampled variance swaps in a general time-homogeneous model for stochastic volatility. For discretely sampled variance swaps, it is difficult to use the elegant and model-free approach of

¹See, for example, Howison, Rafailidis and Rasmussen (2004), Windcliff, Forsyth and Vetzal (2006), Benth, Groth and Kufakunesu (2007) and Broadie and Jain (2008b).

Dupire (1993) and Neuberger (1994), who independently proved that the fair strike for a continuously sampled variance swap on any underlying price process with continuous path is simply two units of the forward price of the log contract. Building on these results, Carr and Madan (1998) published an explicit expression to obtain this forward price from option prices (by synthesizing a forward contract with vanilla options). The Dupire-Neuberger theory was recently extended by Carr, Lee and Wu (2012) to the case when the underlying stock price is driven by a time-changed Lévy process (thus allowing jumps in the path of the underlying stock price). In this paper, we adopt a parametric approach that allows us to derive explicit closed-form expressions and asymptotic behaviors with respect to key parameters such as the maturity of the contract, the risk-free rate, the sampling frequency, the volatility of the variance process, or the correlation between the underlying stock and its volatility. This is in line with the work of Broadie and Jain (2008a) in which the Heston model and the Merton jump diffusion model are considered. See also Itkin and Carr (2010) who study discretely sampled variance swaps in the $3/2$ stochastic volatility model.

Our main contributions are as follows. We give an expression of the fair strike of the discretely sampled variance swap and derive its sensitivity to the interest rate in a general time-homogeneous stochastic volatility model. In the case of the (correlated) Heston (1993) model, the (correlated) Hull-White (1987) model, and the (correlated) Schöbel-Zhu (1999) model, we obtain simple explicit closed-form formulas for the respective fair strikes of continuously and discretely sampled variance swaps. In the Heston model, our formula simplifies the results of Broadie and Jain (2008a) and is easy to analyze. Consequently, we are able to give asymptotic behaviors with respect to key parameters of the model and to the sampling frequency. In particular, we provide explicit conditions under which the fair strike of the discretely sampled variance swap is less valuable than that of the continuously sampled variance swap for high sampling frequencies, although the contrary is commonly observed in the literature (see Bühler (2006) for example). Thus the “convex-order conjecture” formulated by Keller-Ressel and Griessler (2012) may not hold for stochastic volatility models with correlation. We discuss practical implications and illustrate the risk to underestimate or overestimate prices of discretely sampled variance swaps when using a model for the corresponding continuously sampled ones with numerical examples.

The paper is organized as follows. Section 2 deals with the general time-homogeneous stochastic volatility model. Sections 3, 4 and 5 provide formulas for the fair strike of a discrete variance swap in the Heston, the Hull-White and the Schöbel-Zhu models. Section 6 contains asymptotics for the Heston, the Hull-White and the Schöbel-Zhu models and discusses the “convex-order conjecture”. A numerical analysis is given in Section 7.

2 Pricing Discrete Variance Swaps in a Time-homogeneous Stochastic Volatility Model

In this section, we consider the problem of pricing a discrete variance swap under the following general time-homogeneous stochastic volatility model (M) , where the stock price and its volatility can possibly be correlated. We assume a constant risk-free rate $r \geq 0$, and that under a risk-neutral probability measure \mathbb{Q}

$$(M) \quad \begin{cases} \frac{dS_t}{S_t} &= rdt + m(V_t)dW_t^{(1)} \\ dV_t &= \mu(V_t)dt + \sigma(V_t)dW_t^{(2)} \end{cases} \quad (1)$$

where $\mathbb{E}[dW_t^{(1)}dW_t^{(2)}] = \rho dt$, with $W^{(1)}, W^{(2)}$ standard correlated Brownian motions. The state space of the stochastic variance process V is $J = (l, r)$, $-\infty \leq l < r \leq \infty$. Assume that $\mu, \sigma : J \rightarrow \mathbb{R}$ are Borel functions satisfying the following Engelbert-Schmidt conditions, $\forall x \in J, \sigma(x) \neq 0$, $\frac{1}{\sigma^2(x)}, \frac{\mu(x)}{\sigma^2(x)}, \frac{m^2(x)}{\sigma^2(x)} \in L_{loc}^1(J)$. Here $L_{loc}^1(J)$ denotes the class of locally integrable functions, i.e. the functions $J \rightarrow \mathbb{R}$ that are integrable on compact subsets of J . Under the above conditions, the SDE (1) for V has a unique in law weak solution that possibly exits its state space J (see Theorem 5.5.15, page 341, Karatzas and Shreve (1991)). Assume that $\frac{m(x)}{\sigma(x)}$ is differentiable at all $x \in J$.

In particular, this general model includes the Heston, the Hull-White, the Schöbel-Zhu, the 3/2 and the Stein-Stein models as special cases. In what follows, we study discretely and continuously sampled variance swaps with maturity T . In a variance swap, one counterparty agrees to pay at a fixed maturity T a notional amount times the difference between a fixed level and a realized level of variance over the swap’s life. If it is continuously sampled, the realized variance corresponds to the quadratic variation of the

underlying log price. When it is discretely sampled, it is the sum of the squared increments of the log price. Define their respective fair strikes as follows.

Definition 2.1. *The fair strike of the discrete variance swap associated with the partition $0 = t_0 < t_1 < \dots < t_n = T$ of the time interval $[0, T]$ is defined as*

$$K_d^M(n) := \frac{1}{T} \sum_{i=0}^{n-1} \mathbb{E} \left[\left(\ln \frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right], \quad (2)$$

where the underlying stock price S follows the time-homogeneous stochastic volatility model (1) and where the exponent M refers to the model (M) .

Definition 2.2. *The fair strike of the continuous variance swap is defined as*

$$K_c^M := \frac{1}{T} \mathbb{E} \left[\int_0^T m^2(V_s) ds \right], \quad (3)$$

where S follows the time-homogeneous stochastic volatility model (1).

In popular stochastic volatility models, $m(v) = \sqrt{v}$, so that $K_c^M = \frac{1}{T} \mathbb{E} \left[\int_0^T V_s ds \right]$. The derivation of the fair strike of a discrete variance swap in the time-homogeneous stochastic volatility model (1) is based on the following proposition.

Proposition 2.1. *Under the dynamics (1) for the stochastic volatility model (M) , define*

$$f(v) = \int_0^v \frac{m(z)}{\sigma(z)} dz \quad \text{and} \quad h(v) = \mu(v)f'(v) + \frac{1}{2}\sigma^2(v)f''(v).$$

For all $t \leq s \leq t + \Delta$ and $t \leq u \leq t + \Delta$, assume that²

$$\begin{aligned} \mathbb{E} [|h(V_s)h(V_u)|] &< \infty, \quad \mathbb{E} [|h(V_s)m^2(V_u)|] < \infty, \\ \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))(2\rho h(V_s) + m^2(V_s))] &< \infty, \end{aligned} \quad (4)$$

²These conditions ensure that we can apply Fubini's theorem to exchange the order of integration. They are easily verified in specific examples.

Define for $t \leq s \leq t + \Delta$, $t \leq u \leq t + \Delta$,

$$\begin{aligned} m_1(s) &:= \mathbb{E} [m^2(V_s)], & m_2(s, u) &:= \mathbb{E} [m^2(V_s)m^2(V_u)], \\ m_3(s, u) &:= \mathbb{E} [h(V_s)h(V_u)], & m_4(s, u) &:= \mathbb{E} [h(V_s)m^2(V_u)], \\ m_5(t, s) &:= \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))(2\rho h(V_s) + m^2(V_s))], \text{ then} \end{aligned}$$

$$\begin{aligned} \mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] &= r^2 \Delta^2 + (1 - \rho^2 - r\Delta) \int_t^{t+\Delta} m_1(s) ds - \rho \int_t^{t+\Delta} m_5(t, s) ds \\ &+ \frac{1}{4} \int_t^{t+\Delta} \int_t^{t+\Delta} m_2(s, u) ds du + \rho^2 \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))^2] \\ &+ \rho^2 \int_t^{t+\Delta} \int_t^{t+\Delta} m_3(s, u) ds du + \rho \int_t^{t+\Delta} \int_t^{t+\Delta} m_4(s, u) ds du. \end{aligned} \quad (5)$$

Proof. See Appendix A. \square

Proposition 2.1 gives the key equation in the analysis of discrete variance swaps. Observe³ that the final expression (5) only depends on covariances of functionals of V_t . Thus we can derive closed-form formulas for the fair strike of discrete variance swaps in those stochastic volatility models in which the terms m_i from Proposition 2.1 can be computed in closed-form. In the rest of the paper, we provide three examples to apply this formula.

From now on, for simplicity, we consider the equi-distant sampling scheme in (2). Under this scheme, $t_i = iT/n$ and $\Delta = t_{i+1} - t_i = T/n$, for $i = 0, 1, \dots, n$.

Remark 2.1. From (5) it is clear that the fair strike of a discrete variance swap only depends on the risk-free rate r up to the second order, as there is no higher order terms of r . Interestingly, the second order coefficient of this expansion is model-independent whereas the first order coefficient is directly related to the strike of the corresponding continuously-sampled variance swap. Assume a constant sampling period $\frac{T}{n}$, the fair strike of the discrete variance swap can be expressed as

$$K_d^M(n) = b^M(n) - \frac{T}{n} K_c^M r + \frac{T}{n} r^2, \quad (6)$$

where $b^M(n)$ does not depend on r . Its sensitivity⁴ to the risk-free rate r is

³Thanks to the anonymous referee for pointing out this general expression.

⁴The impact of stochastic interest rates on variance swaps is studied by Hörfelt and Torné (2010). Long-dated variance swaps will usually be sensitive to the interest rate

equal to

$$\frac{dK_d^M(n)}{dr} = \frac{T}{n}(2r - K_c^M). \quad (7)$$

so that the minimum of K_d^M as a function of r is attained when the risk-free rate takes the value r^* given by

$$r^* = \frac{K_c^M}{2}.$$

The next proposition deals with the special case when the risk-free rate r and the correlation coefficient ρ are both equal to 0.

Proposition 2.2. *(Fair strike when $r = 0\%$ and $\rho = 0$)*

In the special case when the constant risk-free rate is 0, and the underlying stock price is not correlated to its volatility, we observe that

$$K_d^M(n) \geq K_c^M.$$

Proof. Using Proposition 2.1 when $r = 0\%$ and $\rho = 0$, we obtain

$$\mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] = \frac{1}{4} \mathbb{E} \left[\left(\int_t^{t+\Delta} m^2(V_s) ds \right)^2 \right] + \int_t^{t+\Delta} \mathbb{E} [m^2(V_s)] ds.$$

We then add up the expectations of the squares of the log increments (as in (2)) and find that the fair strike of the discrete variance swap is always larger than the fair strike of the continuous variance swap given in (3). \square

Proposition 2.2 has already appeared in the literature in specific models. See for example Corollary 6.2 of Carr, Lee and Wu (2012), where this result is proved in the more general setting of time-changed Lévy processes with independent time changes. However, we will see in the remainder of this paper that Proposition 2.2 may not hold under more general assumptions, namely when the dynamic of the stock price is correlated to the one of the volatility.

volatility.

3 Fair Strike of the Discrete Variance Swap in the Heston model

Assume that we work under the Heston stochastic volatility model with the following dynamics

$$(H) \quad \begin{cases} \frac{dS_t}{S_t} = rdt + \sqrt{V_t}dW_t^{(1)}, \\ dV_t = \kappa(\theta - V_t)dt + \gamma\sqrt{V_t}dW_t^{(2)} \end{cases} \quad (8)$$

where $\mathbb{E} [dW_t^{(1)}dW_t^{(2)}] = \rho dt$. It is a special case of the general model (1), where we choose

$$m(x) = \sqrt{x}, \quad \mu(x) = \kappa(\theta - x), \quad \sigma(x) = \gamma\sqrt{x}. \quad (9)$$

Using (27) in Lemma A.1 in the Appendix with $f(v) = \frac{v}{\gamma}$ and $h(v) = \frac{\kappa}{\gamma}(\theta - v)$, the stock price is

$$S_t = S_0 e^{rt - \frac{1}{2}\xi_t + (V_t - V_0 - \kappa\theta t + \kappa\xi_t)\frac{\rho}{\gamma} + \sqrt{1-\rho^2} \int_0^t \sqrt{V_s}dW_s^{(3)}} \quad (10)$$

where $\xi_t = \int_0^t V_s ds$ and $W_t^{(3)}$ is such that $dW_t^{(1)} = \rho dW_t^{(2)} + \sqrt{1-\rho^2}dW_t^{(3)}$.

Using Proposition 2.1 for the time-homogeneous stochastic volatility model, we then derive a closed-form expression for the fair strike of a discrete variance swap and compare it with the fair strike of a continuous variance swap.

Proposition 3.1. *(Fair Strikes in the Heston Model)*

In the Heston stochastic volatility model (8), the fair strike (2) of the discrete variance swap is

$$\begin{aligned} K_d^H(n) = \frac{1}{8n\kappa^3T} & \left\{ n \left(\gamma^2 (\theta - 2V_0) + 2\kappa (V_0 - \theta)^2 \right) (e^{-2\kappa T} - 1) \frac{1 - e^{-\frac{\kappa T}{n}}}{1 + e^{-\frac{\kappa T}{n}}} \right. \\ & + 2\kappa T \left(\kappa^2 T (\theta - 2r)^2 + n\theta (4\kappa^2 - 4\rho\kappa\gamma + \gamma^2) \right) \\ & + 4(V_0 - \theta) \left(n(2\kappa^2 + \gamma^2 - 2\rho\kappa\gamma) + \kappa^2 T (\theta - 2r) \right) (1 - e^{-\kappa T}) \\ & \left. - 2n^2\theta\gamma (\gamma - 4\rho\kappa) \left(1 - e^{-\frac{\kappa T}{n}} \right) + 4(V_0 - \theta) \kappa T \gamma (\gamma - 2\rho\kappa) \frac{1 - e^{-\kappa T}}{1 - e^{-\frac{\kappa T}{n}}} \right\}, \end{aligned} \quad (11)$$

where $a = r - \frac{\rho\kappa\theta}{\gamma}$ and $b = \frac{\rho\kappa}{\gamma} - \frac{1}{2}$. The fair strike of the continuous variance

swap is

$$K_c^H = \frac{1}{T} \mathbb{E} \left[\int_0^T V_s ds \right] = \theta + (1 - e^{-\kappa T}) \frac{V_0 - \theta}{\kappa T}. \quad (12)$$

Proof. See Appendix B for the proof of (11). The formula (12) for the fair strike of a continuous variance swap is already well-known and can be found for example in Broadie and Jain (2008a), formula (4.3), page 772. \square

Proposition 3.1 provides an explicit formula for the fair strike of a discrete variance swap as a function of model parameters. This formula simplifies the expressions obtained by Broadie and Jain (2008a) in equations (A-29) and (A-30), page 793, where several sums from 0 to n are involved and can actually be computed explicitly as shown by the expression (11) above. We verified that our formula agrees with numerical examples presented in Table 5 (column ‘SV’) on page 782 of Broadie and Jain (2008a).⁵

Contrary to what is stated in the introduction of the paper by Zhu and Lian (2011), the techniques of Broadie and Jain (2008a) can easily be extended to other types of payoffs. The following proposition gives explicit expressions for the volatility derivative considered by Zhu and Lian (2011).

Proposition 3.2. *For the following set of dates $t_i = \frac{iT}{n}$ with $i = 0, 1, \dots, n$, denote $\Delta = T/n$, and assume $\alpha = 2\kappa\theta/\gamma^2 - 1 \geq 0$, and $\gamma^2 T < 1$. Then the fair price of a discrete variance swap with payoff $\frac{1}{T} \sum_{i=0}^{n-1} \left(\frac{S_{t_{i+1}} - S_{t_i}}{S_{t_i}} \right)^2$ is equal to*

$$K_d^{zl}(n) = \frac{1}{T} \sum_{i=0}^{n-1} \mathbb{E} \left[\left(\frac{S_{t_{i+1}} - S_{t_i}}{S_{t_i}} \right)^2 \right] = \frac{1}{T} \left(a_0 + \sum_{i=1}^{n-1} a_i \right) + \frac{n - 2ne^{r\Delta}}{T},$$

where we define $a_i = \mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right]$, for $i = 0, 1, \dots, n-1$. Then for $i = 0, 1, \dots, n-1$, we have

$$a_i = \frac{e^{2r\Delta}}{S_0^2} M(2, \Delta) e^{q(2)V_0 \left(\frac{\eta(t_i)e^{-\kappa t_i}}{\eta(t_i) - q(2)} - 1 \right)} \left(\frac{\eta(t_i)}{\eta(t_i) - q(2)} \right)^{\alpha+1},$$

⁵This formula has been implemented in Matlab and its code is available upon request from authors as well as for all other formulas that appear in this paper.

where

$$M(u, t) = \mathbb{E}[e^{uX_t}] = S_0^u e^{\frac{\kappa\theta}{\gamma^2} \left((\kappa - \gamma\rho u - d(u))t - 2 \ln \left(\frac{1 - g(u)e^{-d(u)t}}{1 - g(u)} \right) \right)} e^{V_0 \frac{\kappa - \gamma\rho u - d(u)}{\gamma^2} \frac{1 - e^{-d(u)t}}{1 - g(u)e^{-d(u)t}}},$$

with the following auxiliary functions

$$\begin{aligned} d(u) &= \sqrt{(\kappa - \gamma\rho u)^2 + \gamma^2(u - u^2)}, & g(u) &= \frac{\kappa - \gamma\rho u - d(u)}{\kappa - \gamma\rho u + d(u)}, \\ q(u) &= \frac{\kappa - \gamma\rho u - d(u)}{\gamma^2} \frac{1 - e^{-d(u)\Delta}}{1 - g(u)e^{-d(u)\Delta}}, & \eta(u) &= \frac{2\kappa}{\gamma^2} (1 - e^{-\kappa u})^{-1}. \end{aligned}$$

Proof. See Appendix C. □

Remark 3.1. The formula in the above Proposition 3.2 is consistent with the one obtained in equation (2.34) by Zhu and Lian (2011). In particular, we are able to reproduce all numerical results but one presented in Table 3.1, page 246 of Zhu and Lian (2011) using their set of parameters: $\kappa = 11.35$, $\theta = 0.022$, $\gamma = 0.618$, $\rho = -0.64$, $V_0 = 0.04$, $r = 0.1$, $T = 1$ and $S_0 = 1$ (all numbers match except the case when $n = 4$ we get 263.2 instead of 267.6).

Proposition 3.2 gives a formula for pricing the variance swap with payoff $\frac{1}{T} \sum_{i=0}^{n-1} \left(\frac{S_{t_{i+1}} - S_{t_i}}{S_{t_i}} \right)^2$, but it is straightforward to extend its proof to the following payoff $\frac{1}{T} \sum_{i=0}^{n-1} \left(\frac{S_{t_{i+1}} - S_{t_i}}{S_{t_i}} \right)^k$, with an arbitrary integer power k .

4 Fair Strike of the Discrete Variance Swap in the Hull-White model

The correlated Hull-White stochastic volatility model is as follows

$$(HW) \quad \begin{cases} \frac{dS_t}{S_t} = rdt + \sqrt{V_t}dW_t^{(1)} \\ dV_t = \mu V_t dt + \sigma V_t dW_t^{(2)} \end{cases} \quad (13)$$

where $\mathbb{E}[dW_t^{(1)} dW_t^{(2)}] = \rho dt$. Referring to equation (1), we have $m(x) = \sqrt{x}$, $\mu(x) = \mu x$, $\sigma(x) = \sigma x$, so it is straightforward to determine $f(v) = \frac{2}{\sigma} \sqrt{v}$, $h(v) = \left(\frac{\mu}{\sigma} - \frac{\sigma}{4} \right) \sqrt{v}$, and apply (27) in Lemma A.1 in the Appendix to

obtain

$$S_T = S_0 \exp \left\{ rT - \frac{1}{2} \int_0^T V_t dt + \frac{2\rho}{\sigma} (\sqrt{V_T} - \sqrt{V_0}) - \rho \left(\frac{\mu}{\sigma} - \frac{\sigma}{4} \right) \int_0^T \sqrt{V_t} dt + \sqrt{1 - \rho^2} \int_0^T \sqrt{V_t} dW_t^{(3)} \right\}.$$

Proposition 4.1. (*Fair Strikes in the Hull-White Model*)

In the Hull-White stochastic volatility model (13), the fair strike (2) of the discrete variance swap is

$$\begin{aligned} K_d^{HW}(n) = & \frac{r^2 T}{n} + \left(1 - \frac{rT}{n}\right) K_c^{HW} - \frac{V_0^2 \left(e^{(2\mu+\sigma^2)T} - 1\right) \left(e^{\frac{\mu T}{n}} - 1\right)}{2T\mu(\mu + \sigma^2) \left(e^{\frac{(2\mu+\sigma^2)T}{n}} - 1\right)} \\ & + \frac{V_0^2 \left(e^{(2\mu+\sigma^2)T} - 1\right)}{2T(2\mu + \sigma^2)(\mu + \sigma^2)} + \frac{8\rho \left(e^{\frac{3(4\mu+\sigma^2)T}{8}} - 1\right) V_0^{3/2} \sigma \left(e^{\frac{\mu T}{n}} - 1\right)}{\mu T (4\mu + 3\sigma^2) \left(e^{\frac{3(4\mu+\sigma^2)T}{8n}} - 1\right)} \\ & - \frac{64\rho \left(e^{\frac{3(4\mu+\sigma^2)T}{8}} - 1\right) V_0^{3/2} \sigma}{3T(4\mu + \sigma^2) (4\mu + 3\sigma^2)}. \quad (14) \end{aligned}$$

The fair strike of the continuous variance swap is

$$K_c^{HW} = \frac{1}{T} \mathbb{E} \left[\int_0^T V_s ds \right] = \frac{V_0}{T\mu} (e^{\mu T} - 1). \quad (15)$$

Proof. The proof can be found in Appendix D. \square

5 Fair Strike of the Discrete Variance Swap in the Schöbel-Zhu model

The correlated Schöbel-Zhu stochastic volatility model (see Schöbel and Zhu (1999)) can be described by the following dynamics⁶

$$(SZ) \quad \begin{cases} \frac{dS_t}{S_t} = rdt + V_t dW_t^{(1)} \\ dV_t = \kappa(\theta - V_t)dt + \gamma dW_t^{(2)} \end{cases} \quad (16)$$

where $\mathbb{E}[dW_t^{(1)}dW_t^{(2)}] = \rho dt$. Referring to equation (1), we have $m(x) = x$, $\mu(x) = -\kappa(x - \theta)$, $\sigma(x) = \gamma$, so it is straightforward to apply (27) in Lemma A.1 given in the Appendix with $f(v) = \frac{v^2}{2\gamma}$ and $h(v) = \frac{\kappa\theta}{\gamma}v - \frac{\kappa}{\gamma}v^2 + \frac{\gamma}{2}$ to obtain

$$S_T = S_0 \exp \left\{ \left(r - \frac{\gamma\rho}{2} \right) T - \frac{\kappa\theta\rho}{\gamma} \int_0^T V_t dt - \left(\frac{1}{2} - \frac{\rho\kappa}{\gamma} \right) \int_0^T V_t^2 dt + \frac{\rho}{2\gamma} (V_T^2 - V_0^2) + \sqrt{1 - \rho^2} \int_0^T V_t dW_t^{(3)} \right\}.$$

Proposition 5.1. (*Fair Strikes in the Schöbel-Zhu Model*)

In the Schöbel-Zhu stochastic volatility model (16), the fair strike (2) of the discrete variance swap is computed from (5) but does not have a simple expression.⁷ The fair strike of the continuous variance swap is

$$K_c^{SZ} = \frac{\gamma^2}{2\kappa} + \theta^2 + \left(\frac{(V_0 - \theta)^2}{2\kappa T} - \frac{\gamma^2}{4\kappa^2 T} \right) (1 - e^{-2\kappa T}) + \frac{2\theta(V_0 - \theta)}{\kappa T} (1 - e^{-\kappa T}). \quad (17)$$

Proof. The proof can be found in Appendix E. □

6 Asymptotics

In the time-homogeneous stochastic volatility model, this section presents asymptotics for the fair strikes of discrete variance swaps in the Heston, the

⁶We shall note that here $m(V_t) = V_t$ (where $m(\cdot)$ is defined in (1)) instead of $\sqrt{V_t}$, thus the process V_t models the volatility and not the variance. In particular in the Schöbel-Zhu model, the variance process $Y_t = V_t^2$ follows $dY_t = (\gamma^2 + 2\kappa\theta\sqrt{Y_t} - 2\kappa Y_t)dt + 2\gamma\sqrt{Y_t}dW_t^{(2)}$.

⁷See Proposition 6.7 for an explicit expansion.

Hull-White and the Schöbel-Zhu models based on the explicit expressions derived in the previous sections 3, 4 and 5.

The expansions as functions of the number of sampling periods n are given in Propositions 6.1, 6.4 and 6.7 (respectively for the Heston, Hull-White and Schöbel-Zhu models). In the Heston model, our results are consistent with Proposition 4.2 of Broadie and Jain (2008a), in which it is proved that $K_d^H(n) = K_c^H + \mathcal{O}\left(\frac{1}{n}\right)$. The expansion below is more precise in that at least the first leading term in the expansion is given explicitly. See also Theorem 3.8 of Jarrow et al. (2013) in a more general context. In particular, Jarrow et al. (2013) give a sufficient condition for the convergence of the fair strike of a discrete variance swap to that of a continuously monitored variance swap. In our setting, which is in the absence of jumps, their sufficient condition reduces to $E[\int_0^T m^4(V_s)ds] < \infty$. This latter condition is obviously satisfied in the three examples considered in this paper (the Heston, the Hull-White and the Schöbel-Zhu models).

Expansions as a function of the maturity T (for small maturities) are also given in order to complement results of Keller-Ressel and Muhle-Karbe (2012) (see for example Corollary 2.7 which gives qualitative properties of the discretization gap⁸ as the maturity $T \rightarrow 0$).

6.1 Heston Model

We first expand the fair strike of the discrete variance swap with respect to the number of sampling periods n .

Proposition 6.1. *(Expansion of the fair strike $K_d^H(n)$ w.r.t. n)*

In the Heston model, the expansion of the fair strike of a discrete variance swap, $K_d^H(n)$, is given by

$$K_d^H(n) = K_c^H + \frac{a_1^H}{n} + \mathcal{O}\left(\frac{1}{n^2}\right), \quad (18)$$

where

$$a_1^H = r^2T - rTK_c^H + \left(\frac{\gamma(\theta - V_0)}{2\kappa}(1 - e^{-\kappa T}) - \frac{\theta\gamma T}{2}\right)\rho + \left(\frac{\theta^2}{4} + \frac{\theta\gamma^2}{8\kappa}\right)T + c_1$$

⁸See Definition 2.6 on page 112 of Keller-Ressel and Muhle-Karbe (2012).

with

$$c_1^H = \frac{[\gamma^2\theta - 2\kappa(V_0 - \theta)^2] (e^{-2T\kappa} - 1) + 2(V_0 - \theta)(e^{-T\kappa} - 1) [\gamma^2(e^{-T\kappa} - 1) - 4\kappa\theta]}{16\kappa^2}.$$

Proof. This proposition is a straightforward expansion from (11) in Proposition 3.1. \square

We know that $K_d^H(n) = b^H(n) + \frac{T}{n}r(r - K_c^H)$ from (6) in Remark 2.1. It is thus clear that a_1^H contains all the terms in the risk-free rate r and thus that all the higher terms in the expansion (18) with respect to n are independent of the risk-free rate.

Remark 6.1. *The first term in the expansion (18), a_1^H , is a linear function of ρ . Observe that the coefficient in front of ρ , $\frac{\gamma(\theta - V_0)}{2\kappa}(1 - e^{-\kappa T}) - \frac{\theta\gamma T}{2}$ is negative,⁹ so that a_1^H is always a decreasing function of ρ . We have that*

$$a_1^H \geq 0 \iff \rho \leq \rho_0^H$$

$$\text{where } \rho_0^H = \frac{r^2T - rTK_c^H + \left(\frac{\theta^2}{4} + \frac{\theta\gamma^2}{8\kappa}\right)T + c_1^H}{-\left(\frac{\gamma(\theta - V_0)}{2\kappa}(1 - e^{-\kappa T}) - \frac{\theta\gamma T}{2}\right)}.$$

Proposition 6.2. *(Expansion of the fair strike for small maturity)*

In the Heston model, $K_d^H(n)$ can be expanded when $T \rightarrow 0$ as

$$K_d^H(n) = V_0 + b_1^HT + b_2^HT^2 + \mathcal{O}(T^3) \quad (19)$$

where

$$b_1^H = \frac{\kappa(\theta - V_0)}{2} + \frac{1}{4n}((V_0 - 2r)^2 - 2\rho V_0\gamma)$$

$$b_2^H = \frac{\kappa^2(V_0 - \theta)}{6} + \frac{(V_0 - \theta)\kappa(\gamma\rho + 2r - V_0) + \frac{\gamma^2V_0}{2}}{4n} + \frac{\gamma\rho\kappa(V_0 + \theta) - \frac{\gamma^2V_0}{2}}{12n^2}.$$

Note also that $K_c^H = V_0 + \frac{\kappa}{2}(\theta - V_0)T + \frac{\kappa^2}{6}(V_0 - \theta)T^2 + \mathcal{O}(T^3)$ and thus

$$K_d^H(n) - K_c^H = \frac{1}{4n}((V_0 - 2r)^2 - 2\rho V_0\gamma)T + \mathcal{O}(T^2).$$

Proof. This proposition is a straightforward expansion from (11) in Proposition 3.1. \square

⁹This can be easily seen from the fact that for all $x > 0$, $(\theta - V_0)(1 - e^{-x}) - \theta x \leq \theta(1 - e^{-x} - x) < 0$, and note that here $x = \kappa T > 0$.

Proposition 6.2 is consistent with Corollary 2.7 [b] on page 113 of Keller-Ressel and Muhle-Karbe (2012), where it is clear that the limit of $K_d(n) - K_c$ is 0 when $T \rightarrow 0$.

Notice that in the case $\rho \leq 0$, in the Heston model, $K_d^H(n)$ is non-negative and decreasing in n as the maturity T goes to 0. However, this property cannot be generalized to all correlation levels as it depends on the sign of $(V_0 - 2r)^2 - 2\gamma V_0\rho$.

Proposition 6.3. (*Expression of the fair strike w.r.t. γ*)

In the Heston model, $K_d^H(n)$ is a quadratic function of γ :

$$K_d^H(n) = \frac{1}{8n\kappa^3 T} (h_0^H + h_1^H \gamma + h_2^H \gamma^2), \quad (20)$$

where

$$\begin{aligned} h_0^H &= 2n\kappa (V_0 - \theta)^2 (e^{-2\kappa T} - 1) \frac{1 - e^{-\frac{\kappa T}{n}}}{1 + e^{-\frac{\kappa T}{n}}} + 2\kappa T \left(\kappa^2 T (\theta - 2r)^2 + 4\kappa^2 n\theta \right) \\ &\quad + 4(V_0 - \theta) (2\kappa^2 n + \kappa^2 T (\theta - 2r)) (1 - e^{-\kappa T}), \\ h_1^H &= 8\rho\kappa \left(n\theta(n - ne^{-\frac{\kappa T}{n}} - \kappa T) - (V_0 - \theta) \left(n(1 - e^{-\kappa T}) + \kappa T \frac{1 - e^{-\kappa T}}{1 - e^{-\frac{\kappa T}{n}}} \right) \right), \\ h_2^H &= n(\theta - 2V_0) (e^{-2\kappa T} - 1) \frac{1 - e^{-\frac{\kappa T}{n}}}{1 + e^{-\frac{\kappa T}{n}}} - 2n^2\theta \left(1 - e^{-\frac{\kappa T}{n}} \right) \\ &\quad + 4(V_0 - \theta) \left(n - ne^{-\kappa T} + \kappa T \frac{1 - e^{-\kappa T}}{1 - e^{-\frac{\kappa T}{n}}} \right) + 2\kappa T n\theta. \end{aligned}$$

Proposition 6.3 shows that the discrete fair strike in the Heston model is a quadratic function of the volatility of variance γ . From Figure 6, we observe that the discrete fair strikes evolve in a parabolic shape as γ varies.

6.2 Hull-White Model

Proposition 6.4. (*Expansion of $K_d^{HW}(n)$ w.r.t. n*)

In the Hull-White model, the expansion of the fair strike of the discrete variance swap, $K_d^{HW}(n)$, is given by

$$K_d^{HW}(n) = K_c^{HW} + \frac{a_1^{HW}}{n} + \frac{a_2^{HW}}{n^2} + \frac{a_3^{HW}}{n^3} + \mathcal{O}\left(\frac{1}{n^4}\right) \quad (21)$$

where

$$\begin{aligned}
a_1^{HW} &= r^2 T - r T K_c^{HW} + \frac{V_0^2}{4} \frac{e^{(2\mu+\sigma^2)T} - 1}{2\mu + \sigma^2} - \frac{4\rho\sigma V_0^{\frac{3}{2}}}{3} \frac{e^{\frac{3}{8}(4\mu+\sigma^2)T} - 1}{4\mu + \sigma^2}, \\
a_2^{HW} &= -\frac{V_0^2 \sigma^2 T}{24} \frac{e^{(2\mu+\sigma^2)T} - 1}{2\mu + \sigma^2} - \frac{\rho V_0^{\frac{3}{2}} \sigma T (4\mu - 3\sigma^2)}{36} \frac{e^{\frac{3}{8}(4\mu+\sigma^2)T} - 1}{4\mu + \sigma^2}, \\
a_3^{HW} &= -\frac{\mu T^2 V_0^2 (\mu + \sigma^2)}{48} \frac{e^{(2\mu+\sigma^2)T} - 1}{2\mu + \sigma^2} + \frac{\mu T^2 \rho \sigma V_0^{\frac{3}{2}} (4\mu + 3\sigma^2)}{72} \frac{e^{\frac{3}{8}(4\mu+\sigma^2)T} - 1}{4\mu + \sigma^2}.
\end{aligned}$$

Proof. This proposition is a straightforward expansion from (14) in Proposition 4.1. \square

Observe that $K_d^{HW}(n) = b^{HW}(n) - \frac{K_c^{HW} T}{n} r + \frac{T}{n} r^2$ where $b^{HW}(n) = K_d^{HW}(r=0) > K_c^{HW}$ is independent of r .

If we neglect higher order terms in the expansion (21), we observe that the position of the fair strike of the discrete variance swap with respect to the fair strike of the continuous variance swap is driven by the sign of a_1 and we have the following observation.

Remark 6.2. *The first term in the expansion (21), a_1^{HW} , is a linear function of ρ .*

$$\begin{aligned}
a_1^{HW} \geq 0 &\iff \rho \leq \rho_0^{HW} \\
\text{where } \rho_0^{HW} &= \frac{3(4\mu+\sigma^2) \left(r^2 T - r T K_c^{HW} + \frac{V_0^2}{4} \frac{e^{(2\mu+\sigma^2)T} - 1}{2\mu + \sigma^2} \right)}{4\sigma V_0^{\frac{3}{2}} (e^{\frac{3}{8}(4\mu+\sigma^2)T} - 1)} > 0.
\end{aligned}$$

ρ_0^{HW} can take values strictly larger than 1 as it appears clearly in the right panel of Figure 4. In this latter case, the fair strike of the discrete variance swap is larger than the fair strike of the continuous variance swap for all levels of correlation and for sufficiently high values of n . The minimum value of $K_d^{HW}(n)$ as a function of r is obtained when $r = r^* = \frac{K_c^{HW}}{2}$. After replacing r by r^* in the expression of ρ_0^{HW} , ρ_0^{HW} can easily be shown to be positive.¹⁰

¹⁰It reduces to studying the sign of $\frac{e^{(2\mu+\sigma^2)T} - 1}{(2\mu+\sigma^2)T} - \frac{(e^{\mu T} - 1)^2}{\mu^2 T^2}$. It is an increasing function of σ , so it is larger than $\frac{e^{2\mu T} - 1}{2\mu T} - \frac{(e^{\mu T} - 1)^2}{\mu^2 T^2}$, which is always positive because its minimum is 0 obtained when $\mu T = 0$.

Proposition 6.5. (*Expansion of $K_d^{HW}(n)$ for small maturity*)

In the Hull-White model, $K_d^{HW}(n)$ can be expanded when $T \rightarrow 0$ as

$$K_d^{HW}(n) = V_0 + b_1^{HW}T + b_2^{HW}T^2 + \mathcal{O}(T^3), \quad (22)$$

where

$$\begin{aligned} b_1^{HW} &= \frac{V_0 \mu}{2} + \frac{1}{4n} \left((V_0 - 2r)^2 - 2\rho V_0^{3/2} \sigma \right), \\ b_2^{HW} &= \frac{V_0 \mu^2}{6} + \frac{V_0}{4n} \left(\frac{\sigma^2 V_0}{2} - \frac{3\rho V_0^{1/2} \sigma (\sigma^2 + 4\mu)}{8} + \mu(V_0 - 2r) \right) \\ &\quad + \frac{V_0^{3/2} \sigma (\rho(3\sigma^2 - 4\mu) - 4\sigma\sqrt{V_0})}{96n^2}. \end{aligned}$$

Note also that $K_c^{HW} = V_0 + \frac{V_0 \mu}{2}T + \frac{V_0 \mu^2}{6}T^2 + \mathcal{O}(T^3)$, and thus

$$K_d^{HW}(n) - K_c^{HW} = \frac{1}{4n} \left((V_0 - 2r)^2 - 2\rho V_0^{3/2} \sigma \right) T + \mathcal{O}(T^2).$$

Proof. This proposition is a straightforward expansion from (14) in Proposition 4.1. \square

Note that the expansion for small maturities in the Hull White model is similar to the one in the Heston model given in Proposition 6.2.

Proposition 6.6. (*Expansion of $K_d^{HW}(n)$ w.r.t. σ*)

In the Hull-White model, the fair strike of a discrete variance swap, $K_d^{HW}(n)$, verifies

$$K_d^{HW}(n) = h_0^{HW} + h_1^{HW} \sigma + \mathcal{O}(\sigma^2), \quad (23)$$

where

$$\begin{aligned} h_0^{HW} &= \frac{r^2 T}{n} + \left(1 - \frac{rT}{n} \right) V_0 \frac{e^{T\mu} - 1}{T\mu} - \frac{V_0^2}{2} \frac{e^{2T\mu} - 1}{e^{2\frac{T\mu}{n}} - 1} \frac{e^{\frac{T\mu}{n}} - 1}{T\mu^2} + \frac{V_0^2 (e^{2T\mu} - 1)}{4T\mu^2}, \\ h_1^{HW} &= 2\rho \frac{e^{3/2 T\mu} - 1}{e^{3/2 \frac{T\mu}{n}} - 1} V_0^{3/2} \frac{e^{\frac{T\mu}{n}} - 1}{T\mu^2} - \frac{4\rho (e^{3/2 T\mu} - 1) V_0^{3/2}}{3T\mu^2}. \end{aligned}$$

The expansion of the fair strike in the Hull-White model with respect to the volatility of volatility is very different from the one in the Heston model as it is not a quadratic function of σ , and it also involves higher order terms

of σ .

6.3 Schöbel-Zhu Model

We first expand the fair strike of the discrete variance swap with respect to the number of sampling periods n . The following result is similar to Proposition 6.1 and 6.4. In particular we find that the first term in the expansion is also linear in ρ and has a similar behaviour as in the Heston and Hull-White model.

Proposition 6.7. (*Expansion of $K_d^{SZ}(n)$ w.r.t. n*)

In the Schöbel-Zhu model, the expansion of the fair strike of the discrete variance swap, $K_d^{SZ}(n)$, is given by

$$K_d^{SZ}(n) = K_c^{SZ} + \frac{a_1^{SZ}}{n} + \mathcal{O}\left(\frac{1}{n^2}\right), \quad (24)$$

where

$$a_1^{SZ} = r^2 T - r T K_c^{SZ} + d_1 - d_2 \frac{\gamma}{2\kappa} \rho, \quad (25)$$

with

$$\begin{aligned} d_1 := & \frac{TV_0^4}{4} - \frac{E(T+D)}{16\kappa^2} + \left(\frac{3V_0^2\gamma^2}{4} + \frac{E}{32\kappa} + \frac{\kappa V_0^3(\theta - V_0)}{2} \right) D^2 \\ & + \left(\frac{2\theta\kappa^2 V_0^3}{3} - \frac{V_0^4\kappa^2}{6} - \frac{E}{48} - \frac{V_0^2\theta^2\kappa^2}{2} - \gamma^2\kappa V_0\theta + \frac{3V_0^2\kappa\gamma^2}{4} - \frac{\gamma^4}{4} \right) D^3 \\ & + \left(\frac{E}{8\kappa} + 3\gamma^2(\theta - V_0)\theta + \frac{3V_0^2\gamma^2}{2} + V_0\kappa(\theta - V_0)(2\theta^2 - \theta V_0 + V_0^2) \right) \frac{\kappa^2 D^4}{8}, \end{aligned}$$

and

$$d_2 = T(\gamma^2 + 2\kappa\theta^2) + (2\kappa(\theta^2 - V_0^2) + \gamma^2)D + \frac{\kappa}{2}(\gamma^2 - 2\kappa(\theta - V_0)^2)D^2,$$

where

$$E := 4V_0^4\kappa^2 - 4\theta^4\kappa^2 - 3\gamma^4 - 12\gamma^2\theta^2\kappa, \quad D := \frac{e^{-\kappa T} - 1}{\kappa}.$$

Proof. This proposition is a straightforward expansion from the formula of $K_d^{SZ}(n)$ in Proposition 5.1. Note that although the formula of $K_d^{SZ}(n)$ does

not have a simple form, its asymptotic expansion can be easily computed with Maple for instance. \square

Remark 6.3. *Similarly as in the Heston and the Hull-White models, the first term in the expansion (24), a_1^{SZ} , is a linear function of ρ , but the sign of its slope is not clear in general.*

Proposition 6.8. *(Expansion of the fair strike for small maturity)*

In the Schöbel-Zhu model, $K_d^{SZ}(n)$ can be expanded when $T \rightarrow 0$ as

$$K_d^{SZ}(n) = V_0^2 + b_1^{SZ}T + \mathcal{O}(T^2) \quad (26)$$

where

$$b_1^{SZ} = \kappa V_0(\theta - V_0) + \frac{\gamma^2}{2} + \frac{1}{n} \left(r^2 - rV_0^2 + \frac{V_0^2(V_0^2 - 4\rho\gamma)}{4} \right).$$

Note also that $K_c^{SZ} = V_0^2 + \left(V_0\kappa(\theta - V_0) + \frac{\gamma^2}{2} \right) T + \mathcal{O}(T^2)$ and thus,

$$K_d^{SZ}(n) - K_c^{SZ} = \frac{1}{4n} \left((V_0^2 - 2r)^2 - 4\rho V_0^2\gamma \right) T + \mathcal{O}(T^2).$$

Proof. This proposition is a straightforward expansion from the formula of $K_d^{SZ}(n)$ in Proposition 5.1. \square

Note that the form of the expansion is similar for the three models under study (compare Propositions 6.2, 6.5 and 6.8). We find that the difference between the discrete and the continuous strikes has a first term involving the product of 2ρ by a function of the initial variance value and the volatility of the variance process, and respectively γ in the Heston, σ in the Hull-White and 2γ in the Schöbel-Zhu model. See for example footnote 6 where the dynamics of the variance is derived in the Schöbel-Zhu model.

6.4 Discussion on the convex-order conjecture

As motivated in Keller-Ressel and Griessler (2012), it is of interest to study the *systematic bias* for fixed n and T when using the quadratic variation to approximate the realized variance. Bühler (2006) and Keller-Ressel and Muhle-Karbe (2012) show numerical evidence of this bias (see also Section 7 for further evidence in the Heston and the Hull-White models). Keller-Ressel

and Griessler (2012) propose the following “**convex-order conjecture**”:

$$\mathbb{E}[f(RV(X, \mathcal{P}))] \geq \mathbb{E}[f([X, X]_T)]$$

where f is convex, \mathcal{P} refers to the partition of $[0, T]$ in $n + 1$ division points and $X = \log(S_T/S_0)$. $RV(X, \mathcal{P})$ is the discrete realized variance $(\sum_{i=1}^n (\log(S_{t_i}/S_{t_{i-1}}))^2)$ and $[X, X]_T$ is the continuous quadratic variation $(\int_0^T m^2(V_s)ds)$ in our setting).

When $f(x) = x/T$ and the correlation can be positive, the conjecture is violated, see for example Figure 1 to 4 where $K_d^M(n)$ can be below K_c^M . When $\rho = 0$, the process has conditionally independent increments and satisfies other assumptions in Keller-Ressel and Griessler (2012). Proposition 2.2 ensures that $K_d^M(n) \geq K_c^M$, which is consistent with their results.

7 Numerics

This section illustrates with numerical examples in the Heston, the Hull-White and the Schöbel-Zhu models.

7.1 Heston and Hull-White models

Given parameters for the Heston model, we then choose the parameters in the Hull-White model so that the continuous strikes match. Precisely, we obtain μ by solving numerically $K_c^H = K_c^{HW}$, and find σ such that the variances of V_T in the respective Heston and the Hull-White models match. From (33) and (34), the variance for V_T for the Heston model is given by

$$Var^H(V_T) = \frac{\gamma^2}{2\kappa}(\theta + 2e^{-\kappa T}(V_0 - \theta) + e^{-2\kappa T}(\theta - 2V_0)).$$

The variance for V_T for the Hull-White model can be computed using (41)

$$Var^{HW}(V_T) = V_0^2 e^{2\mu T} (e^{\sigma^2 T} - 1).$$

The parameters for the Heston model are taken from reasonable parameter sets in the literature. Precisely the first set of parameters is similar to the one used by Broadie and Jain (2008a). The second set corresponds to Table 2 in Broadie and Kaya (2006). The values for the parameters of the

Hull-White model are obtained consistently using the procedure described above¹¹.

					Heston			(matched) Hull-White	
	T	r	V_0	ρ	γ	θ	κ	μ	σ
Set 1	1	3.19%	0.010201	-0.7	0.31	0.019	6.21	1.003	0.42
Set 2	5	5%	0.09	-0.3	1	0.09	2	2.9×10^{-9}	0.52

Table 1: Parameter sets

Insert Figure 1

Figure 1 displays cases when the fair strike of the discrete variance swap $K_d^M(n)$ may be smaller than the fair strike of the continuous variance swap K_c^M . The first graph obtained in the Heston model (the model M is denoted by the exponent H for Heston) shows that K_d^H is first higher than K_c^H , crosses this level and stays below K_c^H until it converges to the value K_c^H as $n \rightarrow \infty$. It means that options on discrete realized variance may be overvalued when the continuous quadratic variation is used to approximate the discrete realized variance. Note that this unusual pattern happens when $\rho = 0.7$, which may happen for example in foreign exchange markets.

Insert Figure 2

Figure 2 highlights another type of convergence showing the complexity of the behaviour of the fair strike of the discrete variance swap with respect to that of the continuous variance swap.

Insert Figure 3

Figure 3 displays on the same graphs the discrete fair strike $K_d(n)$ and the first two terms of the expansion formula $K_c^H + \frac{a_1^H}{n}$ for the Heston model and $K_c^{HW} + \frac{a_1^{HW}}{n}$ for the Hull-White model (see Propositions 6.1 and 6.4 for the exact expressions of a_1^H and a_1^{HW}). It shows that the first term of this

¹¹For the two sets of parameters above, we compute the critical interest rate r^* as defined in Remark 2.1. Set 1: $r^* = 0.88\%$; Set 2: $r^* = 0.605\%$, and we can see that the interest rates are both larger than r^* .

expansion is already highly informative as it clearly appears to fit very well for small values of n in both models.

Insert Figure 4

Figure 4 further illustrates that the discrete fair strike (for a daily monitoring) can be lower than the continuous fair strike as $K_d^M - K_c^M$ may be negative for high values of the correlation coefficient both in the Heston and the Hull-White models. In Remark 6.1 and 6.2, it is noted that the first term in the asymptotic expansion with respect to n is linear in ρ . From Figure 3 it is clear that the first term has an important explanatory power. This justifies the linear behavior observed in Figure 4 of the difference between discrete and continuous fair strikes with respect to ρ . Computations of ρ_0^H and ρ_0^{HW} for each of the risk-free rate levels $r = 0\%$, $r = 3.2\%$ and $r = 6\%$ confirm that it is always positive when $r = 0\%$ (which is consistent with Proposition 2.2) and that it can be higher than 1, which ensures that for n sufficiently high, the discrete fair strike is always higher than the continuous fair strike.

Insert Figure 5

Figure 5 shows that as the time to maturity T goes to 0, the discrete fair strike is converging to the continuous fair strike at approximately a quadratic rate. This is consistent with Proposition 6.2 and Proposition 6.5.

Insert Figure 6

Figure 6 shows that the discrepancy between the discrete fair strike and the continuous fair strike is exacerbated by the volatility of the underlying variance process. We observe that the gap between the discrete fair strike and the continuous fair strike, with respect to γ , is wider in the Heston model than in the Hull-White model. This illustrates, from a numerical viewpoint, that the discrete fair strike in the Heston model is more sensitive to the volatility of variance parameter than that of the Hull-White model. In particular, the continuous fair strike K_c^H is independent of γ . For each γ we compute the corresponding σ for the Hull-White model such that the variances match as described in Section 7.1. We then observe similar patterns

in the Heston and the Hull-White models. From the left panel of Figure 6, we can see that the shape of the discrete fair strike in the Heston model with respect to γ evolves similar to a parabola, and this is consistent with Proposition 6.3. The right panel of Figure 6 is consistent with Proposition 6.6.

7.2 Schöbel-Zhu model

For the Schöbel-Zhu model, we reproduce a similar numerical analysis and take parameters consistent with the Heston model. Note that the V process in the Schöbel-Zhu model corresponds to the volatility process instead of the variance process¹². Then we choose $\theta = \sqrt{0.019}$ and $V_0 = \sqrt{0.010201}$. Other parameters are taken from set 1 of Table 1.

Insert Figure 7

Both the left and right panels of Figure 7 show that K_d^{SZ} can be below K_c^{SZ} until it converges to the value K_c^{SZ} as $n \rightarrow \infty$. This unusual pattern happens when the correlation is positive similarly in the Heston and the Hull-White models.

Insert Figure 8

Figure 8 illustrates that the discrete fair strike (for a daily monitoring) can be lower than the continuous fair strike as $K_d^{SZ} - K_c^{SZ}$ may be negative for high values of the correlation coefficient. From Figure 8 it is clear that the first term also has an important explanatory power. This justifies the linear behavior observed in Figure 8 of the difference between discrete and continuous fair strikes with respect to ρ . Computations of ρ_0^{SZ} (defined as the zero of a_1^{SZ} computed in Proposition 6.7) for each of the risk-free rate levels $r = 0\%$, $r = 3.2\%$ and $r = 6\%$ confirm that it is always positive when $r = 0\%$ (which is consistent with Proposition 2.2).

¹²The notation V_t in the Schöbel-Zhu model corresponds to the square root of what is denoted by V_t in the Heston model.

8 Conclusions

This paper provides explicit expressions of the fair strikes of discretely sampled variance swaps in the Heston, the Hull-White and the Schöbel-Zhu models. For the Heston model, the explicit closed-form formula simplifies the expressions obtained by Broadie and Jain (2008a) in equations (A-29) and (A-30) on page 793, where several sums from 0 to n are involved. Our formulae are more explicit (as there are no sums involved in the discrete fair strikes), and easier to use. The explicit closed-form formulas for the Hull-White and the Schöbel-Zhu models are new. Asymptotics of the fair strikes with respect to key parameters such as $n \rightarrow \infty$, $T \rightarrow 0$, $\gamma \rightarrow 0$ are new and consistent with theoretical results obtained in Keller-Ressel and Muhle-Karbe (2012).

There are several potential research directions. For example this work can be extended to mixed exponential jump diffusions models (proposed by Cai and Kou (2011) and to the calculations of fair strikes for gamma swaps (see for example Lee (2010) for a definition of the payoff). For the 3/2 stochastic volatility model treated in Itkin and Carr (2010), the difficulty lies in obtaining closed-form expressions of the covariance terms $E[h(V_t)h(V_s)]$ for some functional h , and the determination of a closed-form formula for the fair strike in this model is left as an open problem. Pricing discrete volatility derivatives with non-linear payoffs (e.g. puts and calls on realized variance) in time-homogeneous stochastic volatility models is also left as a future research direction.

A Proof of Proposition 2.1

Using Itô's lemma and Cholesky decomposition, (1) becomes

$$\begin{aligned} d(\ln(S_t)) &= \left(r - \frac{1}{2}m^2(V_t)\right)dt + \rho m(V_t)dW_t^{(2)} + \sqrt{1-\rho^2}m(V_t)dW_t^{(3)} \\ dV_t &= \mu(V_t)dt + \sigma(V_t)dW_t^{(2)}. \end{aligned}$$

where $W_t^{(2)}$ and $W_t^{(3)}$ are two standard independent Brownian motions.

Proposition 2.1 is then a direct application of the following lemma (see Lemma 3.1 of Bernard and Cui (2011) for its proof).

Lemma A.1. *Under the model given in (1), we have*

$$S_T = S_0 \exp \left\{ rT - \frac{1}{2} \int_0^T m^2(V_t)dt + \rho(f(V_T) - f(V_0)) - \rho \int_0^T h(V_t)dt + \sqrt{1-\rho^2} \int_0^T m(V_t)dW_t^{(3)} \right\}, \quad (27)$$

where $f(v) = \int_0^v \frac{m(z)}{\sigma(z)}dz$ and $h(v) = \mu(v)f'(v) + \frac{1}{2}\sigma^2(v)f''(v)$.

Now from equation (27) in Lemma A.1, we compute the following key elements in the fair strike of the discrete variance swap. Assume that the time interval is $[t, t + \Delta]$, then

$$\begin{aligned} \ln \left(\frac{S_{t+\Delta}}{S_t} \right) &= r\Delta - \frac{1}{2} \int_t^{t+\Delta} m^2(V_s)ds + \rho \left(f(V_{t+\Delta}) - f(V_t) - \int_t^{t+\Delta} h(V_s)ds \right) \\ &\quad + \sqrt{1-\rho^2} \int_t^{t+\Delta} m(V_s)dW_s^{(3)}. \end{aligned}$$

Then we can compute

$$\begin{aligned} \mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] &= r^2\Delta^2 + \frac{1}{4}\mathbb{E} \left[\left(\int_t^{t+\Delta} m^2(V_s)ds \right)^2 \right] - r\Delta\mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s)ds \right] + \mathbb{E} [A^2] \\ &\quad + \mathbb{E} \left[\left(2r\Delta - \int_t^{t+\Delta} m^2(V_s)ds \right) A \right] + (1-\rho^2)\mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s)ds \right], \end{aligned} \quad (28)$$

where $A = \rho \left(f(V_{t+\Delta}) - f(V_t) - \int_t^{t+\Delta} h(V_s)ds \right)$, and

$$A^2 = \rho^2 \left((f(V_{t+\Delta}) - f(V_t))^2 + \left(\int_t^{t+\Delta} h(V_s)ds \right)^2 - 2(f(V_{t+\Delta}) - f(V_t)) \int_t^{t+\Delta} h(V_s)ds \right).$$

Using the above expressions for A and A^2 in (28), we obtain

$$\begin{aligned}
& \mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] \\
&= r^2 \Delta^2 + \frac{1}{4} \mathbb{E} \left[\left(\int_t^{t+\Delta} m^2(V_s) ds \right)^2 \right] + (1 - \rho^2 - r\Delta) \mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s) ds \right] \\
&\quad + \rho^2 \mathbb{E}[(f(V_{t+\Delta}) - f(V_t))^2] + \rho^2 \mathbb{E} \left[\left(\int_t^{t+\Delta} h(V_s) ds \right)^2 \right] + 2r\rho\Delta \mathbb{E}[(f(V_{t+\Delta}) - f(V_t))] \\
&\quad - \mathbb{E} \left[(f(V_{t+\Delta}) - f(V_t)) \int_t^{t+\Delta} (2\rho^2 h(V_s) + \rho m^2(V_s)) ds \right] - 2r\rho\Delta \mathbb{E} \left[\int_t^{t+\Delta} h(V_s) ds \right] \\
&\quad + \rho \mathbb{E} \left[\left(\int_t^{t+\Delta} h(V_s) ds \right) \left(\int_t^{t+\Delta} m^2(V_s) ds \right) \right]. \tag{29}
\end{aligned}$$

By Itô's lemma, f defined in Lemma A.1 verifies $df(V_t) = h(V_t)dt + m(V_t)dW_t^{(2)}$. Integrating the above SDE from t to $t + \Delta$, we have

$$f(V_{t+\Delta}) - f(V_t) = \int_t^{t+\Delta} h(V_s) ds + \int_t^{t+\Delta} m(V_s) dW_s^{(2)}.$$

Thus

$$\mathbb{E} [f(V_{t+\Delta}) - f(V_t)] - \mathbb{E} \left[\int_t^{t+\Delta} h(V_s) ds \right] = \mathbb{E} \left[\int_t^{t+\Delta} m(V_s) dW_s^{(2)} \right] = 0. \tag{30}$$

Rearrange (29) and use (30) to simplify the terms, and we obtain

$$\begin{aligned}
& \mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] = r^2 \Delta^2 - r\Delta \mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s) ds \right] + \frac{1}{4} \mathbb{E} \left[\left(\int_t^{t+\Delta} m^2(V_s) ds \right)^2 \right] \\
&\quad + (1 - \rho^2) \mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s) ds \right] + \rho^2 \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))^2] \\
&\quad + \rho^2 \mathbb{E} \left[\left(\int_t^{t+\Delta} h(V_s) ds \right)^2 \right] + \rho \mathbb{E} \left[\int_t^{t+\Delta} h(V_s) ds \int_t^{t+\Delta} m^2(V_s) ds \right] \\
&\quad - \rho \mathbb{E} \left[(f(V_{t+\Delta}) - f(V_t)) \int_t^{t+\Delta} (2\rho h(V_s) + m^2(V_s)) ds \right]. \tag{31}
\end{aligned}$$

Now we apply Fubini's theorem and partial integration to further simplify (31). Note that $m^2(V_s) \geq 0$, Q-a.s., then by Fubini's theorem for non-negative measurable functions, $\mathbb{E} \left[\int_t^{t+\Delta} m^2(V_s) ds \right] = \int_t^{t+\Delta} \mathbb{E} [m^2(V_s)] ds$.

Similarly we have $\mathbb{E} \left[\left(\int_t^{t+\Delta} m^2(V_s) ds \right)^2 \right] = \int_t^{t+\Delta} \int_t^{t+\Delta} \mathbb{E} [m^2(V_s) m^2(V_u)] ds du$ for any $t \leq s \leq t + \Delta$ and any $t \leq u \leq t + \Delta$,

If $\mathbb{E} [|h(V_s)h(V_u)|] < \infty$ for any $t \leq s \leq t + \Delta$ and any $t \leq u \leq t + \Delta$, then we have $\mathbb{E} \left[\left(\int_t^{t+\Delta} h(V_s) ds \right)^2 \right] = \int_t^{t+\Delta} \int_t^{t+\Delta} \mathbb{E} [h(V_s)h(V_u)] ds du$.

If $\mathbb{E} [|h(V_s)m^2(V_u)|] < \infty$ for any $t \leq s \leq t + \Delta$ and any $t \leq u \leq t + \Delta$, then we have

$$\mathbb{E} \left[\int_t^{t+\Delta} h(V_s) ds \int_t^{t+\Delta} m^2(V_s) ds \right] = \int_t^{t+\Delta} \int_t^{t+\Delta} \mathbb{E} [h(V_s)m^2(V_u)] ds du.$$

If $\mathbb{E} [(f(V_{t+\Delta}) - f(V_t))(2\rho h(V_s) + m^2(V_s))] < \infty$ for all $t \leq s \leq t + \Delta$, then we have

$$\begin{aligned} & \mathbb{E} \left[(f(V_{t+\Delta}) - f(V_t)) \int_t^{t+\Delta} (2\rho h(V_s) + m^2(V_s)) ds \right] \\ &= \int_t^{t+\Delta} \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))(2\rho h(V_s) + m^2(V_s))] ds. \end{aligned}$$

Thus we finally have proved (5) from Proposition 2.1. This completes the proof. \square

B Proof of Proposition 3.1

Proof. We apply Proposition 2.1 to the Heston stochastic volatility model. We first compute $f(x) = \frac{x}{\gamma}$ and $h(x) = \frac{\kappa\theta - \kappa x}{\gamma}$, then we have

$$\begin{aligned} K_d^H &= \frac{1}{T} \sum_{i=0}^{n-1} \mathbb{E} \left[\left(\ln \frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] \\ &= \frac{1}{T} \left(\frac{a^2 T^2}{n} + b^2 \sum_{i=0}^{n-1} \int_{t_i}^{t_{i+1}} \int_{t_i}^{t_{i+1}} \mathbb{E}[V_s V_u] ds du + \left(\frac{2abT}{n} + 1 - \rho^2 \right) \int_0^T \mathbb{E}[V_s] ds \right. \\ &\quad + \frac{\rho^2}{\gamma^2} \sum_{i=0}^{n-1} \mathbb{E}[(V_{t_{i+1}} - V_{t_i})^2] + \frac{2\rho a T}{n\gamma} (\mathbb{E}[V_T] - \mathbb{E}[V_0]) \\ &\quad \left. + \frac{2\rho b}{\gamma} \sum_{i=0}^{n-1} \left(\int_{t_i}^{t_{i+1}} \mathbb{E}[V_{t_{i+1}} V_s] ds - \int_{t_i}^{t_{i+1}} \mathbb{E}[V_{t_i} V_s] ds \right) \right). \end{aligned} \quad (32)$$

Furthermore, for all $t \geq 0$

$$\mathbb{E}[V_t] = \theta + e^{-\kappa t} (V_0 - \theta), \quad (33)$$

and for all $0 < s \leq t$

$$\begin{aligned} \mathbb{E}[V_t V_s] &= \theta^2 + e^{-\kappa t}(V_0 - \theta) \left(\theta + \frac{\gamma^2}{\kappa} \right) + e^{-\kappa s} \theta (V_0 - \theta) \\ &\quad + e^{-\kappa(t+s)} \left((\theta - V_0)^2 + \frac{\gamma^2}{2\kappa} (\theta - 2V_0) \right) + \frac{\gamma^2}{2\kappa} \theta e^{-\kappa(t-s)}. \end{aligned} \quad (34)$$

In particular, this formula holds for $t = s$ and gives $\mathbb{E}[V_t^2]$. These formulas already appear in Broadie and Jain (2008a) (formula (A-15)). To compute K_d^H , (33) and (34) are the only expressions needed, and they should then be integrated and summed.

We have computed all terms in (32) with the help of Maple and also have simplified the final expression given by Maple. It turns out that in the case of the Heston model, all terms can be computed explicitly and the final simplified expression for (32) does not require any sums or integrals. We finally obtain an explicit formula for K_d^H as a function of the parameters of the model. This completes the proof. \square

C Proof of Proposition 3.2

Proof. Denote the log stock price without drift as $X_t = \ln S_t - rt$, and $X_0 = x_0$. Denote $V_0 = v_0$, $\Delta = T/n$. We have that $\mathbb{E} \left[\left(\frac{S_{t_{i+1}} - S_{t_i}}{S_{t_i}} \right)^2 \right] = \mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] + 1 - 2e^{r\Delta}$. Thus the goal is to calculate the second moment $\mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right]$, and note that it is closely linked to the moment generating function of the log stock price X . Recall the following formulation of the moment generating function $M(u, t) = \mathbb{E}[e^{uX_t}]$ from Albrecher et al. (2007)

$$\begin{aligned} M(u, t) &= S_0^u \exp \left\{ \frac{\kappa\theta}{\gamma^2} \left((\kappa - \gamma\rho u - d(u))t - 2 \ln \left(\frac{1 - g(u)e^{-d(u)t}}{1 - g(u)} \right) \right) \right\} \\ &\quad \times \exp \left\{ V_0 \frac{\kappa - \gamma\rho u - d(u)}{\gamma^2} \frac{1 - e^{-d(u)t}}{1 - g(u)e^{-d(u)t}} \right\}, \end{aligned} \quad (35)$$

where the auxiliary functions are given by

$$d(u) = \sqrt{(\kappa - \gamma\rho u)^2 + \gamma^2(u - u^2)}, \quad g(u) = \frac{\kappa - \gamma\rho u - d(u)}{\kappa - \gamma\rho u + d(u)}.$$

We first separate out the case of $i = 0$ and $i = 1, \dots, n-1$. For the first case,

we have

$$\mathbb{E} \left[\left(\frac{S_{t_1}}{S_0} \right)^2 \right] = \frac{1}{S_0^2} \mathbb{E} \left[e^{2 \ln S_{t_1}} \right] = \frac{e^{2r t_1}}{S_0^2} M(2, t_1) = \frac{e^{2r \Delta}}{S_0^2} M(2, \Delta). \quad (36)$$

For the second case, with $i = 1, 2, \dots, n-1$, we have

$$\begin{aligned} \mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] &= \mathbb{E} \left[e^{2 \ln \left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)} \right] = e^{2r \Delta} \mathbb{E} \left[\mathbb{E} \left[e^{2(X_{t_{i+1}} - X_{t_i})} \mid \mathcal{F}_{t_i} \right] \right] \\ &= \exp \left\{ 2r \Delta + \frac{\kappa \theta}{\gamma^2} \left((\kappa - 2\gamma\rho - d(2))\Delta - 2 \ln \frac{1 - g(2)e^{-d(2)\Delta}}{1 - g(2)} \right) \right\} \\ &\times \mathbb{E} \left[\exp \left\{ V_{t_i} \frac{\kappa - 2\gamma\rho - d(2)}{\gamma^2} \frac{1 - e^{-d(2)\Delta}}{1 - g(2)e^{-d(2)\Delta}} \right\} \right]. \end{aligned} \quad (37)$$

We first define $\alpha = 2\kappa\theta/\gamma^2 - 1 \geq 0$, and $\eta(t) = \frac{2\kappa}{\gamma^2}(1 - e^{-\kappa t})^{-1}$. Then from Theorem 3.1¹³ in Hurd and Kuznetsov (2008), we have

$$\mathbb{E}[e^{uV_T}] = \left(\frac{\eta(T)}{\eta(T) - u} \right)^{\alpha+1} e^{V_0 \frac{\eta(T)u}{\eta(T)-u} e^{-\kappa T}}. \quad (38)$$

Combine equations (37) and (38), for $i = 1, \dots, n-1$, we finally have

$$\mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] = e^{2r \Delta + \frac{\kappa \theta}{\gamma^2} \left((\kappa - 2\gamma\rho - d(2))\Delta - 2 \ln \frac{1 - g(2)e^{-d(2)\Delta}}{1 - g(2)} \right)} e^{V_0 \frac{\eta(t_i)q(2)}{\eta(t_i) - q(2)} e^{-\kappa t_i}} \left(\frac{\eta(t_i)}{\eta(t_i) - q(2)} \right)^{\alpha+1}, \quad (39)$$

where $q(u) = \frac{\kappa - \gamma\rho u - d(u)}{\gamma^2} \frac{1 - e^{-d(u)\Delta}}{1 - g(u)e^{-d(u)\Delta}}$. Using the definition of $M(u, t)$, we can factor out $M(2, \Delta)$ from (39) and finally we have

$$\mathbb{E} \left[\left(\frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] = \frac{e^{2r \Delta}}{S_0^2} M(2, \Delta) e^{q(2)V_0 \left(\frac{\eta(t_i)e^{-\kappa t_i}}{\eta(t_i) - q(2)} - 1 \right)} \left(\frac{\eta(t_i)}{\eta(t_i) - q(2)} \right)^{\alpha+1}. \quad (40)$$

When $i = 0$, we have $t_i = 0$ and since $\eta_u \rightarrow \infty$ as $u \rightarrow 0$, we use L'Hôpital's rule

$$\frac{\eta_{t_0}}{\eta_{t_0} - q(2)} = \lim_{u \rightarrow 0} \frac{\eta_u}{\eta_u - q(2)} = \lim_{u \rightarrow 0} \frac{\eta'_u}{\eta'_u} = 1.$$

Thus a_0 is a special case of the formula in (40) when $i = 0$. From Theorem 3.1 in Hurd and Kuznetsov (2008), equation (35) and consequently the above

¹³Note that in terms of our notation, the parameters in Hurd and Kuznetsov (2008) and our parameters have the correspondence $a = \kappa\theta$, $b = \kappa$, $c = \gamma$.

(37), (38) are well-defined if $u < \eta(T)^{14}$. Note that the formula (40) involves the $u = 2$ case. A sufficient condition for $u = 2 < \eta(T)$ to hold is $\gamma^2 T < 1$ (since $2 < \eta(T)$ is equivalent to $1 - \frac{\kappa}{\gamma^2} < e^{-\kappa T}$).

Then the final formula for the discrete fair strike follows by summing the above terms $a_i, i = 0, 1, \dots, n-1$. This completes the proof. \square

D Proof of Proposition 4.1

Proof. For the Hull-White model, from Proposition 2.1, we first compute $f(x) = \frac{2}{\sigma}\sqrt{x}$ and $h(x) = \left(\frac{\mu}{\sigma} - \frac{\sigma}{4}\right)\sqrt{x}$, then we have

$$\begin{aligned} \mathbb{E} \left[\left(\ln \frac{S_{t_{i+1}}}{S_{t_i}} \right)^2 \right] &= (1 - \rho^2 - \frac{rT}{n}) \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \mathbb{E}[V_s] ds + r^2 \frac{T^2}{n^2} \\ &\quad - \frac{2\rho}{\sigma} \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \mathbb{E} \left[\left(\sqrt{V_{\frac{(i+1)T}{n}}} - \sqrt{V_{\frac{iT}{n}}} \right) V_s \right] ds + 2\rho^2 q^2 \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \int_{\frac{iT}{n}}^u \mathbb{E} \left[\sqrt{V_s} \sqrt{V_u} \right] ds du \\ &\quad + \frac{4\rho^2}{\sigma^2} \mathbb{E} \left[\left(\sqrt{V_{\frac{(i+1)T}{n}}} - \sqrt{V_{\frac{iT}{n}}} \right)^2 \right] - \frac{4\rho^2 q}{\sigma} \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \mathbb{E} \left[\left(\sqrt{V_{\frac{(i+1)T}{n}}} - \sqrt{V_{\frac{iT}{n}}} \right) \sqrt{V_s} \right] ds \\ &\quad + \frac{1}{2} \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \int_{\frac{iT}{n}}^u \mathbb{E}[V_s V_u] ds du + \rho q \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \int_{\frac{iT}{n}}^u \mathbb{E} \left[\sqrt{V_s} V_u \right] ds du \\ &\quad + \rho q \int_{\frac{iT}{n}}^{\frac{(i+1)T}{n}} \int_u^{\frac{(i+1)T}{n}} \mathbb{E} \left[\sqrt{V_s} V_u \right] ds du, \end{aligned}$$

with $q = \frac{\mu}{\sigma} - \frac{\sigma}{4}$.

We now compute the following covariance terms that are useful in the simplification of the fair strike $K_d^{HW}(n)$. In the Hull-White model, the stochastic variance process V_t follows a geometric Brownian motion. Thus we have $V_t = V_0 \exp \left(\left(\mu - \frac{\sigma^2}{2} \right) t + \sigma W_t^{(2)} \right)$. Note that

$$\mathbb{E}[V_s^a] = V_0^a e^{a\mu s} e^{\frac{a^2 - a}{2} \sigma^2 s},$$

which will be useful below for $a = 1/2$, $a = 1$ and $a = 2$.

$$\mathbb{E}[V_s] = V_0 e^{\mu s}, \quad \mathbb{E}[\sqrt{V_s}] = \sqrt{V_0} e^{\frac{\mu}{2}s - \frac{1}{8}\sigma^2 s} = \sqrt{V_0} e^{\frac{\sigma}{2}qs}, \quad \mathbb{E}[V_s^2] = V_0^2 e^{2\mu s + \sigma^2 s}.$$

The fair strike for the continuous variance swap is straightforward and

¹⁴Note that $\eta(t)$ is a decreasing function in t , thus $u < \eta(T)$ is sufficient for $u < \eta(t_i)$ for all $i = 0, 1, \dots, n$.

is equal to $\mathbb{E} \left[\int_0^T V_s ds \right] = \frac{V_0}{\mu} (e^{\mu T} - 1)$. Similarly

$$\mathbb{E} \left[\int_0^T \sqrt{V_s} ds \right] = \int_0^T \sqrt{V_0} e^{\frac{\sigma}{2} qs} ds = \sqrt{V_0} \frac{2}{\sigma q} \left(e^{\frac{\sigma q T}{2}} - 1 \right),$$

and for $s < u$, we have the following results

$$\begin{aligned} \mathbb{E} [V_s V_u] &= V_0^2 \exp(\mu(u+s) + \sigma^2 s), \\ \mathbb{E} [\sqrt{V_s} \sqrt{V_u}] &= V_0 \exp\left(\frac{\mu}{2}(u+s) - \frac{\sigma^2}{8}(u-s)\right), \\ \mathbb{E} [\sqrt{V_s} V_u] &= V_0^{\frac{3}{2}} \exp\left(\mu\left(\frac{s}{2} + u\right) + \frac{3\sigma^2}{8}s\right), \\ \mathbb{E} [V_s \sqrt{V_u}] &= V_0^{\frac{3}{2}} \exp\left(\mu\left(s + \frac{u}{2}\right) - \frac{\sigma^2}{8}u + \frac{\sigma^2}{2}s\right). \end{aligned} \quad (41)$$

After some tedious calculations with the help of Maple, we can obtain an explicit formula as the one appearing in Proposition 4.1. This completes the proof. \square

E Proof of Proposition 5.1

Proof. For the Schöbel-Zhu model, from the key equation in Proposition 2.1, we have

$$\begin{aligned} \mathbb{E} \left[\left(\ln \frac{S_{t+\Delta}}{S_t} \right)^2 \right] &= r^2 \Delta^2 + (1 - \rho^2 - r\Delta) \int_t^{t+\Delta} m_1(s) ds - \rho \int_t^{t+\Delta} m_5(t, s) ds \\ &\quad + \frac{1}{4} \int_t^{t+\Delta} \int_t^{t+\Delta} m_2(s, u) ds du + \frac{\rho^2}{4\gamma^2} \mathbb{E} \left[(V_{t+\Delta}^2 - V_t^2)^2 \right] \\ &\quad + \rho^2 \int_t^{t+\Delta} \int_t^{t+\Delta} m_3(s, u) ds du + \rho \int_t^{t+\Delta} \int_t^{t+\Delta} m_4(s, u) ds du, \end{aligned} \quad (42)$$

where

$$\begin{aligned} m_1(s) &:= \mathbb{E} [m^2(V_s)] = \mathbb{E} [V_s^2], t \leq s \leq t + \Delta, \\ m_2(s, u) &:= \mathbb{E} [m^2(V_s) m^2(V_u)] = \mathbb{E} [V_s^2 V_u^2], t \leq s \leq t + \Delta, \quad t \leq u \leq t + \Delta, \\ m_3(s, u) &:= \mathbb{E} [h(V_s) h(V_u)], t \leq s \leq t + \Delta, \quad t \leq u \leq t + \Delta, \\ m_4(s, u) &:= \mathbb{E} [h(V_s) m^2(V_u)], t \leq s \leq t + \Delta, \quad t \leq u \leq t + \Delta, \\ m_5(t, s) &:= \mathbb{E} [(f(V_{t+\Delta}) - f(V_t))(2\rho h(V_s) + m^2(V_s))], t \leq s \leq t + \Delta, \end{aligned} \quad (43)$$

and $\mathbb{E} [(V_{t+\Delta}^2 - V_t^2)^2] = \mathbb{E} [V_{t+\Delta}^4] + \mathbb{E} [V_t^4] - 2\mathbb{E} [V_{t+\Delta}^2 V_t^2]$. We compute

the following two terms in (42) by expanding the products out. For $s \leq u$

$$\begin{aligned} m_3(s, u) &= \mathbb{E} \left[\left(\frac{\kappa\theta}{\gamma} V_s - \frac{\kappa}{\gamma} V_s^2 + \frac{\gamma}{2} \right) \left(\frac{\kappa\theta}{\gamma} V_u - \frac{\kappa}{\gamma} V_u^2 + \frac{\gamma}{2} \right) \right] \\ &= \mathbb{E} \left[\frac{\kappa^2\theta^2}{\gamma^2} V_s V_u - \frac{\kappa^2\theta}{\gamma^2} (V_s V_u^2 + V_s^2 V_u) + \frac{\kappa\theta}{2} (V_s + V_u) \right. \\ &\quad \left. - \frac{\kappa}{2} (V_s^2 + V_u^2) + \frac{\kappa^2}{\gamma^2} V_s^2 V_u^2 + \frac{\gamma^2}{4} \right], \end{aligned}$$

and for $t \leq s \leq t + \Delta$

$$\begin{aligned} m_5(t, s) &= \frac{1}{2\gamma} \mathbb{E} \left[(V_{t+\Delta}^2 - V_t^2) \left(2\rho \left(\frac{\kappa\theta}{\gamma} V_s - \frac{\kappa}{\gamma} V_s^2 + \frac{\gamma}{2} \right) + V_s^2 \right) \right] \\ &= \mathbb{E} \left[\frac{\rho\kappa\theta}{\gamma^2} (V_{t+\Delta}^2 V_s - V_t^2 V_s) + \frac{\gamma - 2\rho\kappa}{2\gamma^2} (V_{t+\Delta}^2 V_s^2 - V_t^2 V_s^2) \right. \\ &\quad \left. + \frac{\rho}{2} (V_{t+\Delta}^2 - V_t^2) \right]. \end{aligned}$$

It is clear from the above expressions of m_i for $i = 1, 2, \dots, 5$ that they are all functions of $\mathbb{E}[V_s]$, $\mathbb{E}[V_s^2]$, $\mathbb{E}[V_s^4]$, $\mathbb{E}[V_s V_u]$, $\mathbb{E}[V_s^2 V_u]$, $\mathbb{E}[V_s V_u^2]$ and $\mathbb{E}[V_s^2 V_u^2]$. We now compute these seven expressions.

Lemma E.1. *For the Ornstein-Uhlenbeck process V , introduce the auxiliary deterministic functions $e_s := (V_0 - \theta)e^{-\kappa s} + \theta$, and $v(s) := \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa s})$, then*

$$\mathbb{E}[V_s] = e_s, \quad (44)$$

$$\mathbb{E}[V_s^2] = e_s^2 + v(s), \quad (45)$$

$$\mathbb{E}[V_s^3] = e_s^3 + 3e_s v(s), \quad (46)$$

$$\mathbb{E}[V_s^4] = e_s^4 + 6e_s^2 v(s) + 3v^2(s). \quad (47)$$

For $t \leq s \leq u \leq t + \Delta$

$$\begin{aligned} \mathbb{E}[V_s V_u] &= e^{-\kappa(u-s)} \mathbb{E}[V_s^2] + \theta(1 - e^{-\kappa(u-s)}) \mathbb{E}[V_s], \\ \mathbb{E}[V_s^2 V_u^2] &= e^{-2\kappa(u-s)} \mathbb{E}[V_s^4] + 2\theta e^{-\kappa(u-s)} (1 - e^{-\kappa(u-s)}) \mathbb{E}[V_s^3] \\ &\quad + \left(\theta^2 (1 - e^{-\kappa(u-s)})^2 + \frac{\gamma^2}{2\kappa} (1 - e^{-2\kappa(u-s)}) \right) \mathbb{E}[V_s^2]. \end{aligned} \quad (48)$$

For $t \leq s \leq u \leq t + \Delta$

$$\begin{aligned} \mathbb{E}[V_s V_u^2] &= e^{-2\kappa(u-s)} \mathbb{E}[V_s^3] + 2\theta e^{-\kappa(u-s)} (1 - e^{-\kappa(u-s)}) \mathbb{E}[V_s^2] \\ &\quad + \left(\theta^2 (1 - e^{-\kappa(u-s)})^2 + \frac{\gamma^2}{2\kappa} (1 - e^{-2\kappa(u-s)}) \right) \mathbb{E}[V_s]. \end{aligned} \quad (49)$$

For $t \leq s \leq u \leq t + \Delta$

$$\mathbb{E} [V_s^2 V_u] = e^{-\kappa(u-s)} \mathbb{E} [V_s^3] + \theta(1 - e^{-\kappa(u-s)}) \mathbb{E} [V_s^2]. \quad (50)$$

Proof. The stochastic variance process V_s follows

$$dV_s = -\kappa(V_s - \theta)ds + \gamma dW_s^{(2)}.$$

On page 120 of Jeanblanc, Yor and Chesney (2009), one finds that the exact solution of the above SDE is

$$V_s = (V_0 - \theta)e^{-\kappa s} + \theta + \gamma \int_0^s e^{-\kappa(s-t)} dW_t^{(2)}.$$

We can compute

$$e_s := \mathbb{E} [V_s] = (V_0 - \theta)e^{-\kappa s} + \theta, \quad (51)$$

$$v(s) := \text{Var} [V_s] = \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa s}), \quad (52)$$

and the higher moments can also be computed

$$\mathbb{E} [V_s^2] = e_s^2 + v(s). \quad (53)$$

$$\mathbb{E} [V_s^3] = e_s^3 + 3e_s v(s). \quad (54)$$

$$\mathbb{E} [V_s^4] = e_s^4 + 6e_s^2 v(s) + 3v^2(s). \quad (55)$$

For $s \leq u$, $\mathbb{E} [V_u | V_s] = \mathbb{E} [(V_s - \theta)e^{-\kappa(u-s)} + \theta]$, and

$$\begin{aligned} \mathbb{E} [V_s V_u] &= \mathbb{E} [V_s \mathbb{E} [V_u | V_s]] \\ &= \mathbb{E} [V_s ((V_s - \theta)e^{-\kappa(u-s)} + \theta)] \\ &= e^{-\kappa(u-s)} \mathbb{E} [V_s^2] + \theta(1 - e^{-\kappa(u-s)}) \mathbb{E} [V_s]. \end{aligned}$$

Now we can compute the continuous fair strike as

$$\begin{aligned} K_c &= \frac{1}{T} \mathbb{E} \left[\int_0^T V_s^2 ds \right] = \frac{1}{T} \int_0^T \left[((V_0 - \theta)e^{-\kappa s} + \theta)^2 + \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa s}) \right] ds \\ &= \left((V_0 - \theta)^2 - \frac{\gamma^2}{2\kappa} \right) \frac{1 - e^{-2\kappa T}}{2\kappa T} + 2\theta(V_0 - \theta) \frac{1 - e^{-\kappa T}}{\kappa T} + \theta^2 + \frac{\gamma^2}{2\kappa}. \quad (56) \end{aligned}$$

For $s \leq u$

$$\begin{aligned}
\mathbb{E} [V_s^2 V_u^2] &= \mathbb{E} [V_s^2 \mathbb{E} [V_u^2 | V_s]] \\
&= \mathbb{E} \left[V_s^2 \left(((V_s - \theta)e^{-\kappa(u-s)} + \theta)^2 + \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa(u-s)}) \right) \right] \\
&= e^{-2\kappa(u-s)} \mathbb{E} [V_s^4] + 2\theta e^{-\kappa(u-s)}(1 - e^{-\kappa(u-s)}) \mathbb{E} [V_s^3] \\
&\quad + \left(\theta^2(1 - e^{-\kappa(u-s)})^2 + \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa(u-s)}) \right) \mathbb{E} [V_s^2]. \\
\mathbb{E} [V_s V_u^2] &= \mathbb{E} [V_s \mathbb{E} [V_u^2 | V_s]] \\
&= \mathbb{E} \left[V_s \left(((V_s - \theta)e^{-\kappa(u-s)} + \theta)^2 + \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa(u-s)}) \right) \right] \\
&= e^{-2\kappa(u-s)} \mathbb{E} [V_s^3] + 2\theta e^{-\kappa(u-s)}(1 - e^{-\kappa(u-s)}) \mathbb{E} [V_s^2] \\
&\quad + \left(\theta^2(1 - e^{-\kappa(u-s)})^2 + \frac{\gamma^2}{2\kappa}(1 - e^{-2\kappa(u-s)}) \right) \mathbb{E} [V_s]. \\
\mathbb{E} [V_s^2 V_u] &= \mathbb{E} [V_s^2 \mathbb{E} [V_u | V_s]] \\
&= \mathbb{E} \left[V_s^2 ((V_s - \theta)e^{-\kappa(u-s)} + \theta) \right] \\
&= e^{-\kappa(u-s)} \mathbb{E} [V_s^3] + \theta(1 - e^{-\kappa(u-s)}) \mathbb{E} [V_s^2].
\end{aligned}$$

In the above expressions, the moments $\mathbb{E} [V_s]$, $\mathbb{E} [V_s^2]$, $\mathbb{E} [V_s^3]$ and $\mathbb{E} [V_s^4]$ are already calculated in (51), (53), (54), and (55). Then we can substitute the corresponding inputs into equation (42), sum up the terms, and obtain $K_d^{SZ}(n)$. This completes the proof. \square

References

- ALBRECHER, H., P. MEYER, W. SHOUTENS, AND J. TISTAERT (2007): “The little Heston trap,” *Wilmott Magazine*, pp. 83–92.
- BENTH, F., M. GROTH, AND R. KUFAKUNESU (2007): “Valuing volatility and variance swaps for a non-Gaussian Ornstein-Uhlenbeck stochastic volatility model,” *Applied Mathematical Finance*, 14(4), 347–363.
- BERNARD, C., AND Z. CUI (2011): “Pricing Timer Options,” *Journal of Computational Finance*, 15(1), 69–104.

- BROADIE, M., AND A. JAIN (2008a): “The Effect of Jumps and Discrete Sampling on Volatility and Variance Swaps,” *International Journal of Theoretical and Applied Finance*, 11(8), 761–797.
- (2008b): “Pricing and hedging volatility derivatives,” *Journal of Derivatives*, 15(3), 7–24.
- BROADIE, M., AND Ö. KAYA (2006): “Exact Simulation of Stochastic Volatility and other Affine Jump Diffusion Processes,” *Operations Research*, 54(2), 217–231.
- BÜHLER, H. (2006): “Volatility markets: consistent modelling, hedging and practical implementation,” *Ph.D. dissertation, Technical Universität Berlin*.
- CAI, N., AND S. KOU (2011): “Option pricing under a mixed-exponential jump diffusion model,” *Management Science*, 57(11), 2067–2081.
- CARR, P., AND R. LEE (2007): “Realized Volatility and Variance Options via Swaps,” *Risk*, May, 76–83.
- (2009): “Volatility Derivatives,” *Annual Review of Financial Economics*, 1, 319–339.
- CARR, P., R. LEE, AND L. WU (2012): “Variance swaps on Time-changed Lévy processes,” *Finance and Stochastics*, 16(2), 335–355.
- CARR, P., AND D. MADAN (1998): “Towards a theory of volatility trading,” *Jarrow, R. (ed) Volatility, Risk publications, London*, pp. 417–427.
- DUPIRE, B. (1993): “Model art,” *Risk*, 6, 118–124.
- HESTON, S. (1993): “A Closed-form Solution for Options with Stochastic Volatility with Applications to Bond and Currency Options.,” *The Review of Financial Studies*, 6, 327–343.
- HÖRFELT, P., AND O. TORNÉ (2010): “The Value of a Variance Swap - a Question of interest,” *Risk*, June, 82–85.
- HOWISON, S., A. RAFAILIDIS, AND H. RASMUSSEN (2004): “On the pricing and hedging of volatility derivatives,” *Applied Mathematical Finance*, 11(4), 317–346.
- HULL, J., AND A. WHITE (1987): “The Pricing of Options on Assets with Stochastic Volatilities,” *Journal of Finance*, 42(2), 281–300.
- HURD, T., AND A. KUZNETSOV (2008): “Explicit formulas for Laplace transform of stochastic integrals,” *Markov Processes and Related Fields*, 14, 277–290.

- ITKIN, A., AND P. CARR (2010): “Pricing swaps and options on quadratic variation under stochastic time change models—discrete observations case,” *Review of Derivatives Research*, 13(1), 141–176.
- JARROW, R., Y. KCHIA, M. LARSSON, AND P. PROTTER (2013): “Discretely Sampled Variance and Volatility Swaps versus their Continuous Approximations,” *Finance and Stochastics*, 17(2), 305–324.
- JEANBLANC, M., M. YOR, AND M. CHESNEY (2009): *Mathematical Methods for Financial Markets*.
- KARATZAS, I., AND S. SHREVE (1991): “Brownian Motion and Stochastic Calculus,” *Graduate Texts in Mathematics*. vol. 113, 2nd edn. Springer, New York.
- KELLER-RESSEL, M., AND C. GRIESSLER (2012): “Convex order of discrete, continuous and predictable quadratic variation and applications to options on variance,” *Working paper, available at: <http://arxiv.org/abs/1103.2310>*.
- KELLER-RESSEL, M., AND J. MUHLE-KARBE (2012): “Asymptotic and exact pricing of options on variance,” *Finance and Stochastics*, forthcoming.
- LEE, R. (2010): “Gamma Swap,” *Encyclopedia of Quantitative Finance*, Edited by Rama Cont.
- NEUBERGER, A. (1994): “The Log Contract,” *Journal of Portfolio Management*, 6, 118–124.
- SCHÖBEL, R., AND J. ZHU (1999): “Stochastic volatility with an Ornstein-Uhlenbeck process: An extension,” *European Finance Review*, 3(1), 23–46.
- SEPP, A. (2012): “Pricing options on realized variance in the Heston model with jumps in returns and volatility II: An approximate distribution of the discrete variance,” *Journal of Computational Finance*, forthcoming.
- WINDCLIFF, H., P. FORSYTH, AND K. VETZAL (2006): “Pricing methods and hedging strategies for volatility derivatives,” *Journal of Banking and Finance*, 30, 409–431.
- ZHU, S., AND G. LIAN (2011): “A Closed-form Exact Solution for Pricing Variance Swaps with Stochastic Volatility,” *Mathematical Finance*, 21, 233–256.




Figure 1: Sensitivity to the number of sampling periods n and to ρ

Parameters correspond to Set 1 in Table 1 except for ρ that can take three possible values $\rho = -0.7$, $\rho = 0$ or $\rho = 0.7$ and for T that is equal to $T = 1$ for the two upper graphs and $T = 1/12$ for the two lower graphs. When $T = 1/12$, the parameters for the Hull-White model are adjusted according to the procedure described in Section 7.1. In the case when $T = 1/12$, one has $\mu = 4.03$ and $\sigma = 1.78$.




Figure 2: Sensitivity to the number of sampling periods n and to γ

Parameters are set to unusual values to show that any types of behaviors can be expected.
 $\rho = 0.6$, $r = 3.19\%$, $\theta = 0.019$, $\kappa = .1$, $V_0 = 0.8$ and γ takes three possible values: 0.5, 1.5 and 2.




Figure 3: Asymptotic expansion $K_c^M + a/n$ with respect to the number of sampling periods n and to ρ

Parameters correspond to Set 2 in Table 1 except for ρ that can take three possible values $\rho = -0.3$, $\rho = 0$ or $\rho = 0.3$. The upper graphs correspond to large number of discretization steps whereas lower graphs have relatively small values of n .




Figure 4: Asymptotic expansion with respect to the correlation coefficient ρ and the risk-free rate r

Parameters correspond to Set 1 in Table 1 except for r that can take three possible values $r = 0\%$, $r = 3.2\%$ or $r = 6\%$. Here $n = 250$, which corresponds to a daily monitoring as $T = 1$.




Figure 5: Discrete and continuous fair strikes with respect to the maturity T and to V_0

Parameters correspond to Set 2 in Table 1 except for T and V_0 . Also we choose a monthly monitoring to compute the discrete fair strike. When $\theta = V_0$, K_c^H is independent of the maturity T .




Figure 6: Discrete and continuous fair strikes with respect to the parameter γ and to V_0

Parameters correspond to Set 2 in Table 1 except for γ and V_0 that are indicated on the graphs. A monthly monitoring is used to compute the discrete fair strike. The continuous fair strike K_c^H is independent of γ , so that it is easy to identify the different curves on the graph.




Figure 7: Sensitivity to the number of sampling periods n and to ρ

Parameters are similar to Set 1 in Table 1 for the Heston model except for ρ that can take three possible values $\rho = -0.7$, $\rho = 0$ or $\rho = 0.7$ and for T that is equal to $T = 1$ for the left panel and $T = 1/12$ for the right panel. Precisely, we use the following parameters for the Schöbel-Zhu model. $\kappa = 6.21$, $\theta = \sqrt{0.019}$, $\gamma = 0.31$, $r = 0.0319$, $V_0 = \sqrt{0.010201}$.




Figure 8: Asymptotic expansion with respect to the correlation coefficient ρ and the risk-free rate r .

Parameters are similar to Set 1 in Table 1 for the Heston model except for r that can take three possible values $r = 0\%$, $r = 3.2\%$ or $r = 6\%$. Precisely, we use the following parameters for the Schöbel-Zhu model: $\kappa = 6.21$, $\theta = \sqrt{0.019}$, $\gamma = 0.31$, $\rho = -0.7$, $T = 1$, $V_0 = \sqrt{0.010201}$. Here $n = 250$, which corresponds to a daily monitoring as $T = 1$.