

# Distributed Nash Equilibrium Seeking For Second-Order Systems With Unknown Dynamics

1

Abstract—1

Index Terms—1

## I. INTRODUCTION

OVER the past two decades, game theory has spread across diverse research fields, including biology, economics, and computer science. As game theory evolves, the quest for Nash equilibrium in noncooperative games gains significance in theory and practice, as evidenced by studies [1], [2], [3], [4], [5], [6]. Substantial progress has been made in distributed control and optimization of networked systems based on Nash equilibrium seeking and its applications [7], [8], [9], [10], [11]. For example, building upon the framework introduced in [12], numerous consensus-based distributed Nash equilibrium seeking strategies have been devised, such as distributed Nash equilibrium seeking in multiagent game under switching communication topologies digraph [13] and fully distributed Nash equilibrium seeking [14]. Nevertheless, the majority of existing findings don't account for the influence of system disturbances, which is unrealistic considering that many practical engineering systems frequently encounter disturbances.

## II. NOTATIONS AND PRELIMINARIES

Notations: The real number set is denote as  $\mathbb{R}$ .  $\|z\|$  denotes the  $\ell_2$ -norm of  $z$ .  $[z_i]_{vec}$  where  $i \in \{1, 2, \dots, N\}$  is defined as a column vector whose dimension is  $N \times 1$  and the  $i$ th element is  $z_i$ .  $\text{diag}\{k_i\}$  for  $i \in \{1, 2, \dots, N\}$  is a diagonal matrix whose dimension is  $N \times N$  and the  $i$ th diagonal element is  $k_i$ .  $\text{diag}\{a_{ij}\}$  where  $i, j \in \{1, 2, \dots, N\}$  gives a diagonal matrix whose dimension is  $N^2 \times N^2$  and diagonal elements are  $a_{11}, a_{12}, \dots, a_{1N}, a_{21}, \dots, a_{NN}$ , successively.  $\mathcal{A} = [a_{ij}]$  is a matrix whose  $(i, j)$ th entry is  $a_{ij}$ . Given that matrix  $Q$  is symmetric and real,  $\lambda_{\min}(Q)$  ( $\lambda_{\max}(Q)$ ) stands for the smallest (largest) eigenvalue of  $Q$ .  $\max_{i \in \{1, 2, \dots, N\}} \{l_i\}$  denotes the largest value of  $l_i$  for  $i \in \{1, 2, \dots, N\}$ .  $\mathbf{I}_{N \times N}$  is an identity matrix with its dimension being  $N \times N$  and  $\mathbf{1}(0)$  is a column vector with its entries being 1(0). Moreover,  $\otimes$  is the Kronecker product.

Algebraic Graph Theory: A graph  $\mathcal{G}$  is given by  $\mathcal{G} = (\mathcal{V}, \mathcal{E}_g)$ , in which  $\mathcal{V} = \{1, 2, \dots, N\}$ ,  $\mathcal{E}_g \subseteq \mathcal{V} \times \mathcal{V}$  respectively are the node set and edge set. The edge  $(i, j) \in \mathcal{E}_g$  indicates are the node  $j$  can receive information from node  $i$ , but not necessarily vice versa. The in-neighbor set of node  $i$  is given as  $\mathcal{N}_i^{\text{in}} = \{j | (j, i) \in \mathcal{E}_g\}$ . A directed path is a

sequence of edges of the form  $(i_1, i_2), (i_2, i_3), \dots$ . A directed graph is strongly connected if for every pair of two distinct nodes, there is a path. Let  $\mathcal{A} = [a_{ij}]$  be the adjacency matrix in which  $a_{ij} > 0$  if  $(i, j) \in \mathcal{E}_g$  and  $a_{ij} = 0$  otherwise. The Laplacian matrix  $\mathcal{L}$  is defined as  $\mathcal{L} = \mathcal{D} - \mathcal{A}$ , where  $\mathcal{D} = \text{diag}\{d_i\}$  and  $d_i = \sum_{j=1}^N a_{ij}$ .

## III. PROBLEM STATEMENT

In the concerned game,  $N$  players with labels from 1 to  $N$  are engaged and each player  $i$  has a local objective function  $f_i(\mathbf{x}) : \mathbb{R}^N \rightarrow \mathbb{R}$ , in which  $\mathbf{x} = [x_1, x_2, \dots, x_N]^T$  and  $x_i \in \mathbb{R}$  is the action of player  $i$ . Moreover, for second-order players, player  $i$ 's action is governed by

$$\begin{cases} \dot{x}_i(t) = v_i(t) \\ \dot{v}_i(t) = u_i + d_i(t) \end{cases} \quad (1)$$

where  $v_i(t)$  is the velocity-like state of player  $i$ ,  $u_i$  is the control input and  $d_i(t)$  is the disturbance for  $i \in \mathcal{V}$ . Each player  $i$  aims to minimize its own objective function  $f_i(\mathbf{x})$  by adjusting its action  $x_i(t)$ , that is

$$\begin{aligned} \min_{x_i} \quad & f_i(\mathbf{x}) \\ \text{s.t.} \quad & (1) \text{ for second-order players} \end{aligned} \quad (2)$$

Definition 1: An action profile  $\mathbf{x}^* = (x_i^*, \mathbf{x}_{-i}^*)$  is a Nash equilibrium if for all  $i \in \mathcal{V}$ , we have

$$f_i(x_i^*, \mathbf{x}_{-i}^*) \leq f_i(x_i, \mathbf{x}_{-i}^*), \forall x_i \in \mathbb{R} \quad (3)$$

where  $\mathbf{x}_{-i} = [x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_N]^T$ .

To facilitate later analysis, we made the following assumptions.

Assumptions 1: For  $i \in \mathcal{V}$ ,  $f_i(\mathbf{x})$  is  $\mathcal{C}^2$  and  $\nabla f_i(\mathbf{x})$  is globally Lipschitz with  $l_i$ .

Assumptions 2: The digraph  $\mathcal{G}$  is strongly connected.

Lemma 1: Let  $D$  be a nonnegative diagonal matrix and  $\mathcal{H} = \mathcal{L} + D$ . Under Assumptions 2, there are symmetric positive definite matrices  $P$  and  $Q$  such that

$$\mathcal{H}^T P + P \mathcal{H} = Q \quad (4)$$

Assumptions 3: For  $\mathbf{x}, \mathbf{z} \in \mathbb{R}^N$ ,

$$(\mathbf{x} - \mathbf{z})^T ([\nabla f_i(\mathbf{x})]_{vec} - [\nabla f_i(\mathbf{z})]_{vec}) \geq m \|\mathbf{x} - \mathbf{z}\|^2 \quad (5)$$

where  $\nabla f_i(\mathbf{x}) = \partial f_i(\mathbf{x}) / \partial x_i$  and  $m$  is a positive constant.

#### IV. MAIN RESULTS

Assumptions 4: For  $i \in \mathcal{V}$ ,  $\nabla_{ij}^2 f_i(\mathbf{x}) = \partial^2 f_i(\mathbf{x}) / \partial x_i \partial x_j$  is bounded.

Assumptions 5: The disturbance  $d(t)$  is bounded, i.e.,  $\forall i \in \mathcal{V}, |d_i(t)| \leq \bar{d}_i(t)$  for a positive constant  $\bar{d}_i$ .

To realize asymptotic Nash equilibrium seeking for games with second-order integrator-type players distributively, the control input is designed as

$$\begin{cases} u_i = -\tau k_i s_i - \beta_i \xi_i(t) \\ s_i = v_i + \nabla_i f_i(\mathbf{y}_i) \\ \dot{y}_{ij} = -\theta \bar{\theta}_{ij} \left( \sum_{k=1}^N a_{ik}(y_{ij} - y_{kj}) + a_{ij}(y_{ij} - x_j) \right) \\ \dot{\varrho}_i(t) = -\alpha_i \varrho_i(t) \text{ with } \varrho_i(t) > 0 \end{cases} \quad (6)$$

where  $i, j \in \mathcal{V}$ ,  $\theta, \tau$  are adjustable positive parameters,  $\bar{\theta}_{ij}, k_i, \alpha_i$  are fixed positive parameters,  $\beta_i \geq \bar{d}_i$  is a positive constant,  $s_i$  and  $\varrho_i(t)$  represents an auxiliary variable for player  $i$  and  $\xi_i(t) = s_i(t) / (|s_i(t)| + \varrho_i(t))$ .

By (1) and (6), the closed-loop system is

$$\begin{cases} \dot{\mathbf{x}} = \mathbf{v} \\ \dot{\mathbf{v}} = -\tau \mathbf{k} \mathbf{s} - [\beta_i \xi_i(t)]_{vec} + \mathbf{d}(t) \\ \mathbf{s} = \mathbf{v} + [\nabla_i f_i(\mathbf{y}_i)]_{vec} \\ \dot{\mathbf{y}} = -\theta \bar{\theta} (\mathcal{L} \otimes \mathbf{I}_{N \times N} + \mathcal{A}_q) \tilde{\mathbf{y}} \end{cases} \quad (7)$$

where  $\mathbf{k} = \text{diag}\{k_i\}$ ,  $\bar{\theta} = \text{diag}\{\bar{\theta}_{ij}\}$ ,  $\mathcal{A}_q = \text{diag}\{a_{ij}\}$ ,  $\mathbf{y} = [\mathbf{y}_1^T, \mathbf{y}_2^T, \dots, \mathbf{y}_N^T]^T$  and  $\tilde{\mathbf{y}} = \mathbf{y} - \mathbf{1}_N \otimes \mathbf{x}$ .

Let  $\tilde{\mathbf{x}} = \mathbf{x} - \mathbf{x}^*$ .

Theorem 1: Under Assumptions 1-5, there exist a  $\theta^* > 0$  such that for each  $\theta > \theta^*$ , there exists a  $\tau^* > 0$  such that for each  $\tau > \tau^*$ , players' actions globally exponentially converge to the Nash equilibrium by (7).

Proof: Define  $V = V_1 + V_2 + V_3$ , where

$$\begin{aligned} V_1(t) &= \frac{1}{2} \tilde{\mathbf{x}}^T \tilde{\mathbf{x}} \\ V_2(t) &= \frac{1}{2} \tilde{\mathbf{y}}^T P \tilde{\mathbf{y}} \\ V_3(t) &= \frac{1}{2} \mathbf{s}^T \mathbf{s} + \sum_{i=1}^N \frac{\beta_i}{\alpha_i} \varrho_i(t) \end{aligned} \quad (8)$$

Then, taking the time derivative of  $V_1$  yields

$$\begin{aligned} \dot{V}_1 &= \tilde{\mathbf{x}}^T (\mathbf{s} - [\nabla_i f_i(\mathbf{y}_i)]_{vec}) \\ &\leq -m \|\tilde{\mathbf{x}}\|^2 + \max_{i \in \mathcal{V}} \{l_i\} \|\tilde{\mathbf{x}}\| \|\tilde{\mathbf{y}}\| + \|\tilde{\mathbf{x}}\| \|\mathbf{s}\| \end{aligned} \quad (9)$$

where in the inequality, we have used Assumptions 3 and  $\|[\nabla_i f_i(\mathbf{y}_i)]_{vec} - [\nabla_i f_i(\mathbf{x})]_{vec}\| \leq \max_{i \in \mathcal{V}} \{l_i\} \|\tilde{\mathbf{y}}\|$ .

Then, taking the time derivative of  $V_2$  yields

$$\begin{aligned} \dot{V}_2 &= \dot{\tilde{\mathbf{y}}}^T P \tilde{\mathbf{y}} + \tilde{\mathbf{y}}^T P \dot{\tilde{\mathbf{y}}} \\ &= -\theta \tilde{\mathbf{y}}^T Q \tilde{\mathbf{y}} - 2\tilde{\mathbf{y}}^T P \mathbf{1}_N \otimes \mathbf{s} \\ &\quad + 2\tilde{\mathbf{y}}^T P \mathbf{1}_N \otimes [\nabla_i f_i(\mathbf{y}_i)]_{vec} \\ &\leq -(\theta \lambda_{\min}(Q) - 2\sqrt{N} \|P\| \max_{i \in \mathcal{V}} \{l_i\}) \|\tilde{\mathbf{y}}\|^2 \\ &\quad + 2\sqrt{N} \|P\| \|\tilde{\mathbf{y}}\| \|\mathbf{s}\| + 2N \|P\| \max_{i \in \mathcal{V}} \{l_i\} \|\tilde{\mathbf{y}}\| \|\tilde{\mathbf{x}}\|. \end{aligned} \quad (10)$$

where in the inequality, by Assumption 2 and Lemma 1, we let  $P, Q \in \mathbb{R}^{N^2 \times N^2}$  and  $Q = P\bar{\theta}(\mathcal{L} \otimes \mathbf{I}_{N \times N} + \mathcal{A}_q) + (\mathcal{L} \otimes \mathbf{I}_{N \times N} + \mathcal{A}_q)^T \bar{\theta} P = Q$ .

Moreover, taking the time derivative of  $V_3$  yields

$$\begin{aligned} \dot{V}_3 &= \mathbf{s}^T (\dot{\mathbf{v}} + \bar{H}(\mathbf{y}) \dot{\mathbf{y}}) - \sum_{i=1}^N \beta_i \varrho_i(t) \\ &= \mathbf{s}^T (-\tau \mathbf{k} \mathbf{s} - [\beta_i \xi_i(t)]_{vec} + \mathbf{d}(t) + \bar{H}(\mathbf{y}) \dot{\mathbf{y}}) - \sum_{i=1}^N \beta_i \varrho_i(t) \end{aligned} \quad (11)$$

where  $\bar{H}(\mathbf{y}) = [h_{ij}]$  in which for  $i \neq j$ ,  $h_{ij} = \mathbf{0}_N^T$  and for  $i = j$ ,  $h_{ii} = [\nabla_{i1}^2 f_i(\mathbf{y}_i), \nabla_{i2}^2 f_i(\mathbf{y}_i), \dots, \nabla_{iN}^2 f_i(\mathbf{y}_i)] \nabla_{ij}^2 f_i(\mathbf{y}_i) = \frac{\partial^2 f_i(\mathbf{x})}{\partial x_i \partial x_i} |_{\mathbf{x}=\mathbf{y}_i}$ . By Assumption 4,  $\|\bar{H} \mathbf{y}\|$  is bounded. Hence, There are some positive constant  $L_1$  meet  $\|\bar{H}(\mathbf{y})\| \|\bar{\theta}(\mathcal{L} \otimes \mathbf{I}_{N \times N} + \mathcal{A}_q)\| \leq L_1$ . Therefore,

$$\begin{aligned} \dot{V}_3 &\leq -\tau \lambda_{\min}(\mathbf{k}) \|\mathbf{s}\|^2 + \theta L_1 \|\mathbf{s}\| \|\tilde{\mathbf{y}}\| \\ &\quad - \sum_{i=1}^N \frac{(\beta_i - |d_i(t)|) |s_i(t)| (|s_i(t)| + \varrho_i(t)) + \beta_i \varrho_i^2(t)}{|s_i(t)| + \varrho_i(t)} \end{aligned} \quad (12)$$

Hence,

$$\begin{aligned} \dot{V} &\leq -m \|\tilde{\mathbf{x}}\|^2 - \tau \lambda_{\min}(\mathbf{k}) \|\mathbf{s}\|^2 \\ &\quad - (\theta \lambda_{\min}(Q) - 2\sqrt{N} \|P\| \max_{i \in \mathcal{V}} \{l_i\}) \|\tilde{\mathbf{y}}\|^2 \\ &\quad + (\max_{i \in \mathcal{V}} \{l_i\} + 2N \|P\| \max_{i \in \mathcal{V}} \{l_i\}) \|\tilde{\mathbf{x}}\| \|\tilde{\mathbf{y}}\| \\ &\quad + (2\sqrt{N} \|P\| + \theta L_1) \|\tilde{\mathbf{y}}\| \|\mathbf{s}\| + \|\tilde{\mathbf{x}}\| \|\mathbf{s}\| \\ &\quad - \sum_{i=1}^N \frac{(\beta_i - |d_i(t)|) |s_i(t)| (|s_i(t)| + \varrho_i(t)) + \beta_i \varrho_i^2(t)}{|s_i(t)| + \varrho_i(t)} \\ &\leq -\lambda_{\min}(F) \|\mathbf{E}_1\|^2 - \tau \lambda_{\min}(\mathbf{k}) \|\mathbf{s}\|^2 \\ &\quad + (1 + 2\sqrt{N} \|P\| + \theta L_1) \|\mathbf{E}_1\| \|\mathbf{s}\| \end{aligned} \quad (13)$$

where  $F = \begin{bmatrix} m & a \\ a & \theta \lambda_{\min}(Q) - 2\sqrt{N} \|P\| \max_{i \in \mathcal{V}} \{l_i\} \end{bmatrix}$ ,  $\mathbf{E}_1 = [\tilde{\mathbf{x}}^T, \tilde{\mathbf{y}}^T]^T$ ,  $a = -\frac{\max_{i \in \mathcal{V}} \{l_i\} + 2N \|P\| \max_{i \in \mathcal{V}} \{l_i\}}{2}$ , choose  $\theta > \frac{(\max_{i \in \mathcal{V}} \{l_i\} + 2N \|P\| \max_{i \in \mathcal{V}} \{l_i\})^2}{4m \lambda_{\min}(Q)} + \frac{2\sqrt{N} \|P\| \max_{i \in \mathcal{V}} \{l_i\}}{\lambda_{\min}(Q)}$ . Then,  $\lambda_{\min}(F)$  is a real positive number and  $\lambda_{\min}(F) > 0$ . Thus,

$$\dot{V}(t) \leq -\lambda_{\min}(G) \|\mathbf{E}\|^2 \quad (14)$$

where  $G = \begin{bmatrix} \lambda_{\min}(F) & -\frac{1+2\sqrt{N} \|P\| + \theta L_1}{2} \\ -\frac{1+2\sqrt{N} \|P\| + \theta L_1}{2} & \tau \lambda_{\min}(\mathbf{k}) \end{bmatrix}$ ,  $\mathbf{E} = [\tilde{\mathbf{x}}^T, \tilde{\mathbf{y}}^T, \tilde{\mathbf{s}}^T]^T$ . By choosing  $\tau > \frac{(1+2\sqrt{N} \|P\| + \theta L_1)^2}{4\lambda_{\min}(F) \lambda_{\min}(\mathbf{k})}$ ,  $\lambda_{\min}(G)$  is a real positive number and  $\lambda_{\min}(G) > 0$ .

The conclusion can be drawn.

#### V. SIMULATION STUDIES

#### VI. CONCLUSIONS

#### References

- [1] David Pozo and Javier Contreras. Finding multiple nash equilibria in pool-based markets: A stochastic epec approach. IEEE Transactions on Power Systems, 26(3):1744–1752, 2011.

- [2] Lillian J Ratliff, Samuel A Burden, and S Shankar Sastry. On the characterization of local nash equilibria in continuous games. *IEEE transactions on automatic control*, 61(8):2301–2307, 2016.
- [3] Youcheng Lou, Yiguang Hong, Lihua Xie, Guodong Shi, and Karl Henrik Johansson. Nash equilibrium computation in subnetwork zero-sum games with switching communications. *IEEE Transactions on Automatic Control*, 61(10):2920–2935, 2015.
- [4] Miloš S Stankovic, Karl H Johansson, and Dušan M Stipanovic. Distributed seeking of nash equilibria with applications to mobile sensor networks. *IEEE Transactions on Automatic Control*, 57(4):904–919, 2011.
- [5] Shu-Jun Liu and Miroslav Krstić. Stochastic nash equilibrium seeking for games with general nonlinear payoffs. *SIAM Journal on Control and Optimization*, 49(4):1659–1679, 2011.
- [6] Hans-Bernd Dürr, Miloš S Stanković, Christian Ebenbauer, and Karl Henrik Johansson. Lie bracket approximation of extremum seeking systems. *Automatica*, 49(6):1538–1552, 2013.
- [7] Lei Ding, Qing-Long Han, Boda Ning, and Dong Yue. Distributed resilient finite-time secondary control for heterogeneous battery energy storage systems under denial-of-service attacks. *IEEE Transactions on Industrial Informatics*, 16(7):4909–4919, 2019.
- [8] Jiahui Wu, Yunru Zhu, Yuanshi Zheng, and Huaizhu Wang. Resilient bipartite consensus of second-order multiagent systems with event-triggered communication. *IEEE Systems Journal*, 2021.
- [9] Frank L Lewis, Hongwei Zhang, Kristian Hengster-Movric, and Abhijit Das. Cooperative control of multi-agent systems: optimal and adaptive design approaches. Springer Science & Business Media, 2013.
- [10] Xiaohua Ge, Shunyuan Xiao, Qing-Long Han, Xian-Ming Zhang, and Derui Ding. Dynamic event-triggered scheduling and platooning control co-design for automated vehicles over vehicular ad-hoc networks. *IEEE/CAA Journal of Automatica Sinica*, 9(1):31–46, 2021.
- [11] Maojiao Ye, Qing-Long Han, Lei Ding, Shengyuan Xu, and Guobiao Jia. Distributed nash equilibrium seeking strategies under quantized communication. *IEEE/CAA Journal of Automatica Sinica*, 2022.
- [12] Maojiao Ye and Guoqiang Hu. Distributed nash equilibrium seeking by a consensus based approach. *IEEE Transactions on Automatic Control*, 62(9):4811–4818, 2017.
- [13] Maojiao Ye and Guoqiang Hu. Distributed nash equilibrium seeking in multiagent games under switching communication topologies. *IEEE Transactions on Cybernetics*, 48(11):3208–3217, 2018.
- [14] Maojiao Ye and Guoqiang Hu. Adaptive approaches for fully distributed nash equilibrium seeking in networked games. *Automatica*, 129:109661, 2021.