


General Theory Of Relativity and Cosmology

LECTURE NOTES

Instructor: Prof. Golam Mortuza Hossain



Sagnik Seth

Dept of Physical Sciences
IISER Kolkata

Contents

Lecture 01: Introduction	4
Lecture 02: Galilean Stuffs	5
2.1 Galilean Transformation	5
2.2 A Big Blow to Galileo	7
Lecture 03: Lorentz Transformation	7
3.1 What happens to physical laws?	8
3.2 Classification of Intervals	9
Lecture 04: Vectors and Stuffs	9
Lecture 05: Some more Vectors and Stuffs	11
5.1 Gradient Vector	11
5.2 Tensors	12
5.3 Inner Product	13
Lecture 06: Metric is also a Matrix	14
6.1 Transformation of the metric	14
6.2 Metric is a map!	15
Lecture 07: Metric and STR	15
7.1 Proper Time	16
7.2 Looking back at STR	16
7.2.1 Metric in STR	17
Lecture 08: May the Force be with you!	17
8.1 Emergence of Energy	17
8.2 The many faces of Mass	18
8.2.1 Eötvös Experiment	19
Lecture 09: Einstein's thoughts	19
9.1 How's gravity different?	20
9.2 Gedankenexperiment	20
9.2.1 Lift in free space:	21
9.3 Lift under Gravity	21
Lecture 10: Connections are important	23
10.1 Action tells a lot!	26
Lecture 11: Finding Geodesics	27
Lecture 12: Towards the action for Gravity: Invariant Volumes	30
12.1 Maxwell's action in spacetime	30
12.2 Metric, Transformations and Jacobian	32
Lecture 13: Towards the action for Gravity: Covariant Derivatives	33
Lecture 14: Towards the action for Gravity: Riemann Tensor	35
Lecture 15: Curvatures	39

Lecture 16: Bianchi Identities	41
16.1 Natural Units	41
16.2 First Bianchi Identity	42
16.3 Second Bianchi Identity	43

Lecture 01: Introduction

In the physical sense, the word 'relativity' can be used to relate the measurement of one observer with that of another, for the same object/quantity. So using relativity, we can characterise the difference in the observations of two observers in two different frame of reference.

Note that, physical laws should not matter based on who is observing (that is, physical laws are *universal*), hence relativity becomes very important when dealing with the resolution of conflicts.

Let us consider two reference frames, and let v be the relative velocity between the frames.





Figure 1: Two frames of reference with a relative velocity v

Fig. 1 shows two reference frames. The moving frame is denoted by O' (henceforth, primes will generally denote moving frames). If the relative velocity v between the two frames remain constant, then it falls under the domain of *Special Relativity* and if unfortunately (or fortunately) it doesn't, then it comes under *General Relativity*.


The theory of relativity dates back to the inception of Newtonian mechanics and hence, let us review the laws in brief (without being much technical 😊):

- **The First Law** $\approx$ velocity of a body remains constant unless acted by an external, unbalanced force.
- **The Second Law** $\approx \vec{F} = m\vec{a}$ where $\vec{F}$ is the external force and $\vec{a}$ is the acceleration.
- **The Third Law** $\approx$ Every action has an equal and opposite reaction. Note that for this, two bodies are needed.



From the figure above, we will have $\vec{F}_{21} = -\vec{F}_{12}$

- **The Gravitational Law** $\approx$



In the above situation, if we consider for mass m_1 , we have:

$$\vec{F}_{12} = G \frac{m_1 m_2}{r_{21}^2} \hat{r}_{21}$$

An important observation: in the second law, if we take the force to be zero, then apparently $\vec{a} = \frac{d\vec{v}}{dt} = 0 \implies \vec{v} = \text{const.}$ which is the statement of the first law. However, the second law does not imply the first law, since the first law defines what an *inertial frame* is and the statement of the second law applies only in case of inertial frames. So, a better formulation of the second and third law can be like, "In a frame where the first law is valid, blah blah blah..."

Lecture 02: Galilean Stuffs

In the previous lecture, we saw how Newton's laws define an inertial frame and how we can rephrase the second and third law in terms of inertial frames, to avoid confusion. It happens that, there are reference frames where the first law doesn't hold true.

Imagine a person inside a darkened car, isolated from the outside world, with a ball hanging from the roof of the car. The car suddenly starts accelerating and (as intuition speaks) the ball starts moving. The person claims "The ball has moved!" 🧠.

Note that from the perspective of the person, no force has been applied to the ball, yet it moved, which is a direct violation of the second law. Thus, we can call this frame *non-inertial*.

Now, consider a person inside an ill-fated lift, tragically falling 😊. Everything in the lift frame is accelerating together under gravity and hence 'weightlessness' (free-fall) occurs. If somehow a coin is also there in the lift, it simply floats. And if the person pushes the coin, it moves with an almost constant velocity. Thus, it is a very good approximation to an *inertial frame* (although, extremely tragic for the person!).




Figure 2: A sad guy floating with a coin in a falling lift

We had always wanted the physical laws to stay the same. This is actually the statement of Galilean relativity:

"Laws of physics (nature) must be the same in all inertial frames of reference"

By physical laws, we imply Newton's laws of mechanics and the law of gravitation. To have a more mathematical (and less philosophical) aspect to the above statement, we define the *Galilean transformation*.

2.1 Galilean Transformation

Let an object be at point P and consider two reference frames O and O' , which coincided at $t = 0$ but have moved apart since, along the common x -axis. (Henceforth, reference frames will simply be termed *observers*).




Figure 3: Observing same point from two different frames

Now, the distance between the origins of two frames, at time t , will be $R = vt$. Then accordingly, we can write the relation between the coordinates of the two observers as follows:

$$\begin{aligned} x' &= x - vt \\ y' &= y \\ z' &= z \\ t' &= t \end{aligned} \quad \begin{pmatrix} t' \\ x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -v & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} t \\ x \\ y \\ z \end{pmatrix}$$

An important assumption: The Galilean transformation assumes a *universal time*, that is, time elapsed on clocks in both frames are the same ¹.

Now, it is not necessary for the observers to move along the x axis only. Hence, generalising the transformation to arbitrary direction, we obtain:

$$t' = t \quad \mathbf{r}' = \mathbf{r} - \mathbf{v}t$$

Now, let us look how the physical laws react under the Galilean transformation.

■ Law of Gravitation:

Consider the situation in the diagram below:




Figure 4: Gravitational law from two different frames

According to Galilean transformation, we have:

$$\begin{aligned} \mathbf{z} &= \mathbf{r}_2 - \mathbf{r}_1 \\ &= (\mathbf{r}'_2 + \mathbf{v}t) - (\mathbf{r}'_1 + \mathbf{v}t) \\ &= \mathbf{r}'_2 - \mathbf{r}'_1 \\ &= \mathbf{z}' \end{aligned}$$

¹This was perhaps a fair assumption since time, atleast in the philosophical aspect, has always seemed to be *superior* and all-encompassing, second to none.

Thus, we see that the vector $\hat{\mathbf{r}}$ is unchanged and since this is the only vector appearing in Newton's law of gravitation, we have:

$$\mathbf{F}_{12} = G \frac{m_1 m_2}{r^2} \hat{\mathbf{r}} = G \frac{m_1 m_2}{r'^2} \hat{\mathbf{r}}' = \mathbf{F}'_{12}$$

We see that the expression of the force does not change between frames, thus indicating a *universality*.

■ Second Law:

Using the second law, we can write

$$\mathbf{F} = m \frac{d^2 \mathbf{r}}{dt^2}$$

Now, using Galilean transformation, we obtain:

$$\mathbf{r}(t) = \mathbf{r}'(t') + \mathbf{v}t \implies \frac{d\mathbf{r}}{dt} = \frac{d\mathbf{r}'}{dt} \times \frac{dt'}{dt} + \mathbf{v} \implies m \frac{d^2 \mathbf{r}}{dt^2} = m \frac{d^2 \mathbf{r}'}{dt'^2}$$

Thus, we see that in general, force will be the same for observers in different inertial frames. Note that we assumed the *time universality*, that is, $\frac{dt'}{dt} = 1$

■ Third Law:

Note that since the force expression for gravity is symmetric in m_1 and m_2 , the law of gravitation automatically incorporates the third law. Newton knew only about gravity and perhaps, third law would not have been necessary if he had relied only on gravity, however, he postulated that for all other forces, third law holds.

2.2 A Big Blow to Galileo

Let us consider that in Fig. 3, point P is also moving with some velocity $\mathbf{u}'$ and $\mathbf{u}$ in primed and unprimed frame respectively. Then, we can write:

$$\frac{d\mathbf{u}}{dt} = \frac{d}{dt}(\mathbf{r}' + \mathbf{v}t) = \frac{d\mathbf{r}'}{dt} + \mathbf{v} = \mathbf{u}' + \mathbf{v} \implies \mathbf{u} = \mathbf{u}' + \mathbf{v} \quad : \text{velocity addition formula}$$

However, observations, specially the Michelson-Morley experiment, were made in different frames but found the speed of light (henceforth denoted as c) to be a constant frame-independent quantity, which contradicted the velocity addition formula when applied to speed of light.

Since the velocity addition formula depended entirely on the Galilean transformation, to resolve this conflict, we have to change the transformation rule in its entirety.

Lecture 03: Lorentz Transformation

We saw previously that Galilean relativity failed terribly when speed of light was considered. This needed a change in the transformation rule. The new transformation rule is called the *Lorentz transformation* and is given by (for two frames relatively moving along the common x axis):

$$\begin{aligned} x' &= \gamma(x - vt) \\ y' &= y \\ z' &= z \\ t' &= \gamma\left(t - \frac{vx}{c^2}\right) \end{aligned} \quad \begin{pmatrix} ct' \\ x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} \gamma & -\gamma \frac{v}{c} & 0 & 0 \\ -\gamma \frac{v}{c} & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} ct \\ x \\ y \\ z \end{pmatrix}$$

where $\gamma := \frac{1}{\sqrt{1-\frac{v^2}{c^2}}}$ is called the *Lorentz factor*. Using this transformation, we can derive a new rule for vector addition:

$$u' = \frac{dx'}{dt'} = \frac{dx'/dt}{dt'/dt} = \frac{\gamma(dx/dt - v)}{\gamma(1 - \frac{v}{c^2} dx/dt)} = \frac{u - v}{1 - u \frac{v}{c^2}}$$

If we use the inverse Lorentz transformation, we will obtain:

$$u = \frac{u' + v}{1 + u' \frac{v}{c^2}}$$

Now, if we put $u' = c$, then we have

$$u = \frac{c + v}{1 + \frac{cv}{c^2}} = c = u'$$

Hence the conflict is successfully resolved. However, in changing the rule of transformation, we had put a great risk to the already established laws, since their invariance is now not guaranteed.

3.1 What happens to physical laws?

- Since there is mixing between x and t in the transformation, it is clear that:

$$\frac{d^2 \mathbf{r}}{dt^2} \neq \frac{d^2 \mathbf{r}'}{dt'^2}$$

Thus, the second law is out of the picture! 🤖

- Imagine two balls falling, however the distance between them does not remain same. Then, at some instant, one of the ball conveys its position to the other ball and since the information takes a finite amount of time to reach the other balls, by that time, the first ball has fallen by by some distance and hence the forces $\mathbf{F}_{12}$ and $\mathbf{F}_{21}$ are not equal and opposite. Thus, the third law is out of the picture too! 😞
- Since in general, $r'^2 = (x'_2 - x'_1)^2 + (y'_2 - y'_1)^2 + (z'_2 - z'_1)^2 \neq r^2$, the law of gravitation also falls apart 🤖!

Hence, all the established laws were destroyed by the mere change of a transformation. We desperately needed another theory to reconcile with the physical laws. This led to the formulation of special and general relativity.

It is convenient to consider *infinitesimal transformation*, like:

$$\mathbf{r}_2 = \mathbf{r}_1 + d\mathbf{r} \implies (\mathbf{r}_2 - \mathbf{r}_1)^2 = d\mathbf{r} \cdot d\mathbf{r} = dr^2$$

Note that under Galilean transformation, dr^2 remain unchanged (invariant), however, under Lorentz transformation, it no longer remains constant. So, what is the invariant quantity under Lorentz transformation?

It turns out that if we define something like:

$$ds^2 = -c^2 dt^2 + dx^2 + dy^2 + dz^2 = -c^2 dt^2 + dr^2$$

then, ds^2 is invariant under Lorentz transformation.

Lemma 1 (Invariance of ds):

$$ds'^2 = ds^2$$

Proof. We use brute force to calculate the quantities:

$$\begin{aligned}
 ds'^2 &= -c^2 dt'^2 + dx'^2 + dy'^2 + dz'^2 \\
 &= -c^2 \gamma^2 \left(dt - \frac{v dx}{c^2} \right)^2 + \gamma^2 (dx - v dt)^2 + dy^2 + dz^2 \\
 &= -\gamma^2 c^2 \left(dt^2 + \frac{v^2 dx^2}{c^4} - 2 \frac{dx dt v}{c^2} \right) + \gamma^2 (dx^2 + v^2 dt^2 - 2v dx dt) + dy^2 + dz^2 \\
 &= -\gamma^2 c^2 dt^2 - \gamma^2 v^2 \frac{dx^2}{c^2} + \gamma^2 dx^2 + \gamma^2 v^2 dt^2 + dy^2 + dz^2 \\
 &= -c^2 \cancel{\gamma^2} dt^2 (1 - \cancel{v^2/c^2}) + \cancel{\gamma^2} dx^2 (1 - \cancel{v^2/c^2}) + dy^2 + dz^2 \\
 &= -c^2 dt^2 + dx^2 + dy^2 + dz^2 \\
 &= ds^2
 \end{aligned}$$

3.2 Classification of Intervals

Unlike Galilean transformation where $dr^2 > 0$ always, ds^2 need not be positive. In fact, it can take any value and accordingly we define the intervals. If the interval between two spacetime events is such that:

- $c^2 dt^2 > dr^2 \implies ds^2 < 0$: time-like interval
- $c^2 dt^2 < dr^2 \implies ds^2 > 0$: space-like interval
- $c^2 dt^2 = dr^2 \implies ds^2 = 0$: light-like/null interval

Note that for light, $\frac{dr}{dt} = \pm c \implies dr^2 = c^2 dt^2 \implies ds^2 = 0$. Hence, we call the interval light-like. Trajectory of light is a null trajectory. However, all massive objects move along a time-like trajectory.

Lecture 04: Vectors and Stuffs

Let us consider rotation about the z-axis by an angle θ , in which case, the x and y components transform as:

$$\begin{aligned}
 x' &= x \cos \theta + y \sin \theta \\
 y' &= -x \sin \theta + y \cos \theta
 \end{aligned}$$

Then, we can write the infinitesimal change in the components in the following form:

$$\begin{pmatrix} dx' \\ dy' \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Note that, this can also be written as:

$$\begin{pmatrix} dx' \\ dy' \end{pmatrix} = \begin{pmatrix} \frac{\partial x'}{\partial x} & \frac{\partial x'}{\partial y} \\ \frac{\partial y'}{\partial x} & \frac{\partial y'}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

The matrix in this case is the *Jacobian matrix* and this is a more general structure of coordinate transformations and hence any transformation can be written like this, even the Galilean or the Lorentz transformations.

This shows that transformations are just multiplication of a matrix with a vector. To this extent, we define a vector ¹ $\bar{x}$ with the following components:

$$\bar{x} = (x^0, x^1, x^2, x^3)^T$$

¹This is often called a *four vector*

Here $x^0 = ct$ and $x^1 = x, x^2 = y, x^3 = z$. Note that since t and x are not dimensionally same, we took $x^0 = ct$ to make it in terms of distance. We multiplied by c (and not any other velocity) because c is postulated to be a universal constant.

We can express any infinitesimal transformation as:

$$\begin{pmatrix} dx'^0 \\ dx'^1 \\ dx'^2 \\ dx'^3 \end{pmatrix} = \begin{pmatrix} \frac{\partial x'^0}{\partial x^0} & \cdots & \frac{\partial x'^0}{\partial x^3} \\ \vdots & \ddots & \vdots \\ \frac{\partial x'^3}{\partial x^0} & \cdots & \frac{\partial x'^3}{\partial x^3} \end{pmatrix} \begin{pmatrix} dx^0 \\ dx^1 \\ dx^2 \\ dx^3 \end{pmatrix}$$

In a more compact way, we can write in the vector-matrix form or in the component form using Einstein's summation convention:

$$d\bar{x}' = \Lambda d\bar{x} \iff dx'^\mu = \Lambda^\mu_\nu dx^\nu$$

where $\Lambda^\mu_\nu = \frac{\partial x'^\mu}{\partial x^\nu}$ is an element of the *matrix* Λ .

Summation convention: We drop the summation sign if twice repeated indices are there on the same side of the equation.

Now, let us define what a vector is:

Definition 1 (Vector):

If a mathematical object transforms under a coordinate transformation like a differential $d\bar{x}$, that is, $d\bar{x} \rightarrow d\bar{x}' = \Lambda d\bar{x}$, then it is called a *contravariant vector* or a *contravector* or a *tensor of rank (1, 0)*

Let us show that under the Galilean transformation, velocity, as defined by $\mathbf{u} = \frac{d\mathbf{r}}{dt}$ is a vector. We will define everything in contravariant and four-vector notation. For this, we consider the vector defined as before:

$$d\bar{x} = (ct, x, y, z)^T \equiv (x^0, x^1, x^2, x^3)^T$$

The Galilean transformation is defined by:

$$\lambda \equiv \begin{pmatrix} 1 & 0 & 0 & 0 \\ -v^1/c & 1 & 0 & 0 \\ -v^2/c & 0 & 1 & 0 \\ -v^3/c & 0 & 0 & 1 \end{pmatrix} \iff \begin{aligned} x'^0 &= x^0 \\ x'^1 &= x^1 - (v^1/c)x^0 \\ x'^2 &= x^2 - (v^2/c)x^0 \\ x'^3 &= x^3 - (v^3/c)x^0 \end{aligned}$$

We can then define the four velocity as¹:

$$u^\mu \equiv \frac{dx^\mu}{dt} = (c, u^1, u^2, u^3)^T$$

Now, we have:

$$\begin{aligned} dx'^\mu &= \frac{\partial x'^\mu}{\partial x^\nu} dx^\nu \\ \implies u'^\mu dt' &= \frac{\partial x'^\mu}{\partial x^\nu} u^\nu dt \quad (\text{as } dt = dt') \\ \implies u'^\mu &= \frac{\partial x'^\mu}{\partial x^\nu} u^\nu \end{aligned}$$

Thus, we see that the components of the velocity vector transform similar to the differential displacement and hence, velocity is indeed a vector under Galilean transformation. In a similar way, we can show that

¹Note that four velocity is defined as derivative with respect to something called as *proper time*, however, for Galilean transformations, proper time coincides with the coordinate time and hence this definition is okay.

acceleration $\mathbf{a} = \frac{d\mathbf{v}}{dt}$ is also a vector under Galilean transformation.

Now, let us turn to Lorentz transformation. Things become bad here! For example, consider the same definition of velocity $\mathbf{u} = \frac{d\mathbf{x}}{dt}$. However, here t does not remain invariant under coordinate transformation. Then we will have:

$$u'^{\mu} = \frac{dx'^{\mu}}{dt'} = \frac{\Lambda^{\mu}_{\nu} x^{\nu}}{\Lambda^0_{\alpha} dx^{\alpha}} \neq \frac{dx^{\mu}}{dt}$$

Thus, this definition fails to transform similar to a differential displacement and hence, the way it is defined presently, velocity is not a vector 😞.

Lecture 05: Some more Vectors and Stuffs

Previously, we saw how a vector was defined in terms of the transformation of the differential displacements. In here, we will consider another situation.

First, note that, using a given transformation,

$$dx'^{\mu} = \Lambda^{\mu}_{\nu} dx^{\nu}$$

we can define the inverse transformation,

$$dx^{\nu} = (\Lambda^{-1})^{\nu}_{\mu} dx'^{\mu}$$

The above statement is valid only if the transformation matrix is invertible (in most case, transformations like translations, rotations, etc. are invertible).

Let us now see what the inverse transformation matrix looks like. For that, consider:

$$\begin{aligned} dx'^{\mu} &= \frac{\partial x'^{\mu}}{\partial x^{\nu}} dx^{\nu} \\ \Rightarrow \frac{\partial x^{\beta}}{\partial x'^{\mu}} dx'^{\mu} &= \underbrace{\frac{\partial x^{\beta}}{\partial x'^{\mu}} \frac{\partial x'^{\mu}}{\partial x^{\nu}}}_{\delta^{\beta}_{\nu}} dx^{\nu} \\ \Rightarrow dx^{\beta} &= \frac{\partial x^{\beta}}{\partial x'^{\mu}} dx'^{\mu} \end{aligned}$$

Thus we see that:

$$(\Lambda^{-1})^{\nu}_{\mu} \equiv \frac{\partial x^{\nu}}{\partial x'^{\mu}}$$

5.1 Gradient Vector

In the Cartesian coordinates, we all know:

$$\nabla = \hat{\mathbf{i}} \frac{\partial}{\partial x} + \hat{\mathbf{j}} \frac{\partial}{\partial y} + \hat{\mathbf{k}} \frac{\partial}{\partial z}$$

In line with this, we will define another quantity (often called *four gradient*) such that:

$$\nabla_{\mu} := \frac{\partial}{\partial x^{\mu}}$$

A few points on this:

1. Note that we wrote the index in the down position for ∇_{μ} . This is done since we will see that the four gradient does not follow the transformation rule of a vector, hence it is a different entity. Since in vectors we put the indices upwards, to differentiate it from a vector, we put it down.

2. In spherical or cylindrical or any other coordinates, the *ordinary* gradient is not just the partial derivatives of the coordinates. There are some obnoxious prefactors involved also. Irrespective of that, we define the four gradient as just the derivative of the coordinates ('coz later we will see that those prefactors are just some jugglery with the *metric*).

The four gradient is better denoted by ∂_μ and thus, we have:

$$\bar{\partial} \equiv (\partial_0 \quad \partial_1 \quad \partial_2 \quad \partial_3) = \left(\frac{1}{c} \frac{\partial}{\partial t} \quad \frac{\partial}{\partial x} \quad \frac{\partial}{\partial y} \quad \frac{\partial}{\partial z} \right)$$

Under a coordinate transformation $x^\mu \rightarrow x'^\mu = \Lambda^\mu_\nu x^\nu$, we have using chain rule:

$$\partial'_\mu = \frac{\partial x^\nu}{\partial x'^\mu} \frac{\partial}{\partial x^\nu} \implies \partial'_\mu = (\Lambda^{-1})^\nu_\mu \frac{\partial}{\partial x^\nu}$$

The four gradient apparently transform accordingly as $\bar{\partial}' = \Lambda^{-1} \bar{\partial}$ which is contrary to that of the vector.

Definition 2:

Covector If a mathematical object transforms under a coordinate transformation like the derivative $\bar{\partial}$, then we call it a *covariant vector* or *covector* or a *tensor of rank (0,1)*

Actually vectors and covectors are defined based on how the basis vectors of the space transform. Vectors transform in a way *contrary* to the basis vectors and hence are called *contravariant* while covectors transform in a way similar to how the basis transforms and hence called *covariant*.¹

5.2 Tensors

Consider the product,

$$\omega = \bar{\mathbf{x}} \bar{\mathbf{x}}^T \equiv \begin{pmatrix} dx^0 \\ dx^1 \\ dx^2 \\ dx^3 \end{pmatrix} \begin{pmatrix} dx^0 & dx^1 & dx^2 & dx^3 \end{pmatrix} = \begin{pmatrix} dx^0 dx^0 & \dots & dx^0 dx^3 \\ \vdots & \ddots & \vdots \\ dx^3 dx^0 & \dots & dx^3 dx^3 \end{pmatrix}$$

Note that, defined like this, ω represents a square matrix and thus, has two indices. This operation is often called *outer product* and is denoted as

$$\omega = \bar{\mathbf{x}} \otimes \bar{\mathbf{x}} \quad \omega^{\mu\nu} = dx^\mu dx^\nu$$

Now, let us see how it transforms:

$$\begin{aligned} \omega^{\mu\nu} &= dx'^\mu dx'^\nu \\ &= \left(\frac{\partial x'^\mu}{\partial x^\alpha} dx^\alpha \right) \left(\frac{\partial x'^\nu}{\partial x^\beta} dx^\beta \right) \\ &= \left(\frac{\partial x'^\mu}{\partial x^\alpha} \right) \left(\frac{\partial x'^\nu}{\partial x^\beta} \right) dx^\alpha dx^\beta \end{aligned}$$

This is yet another kind of transformation, different from either vector and covector.

¹In the elegant language of differential geometry, vectors are actually the *tangent vectors* belonging to the *tangent space* of the space (*manifold*) while covectors belong to the *dual space* of the *tangent space*.

Definition 3 (Tensor):

If a mathematical object transforms, under a coordinate transformation, like $\underbrace{d\bar{x} \otimes \dots \otimes d\bar{x}}_{m \text{ times}}$ and like $\underbrace{\bar{d} \otimes \dots \otimes \bar{d}}_{n \text{ times}}$, then it is called a *tensor of rank (m,n)*

A prime example would be the *moment of inertia tensor* I^{ij} or the *electromagnetic tensor* $F_{\mu\nu}$.

We note that tensors have upstairs indices for each contravariant components and downstairs indices for each covariant component ¹.

Definition 4 (Scalar):

If a mathematical object does not transform under a coordinate transformation, then it is called a *scalar* or a *tensor of rank (0,0)*.

Few example of scalars would be the invariant displacement $dr^2 = dx^2 + dy^2 + dz^2$ under Galilean transformation and the invariant spacetime interval $ds^2 = -c^2 dt^2 + dr^2$ under Lorentz transformation.

5.3 Inner Product

Previously we saw the *outer product* and we expect an *inner product* too. Multiplying a vector with the transpose changed it to a square matrix. Here, we do the opposite. Consider the scalar dr^2 under Galilean transformation, which is written as:

$$dr^2 = \mathbf{dr} \cdot \mathbf{dr} = \mathbf{dr}^T \mathbf{dr} = \begin{pmatrix} dx^1 & dx^2 & dx^3 \end{pmatrix} \begin{pmatrix} dx^1 \\ dx^2 \\ dx^3 \end{pmatrix}$$

However, notice the apparent problem if we generalise this to Lorentz transformation too. We have:

$$ds^2 = -c^2 dt^2 + dr^2$$

This cannot be simply written as $d\bar{x}^T d\bar{x}$, all due to that *innocent* minus sign in front of $c^2 dt^2 \equiv (dx^0)^2$ which was necessary to distinguish spatial and temporal components.

How do we generalise this notion of scalar as some matrix product?

For that, let us define another matrix which will take care of the minus sign:

$$[g] \equiv \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}$$

Using this, we can check that:

$$ds^2 = \begin{pmatrix} dx^0 & dx^1 & dx^2 & dx^3 \end{pmatrix} \begin{pmatrix} -1 & & & \\ & 1 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \begin{pmatrix} dx^0 \\ dx^1 \\ dx^2 \\ dx^3 \end{pmatrix} = d\bar{x}^T g d\bar{x}$$

This g is called the *metric tensor* and in component form, the above can be written as:

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$$

¹We can easily change upstairs to downstairs indices by the *metric* and hence it is better to write tensors as T^μ_ν and not T_ν^μ so as to keep a vacant place for the indices to move up/down.

Note that, for Galilean transformation, the metric tensor is just the *identity* matrix,

$$[g]_{\text{GT}} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}$$

Lecture 06: Metric is also a Matrix

Previously, we saw that to get a scalar from a contra-vector, we had to use an additional matrix which led to the notion of the *metric tensor*. Let us analyse this object further...

6.1 Transformation of the metric

For this, we will consider the *sacred* rule that space-time interval should remain invariant under coordinate change. Thus, we have:

$$\begin{aligned} ds^2 &= ds'^2 \\ \implies g_{\mu\nu} dx^\mu dx^\nu &= g'_{\alpha\beta} dx'^\alpha dx'^\beta \\ \implies g_{\mu\nu} \left(\frac{\partial x^\mu}{\partial x'^\alpha} dx'^\alpha \right) \left(\frac{\partial x^\nu}{\partial x'^\beta} dx'^\beta \right) &= g'_{\alpha\beta} dx'^\alpha dx'^\beta \\ \implies g_{\mu\nu} \left(\frac{\partial x^\mu}{\partial x'^\alpha} \right) \left(\frac{\partial x^\nu}{\partial x'^\beta} \right) dx'^\alpha dx'^\beta &= g'_{\alpha\beta} dx'^\alpha dx'^\beta \end{aligned}$$

And now by the eternal rule for arbitrariness (of the different displacements), we conclude that:

$$g'_{\alpha\beta} = g_{\mu\nu} \left(\frac{\partial x^\mu}{\partial x'^\alpha} \right) \left(\frac{\partial x^\nu}{\partial x'^\beta} \right)$$

Thus, we see that the metric tensor transforms as a rank (0,2) tensor and hence, we were not wrong about calling this a *tensor*!

Few Properties of the metric:

- Note the following :

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = g_{\nu\mu} dx^\nu dx^\mu = g_{\nu\mu} dx^\mu dx^\nu$$

In the first step, we interchanged $\mu \leftrightarrow \nu$ since these are dummy indices and then we interchanged the placement of $dx^\mu \leftrightarrow dx^\nu$ since these are just components and hence commutes. Then, from this, we have $g_{\mu\nu} = g_{\nu\mu}$ which leads us to the fact that:

Metric tensor is symmetric

- If we have g to be non-singular, that is, $\det(g) \neq 0$, then we can define an inverse metric such that:

$$g g^{-1} = g^{-1} g = \mathbb{1}$$

In the component form, we can write

$$g_{\mu\nu} (g^{-1})^{\nu\alpha} = \delta^\alpha_\mu$$

We will henceforth use $g^{\nu\alpha}$ instead of $(g^{-1})^{\nu\alpha}$ for the inverse metric, since this is more sleek visually and also, it does not confuse us.

Note that, by the same logic as we showed that metric is a tensor, we can also show that inverse metric is a rank (2,0) tensor.

6.2 Metric is a map!

The metric (and its inverse too) is a map which takes a vector (covector) and changes it to a covector (vector) according as:

$$\begin{aligned} A_\mu &\xrightarrow{g} A^\nu & A_\mu &= g_{\mu\nu} A^\nu \\ A^\mu &\xrightarrow{g^{-1}} A_\nu & A^\mu &= g^{\mu\nu} A_\nu \end{aligned}$$

Using this, the norm of a vector can be written as:

$$\|\bar{\mathbf{A}}\| = \bar{\mathbf{A}} \cdot \bar{\mathbf{A}} = A_\mu A^\mu = A^\mu A_\mu$$

Lecture 07: Metric and STR

Consider the following transformation:

$$(r, \theta) \mapsto (x, y) \quad x = r \cos \theta \quad y = r \sin \theta$$

Let us calculate the transformation matrix for this transformation. Note that, here we have $\mathbf{x} \equiv (r, \theta)$ and $\mathbf{x}' \equiv (x, y)$. We calculate Λ from its definition as:

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \end{aligned} \quad \Lambda \equiv \begin{pmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{pmatrix} = \begin{pmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{pmatrix}$$

Once we have obtained the transformation matrix, we now move on to calculate the metric tensor in polar coordinates. For that, note, for Cartesian coordinate,

$$ds^2 = dx^2 + dy^2 \implies g \equiv \begin{pmatrix} 1 & \\ & 1 \end{pmatrix}$$

Then, according to this question, we already have the knowledge of $g'_{\mu\nu}$ and we have to transform it to $g_{\alpha\beta}$ and we already know the transformation rule. Also, since g' is diagonal, we need to only consider the diagonal components. Thus, all of our work is already done!

$$g_{\alpha\beta} = \Lambda^\mu_\alpha \Lambda^\nu_\beta g'_{\mu\nu}$$

- $g_{rr} = \Lambda^x_r \Lambda^x_r g'_{xx} + \Lambda^y_r \Lambda^y_r g'_{yy} = \cos^2 \theta + \sin^2 \theta = 1$
- $g_{\theta\theta} = \Lambda^x_\theta \Lambda^x_\theta g'_{xx} + \Lambda^y_\theta \Lambda^y_\theta g'_{yy} = r^2(\sin^2 \theta + \cos^2 \theta) = r^2$
- $g_{r\theta} = \Lambda^x_r \Lambda^x_\theta g'_{xx} + \Lambda^y_r \Lambda^y_\theta g'_{yy} = -r \cos \theta \sin \theta + r \sin \theta \cos \theta = 0$ (hence this metric is diagonal too!)

Thus, from this, we obtain

$$g \equiv \begin{pmatrix} 1 & \\ & r^2 \end{pmatrix}$$

From this, the invariant distance becomes our know result:

$$ds^2 = g_{ij} x^i x^j = dr^2 + r^2 (d\theta)^2$$

This is a weirdly powerful technique to find the metric in any coordinate system. And once we have found the metric, then calculations of stuff like gradient and Laplacians become very easy.

7.1 Proper Time

Time no longer remains a scalar in special theory of relativity. However, we do need some notion of invariant time which can be useful to define some quantities. At present, we only know two scalars in STR, that is, c and ds^2 . Note that, dimensionally, there is only one way to construct some 'time' using these two:

$$d\tau^2 := -\frac{ds^2}{c^2}$$

We define $d\tau$ to be the *proper time* which is a scalar, since it is constructed out of two scalars. The minus sign in the definition results from the convention of the $(-, +, +, +)$ metric that we had chosen.

Consider two events $\epsilon_1 = (ct_1, x, y, z)$ and $\epsilon_2 = (ct_2, x, y, z)$, which are happening at the same spatial location. Then, since $dr^2 = 0$, for these two events,

$$ds^2 = -c^2 dt^2 \implies dt = d\tau$$

Hence the time interval between two events happening at the same spatial location is the proper time. Note that, using this notion of time (which is a scalar) we can define velocities and momentum.

7.2 Looking back at STR

Let us now define some quantities:

- **four-position:** denoting the position of an object in space-time

$$\bar{\mathbf{x}} = (x^0, x^1, x^2, x^3)^T$$

- **four-velocity:** denoting a velocity of the object in space-time

$$\bar{\mathbf{u}} = \frac{d\bar{\mathbf{x}}}{d\tau} = \begin{pmatrix} \frac{d(ct)}{d\tau} \\ \frac{d\mathbf{x}}{d\tau} \end{pmatrix}$$

Note that, under this definition, $\bar{\mathbf{u}}$ becomes a tensor of rank $(1, 0)$.

Now, consider two reference frames; the moving frame moves with a velocity $\mathbf{v}$ with respect to the lab frame. Consider an observer at rest in the moving frame. Then, in the observer's frame, since spatial coordinates remain same, we have $ds'^2 = -c^2 d\tau^2$ while for the lab frame, we have $ds^2 = -cdt^2 + dx^2 + dy^2 + dz^2 = -c^2 dt^2 + dr^2$. Now, since the observer is at rest, they move with the same velocity $\mathbf{v}$ with respect to the lab frame and hence $\frac{d\mathbf{r}}{dt} = \mathbf{v}$. Hence, by the requirement that $ds'^2 = ds^2$ we have:

$$\begin{aligned} d\tau^2 &= -\frac{ds^2}{c^2} \\ &= -\frac{1}{c^2}(-c^2 dt^2 + dr^2) \\ &= dt^2 \left[1 - \frac{1}{c^2} \left(\frac{d\mathbf{r}}{dt} \right)^2 \right] \end{aligned} \tag{1}$$

From this we obtain that $d\tau = dt/\gamma$ where $\gamma := \frac{1}{\sqrt{1-(v^2/c^2)}}$. Substituting this in the above expression for four-velocity, we obtain:

$$u^\mu \equiv \gamma(c, d\mathbf{x}/dt)^T = \gamma(c, \mathbf{v})^T$$

- **four-momentum:** momentum of an object in space-time

$$\bar{\mathbf{p}} = m_o \bar{\mathbf{u}} = m_o \gamma(c, \mathbf{v})^T$$

Note that m_o as used here is simply the *mass*. The term rest-mass is often used, however, it often confuses us since this implies the existence of some *relativistic mass*.

Earlier, we used to define the relativistic mass as γm_o which kept the expression for spatial components of four-momentum and three momentum similar (that is, mass times velocity).

However, in recent times, the concept of relativistic mass is being done away with and we instead now change the expression of momentum when considering STR. Thus, we have:

$$\bar{\mathbf{p}} = (\gamma m_o c, \mathbf{p})$$

where $\mathbf{p} = \gamma m_o \mathbf{v}$ is in some way a new definition of 3-momentum.

7.2.1 Metric in STR

We have the invariant displacement as:

$$ds^2 = -c^2 dt^2 + dr^2$$

From this, we infer that the metric $g = \text{diag}(-1, 1, 1, 1)$. This metric is also termed as the *Minkowski metric* and we will henceforth denote this special metric by η . Using this metric, we have:

$$\|\bar{\mathbf{u}}\|^2 = \eta_{\mu\nu} u^\mu u^\nu = -\gamma^2 c^2 + \gamma^2 \|\mathbf{v}\|^2 = \gamma^2 c^2 \left(\frac{\|\mathbf{v}\|^2}{c^2} - 1 \right) = \gamma^2 c^2 \times \frac{-1}{\gamma^2} = -c^2$$

Thus, we obtain that the squared-norm of the four-velocity is always $-c^2$ 🤖, which is kinda cool since even without doing anything, we technically *move* at the speed of light!!

In a similar way, we obtain $\|\bar{\mathbf{p}}\|^2 = m_o^2 \|\bar{\mathbf{u}}\|^2 = -m_o^2 c^2$

Lecture 08: May the Force be with you!

From our previous definition of the 4-momentum, we can now define the 4-force as its derivative with respect to proper time. The 4-force is generally denoted by $\bar{\mathbf{K}}$. Thus,

$$\bar{\mathbf{K}} := \frac{d\bar{\mathbf{p}}}{d\tau}$$

Now, if we consider another frame, then we have:

$$\bar{\mathbf{K}}' = \frac{d\bar{\mathbf{p}}'}{d\tau'} = \frac{d\tau}{d\tau'} \frac{d(\Lambda \bar{\mathbf{p}})}{d\tau} = 1 \times \Lambda \frac{d\bar{\mathbf{p}}}{d\tau} = \Lambda \bar{\mathbf{K}}$$

This follows the rule of vector transformation and is thus a 4-vector. Moreover, note that, if $\bar{\mathbf{K}} = 0$ then $\bar{\mathbf{p}}$ is conserved. So, in the relativistic case too, *momentum is conserved in absence of force*.

8.1 Emergence of Energy

Note that when defining the 4-momentum, we obtained:

$$\bar{\mathbf{p}} = \gamma(m_o c, m_o \mathbf{v})$$

While the spatial components do have a clear Newtonian limit in case of $v \ll c$ ($\gamma \rightarrow 1$), it is not a priori clear what $p^0 = \gamma m_o c$ represent. Let us try to find out this quantity in the limit $v \ll c$!

$$\begin{aligned} p^0 &= \frac{m_o c}{\sqrt{1 - v^2/c^2}} \\ &= m_o c (1 - v^2/c^2)^{-1/2} \\ &= m_o c \left[1 + \frac{v^2}{2c^2} + \mathcal{O}(v^4/c^4) \right] \quad (\text{from Taylor expansion}) \end{aligned}$$

From the above thing, we obtain:

$$p^0 c = m_o c + \frac{1}{2} m_o v^2 + \mathcal{O}(v^4/c^4)$$

The second term clearly resembles our familiar kinetic energy while we do not have a clear idea about the first term and the higher order terms. However, this quantity $p^0 c$ seems to be a right fit to be called the *energy* (since it *does* contain one of the familiar energies). Let us then define the total energy of a relativistic particle as:

$$E := p^0 c$$

Earlier we had seen that the norm $\|\bar{\mathbf{p}}\| = -m_o^2 c^2$. Using this we have:

$$\begin{aligned} \bar{\mathbf{p}} \cdot \bar{\mathbf{p}} &= -m_o^2 c^2 \\ \Rightarrow g_{\mu\nu} p^\mu p^\nu &= -m_o^2 c^2 \\ \Rightarrow -(p^0)^2 + \|\mathbf{p}\|^2 &= -m_o^2 c^2 \\ \Rightarrow -E^2/c^2 + \|\mathbf{p}\|^2 &= -m_o^2 c^2 \\ \Rightarrow m_o^2 c^4 + \|\mathbf{p}\|^2 c^2 &= E^2 \\ \Rightarrow \sqrt{m_o^2 c^4 + \|\mathbf{p}\|^2 c^2} &= E \quad (\text{ignoring negative solution}) \end{aligned}$$


In the rest frame of the particle, the 3-momentum vanish, and hence we get the celebrated:

$$E = m_o c^2$$

Since the total energy of the particle $E = \sqrt{m_o^2 c^4 + \|\mathbf{p}\|^2 c^2}$, the kinetic energy can be obtained by subtracting the *rest energy term*, that is, $K.E = E - m_o c^2$

8.2 The many faces of Mass

Let us go back to the second law. Suppose there is a *heavy* ball and a *light* ball ¹. Then we subject them to the same force. Acceleration of both bodies happen, however, we of course expect the heavy ball to accelerate less than the light ball.



From this, we can say that there exists some innate property of an object which controls the amount of acceleration, even when same force is applied to different objects. This property is the proportionality constant between force and acceleration.

We can relate this to some kind of *inertia* (a fashionable way of saying lethargy). We define the *inertial mass* m^I as:

$$m^I := \frac{|\mathbf{F}|}{|\mathbf{a}|}$$

The inertial mass is a measure of 'heaviness to move'. Now, let us consider the Coulomb's law and Newton's Gravitational Law parallelly:

$$\mathbf{F}_e = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \hat{\mathbf{r}} \quad \mathbf{F}_g = G \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}$$

Similar to $\mathbf{F}_e$: electrical force, we have $\mathbf{F}_g$: gravitaional force while $\frac{1}{4\pi\epsilon_0}$: electrostatic coupling constant is analogous to G : gravitational coupling constant.

¹Consider that heaviness or lightness is being subjectively determined by the observer's opinion

Then indeed the electrical charge q_i must have the analogous gravitational charge m_i . Now, note that the gravitational charge, henceforth denoted by m^G has no apriori relation to the previous inertial mass m^I which came from the second law and was a property of the object itself.

We found an electrostatic analog of the gravitational law. It would have been nice to find some *magnetic analog* too, so that, we can also find a combined 'force' thing, similar to the Lorentz force $F = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$

Consider the electric due to a charge Q at a point. Then, the force on a charge q kept at the point is given by:

$$\mathbf{F} = q\mathbf{E} = m^I \mathbf{a} \implies \mathbf{a} = \left(\frac{q}{m^I}\right)\mathbf{E}$$

The acceleration depends on the charge-mass ratio. Similarly then in a gravitational field, we can write:

$$\mathbf{F} = m^G \mathbf{g} = m^I \mathbf{a} \implies \mathbf{a} = \left(\frac{m^G}{m^I}\right)\mathbf{g}$$

Till now we had assumed the equality of the two kind of 'masses' and then we found that the acceleration due to gravity was independent of masses, that is, every body has the same acceleration due to gravity. However, when we relax this assumption, we are at a critical stage.

8.2.1 Eötvös Experiment

So, this experiment was done to measure the different between the inertial and gravitational mass. In the experiment, two balls of different 'masses' were hanged from a rod and maintained in equilibrium.




Figure 5: Setup of the Eötvös Experiment


Then, considering the rotation of the Earth also, the torque was measured in the z-axis. From the measurement, we obtained experimentally:

$$\eta = \left| \frac{\frac{m_1^G}{m_1^I} - \frac{m_2^G}{m_2^I}}{\frac{1}{2} \left(\frac{m_1^G}{m_1^I} + \frac{m_2^G}{m_2^I} \right)} \right| \sim 10^{-9}$$

This proved that the ratio between the inertial and gravitational masses for different objects is nearly the same. However, experiments are limited by precision of the instruments. We can never definitely say whether $\frac{m_1^G}{m_1^I} = \frac{m_2^G}{m_2^I}$ by a mere experiment.

Lecture 09: Einstein's thoughts

Consider a planet with inertial mass m^I and gravitational mass m^G , rotating around a star with gravitational mass M^G .



Then, by the force balance condition and using $v = r\omega$, we can write the equation of motion as:

$$\frac{m^I v^2}{r} = G \frac{m^G M^G}{r^2} \implies \omega^2 = \left(\frac{m^G}{m^I} \right) \left(\frac{GM^G}{r^3} \right)$$

What this tells us is that, if indeed the ratio between inertial and gravitational mass is not constant, ω will be different for different objects and hence, two objects launched from the same point, will have their trajectories diverge at some point, which did not comply with the experimental fact.

Principle 1 (Weak Equivalence):

The ratio between gravitational and inertial mass is unity, that is,

$$\frac{m^I}{m^G} = 1$$


In other words, the principle states that the trajectory of a freely falling object is not affected by its mass. Let us see why!

$$m^I \mathbf{a} = -G \frac{m^G M^G}{r^2} \hat{\mathbf{r}} \implies \mathbf{a} = -\frac{GM^G}{r^2} \hat{\mathbf{r}}$$

Thus, we see that the acceleration of an object under the gravitational force is independent of its mass. In other words, the trajectory is independent on its mass.

9.1 How's gravity different?

Consider the following two cases:



In case (a), we see two charges q being thrown with same horizontal velocity v , from a building, under an electric field E . We previously saw that the acceleration depends on the charge-mass ratio and hence, depending on the mass of the charges, the trajectories will be different.


In case (b) however, if two masses m_1 and m_2 are thrown from building with same horizontal velocity v , both of them follow the same trajectory since acceleration under the force of gravity is independent of masses.

Thus we see an apparent disparity between gravity and other kind of forces (viz. *electromagnetic, strong nuclear, weak nuclear*).


9.2 Gedankenexperiment

We will now analyse a series of experiments in our minds and see what conclusions we can draw from each of them. All of them will consist of a blackened lift with an observer inside it and an 'all-knowing' outside observer who knows everything what is going on inside and outside the lift.

9.2.1 Lift in free space:



Case 1 : Free lift



Case 2 : Lift with rocket

Consider, there is a lift in free space (frame O'). The lift contains an observer with a ball and is shielded from outside, that is, that observer cannot see what is happening outside the lift. There is another observer outside on the ground (frame O), who has, by some divine grace, information both inside and outside the lift. Let the position of the ball from ground be y and the position of the lift from the ground be y_0 and the position of the ball from the lift be y' . Then, the constraint equation gives:

$$y = y' + y_0$$

Case 1: The lift is stationary and hence frame O' is inertial. As the lift is stationary, after the ball is released, no force acts on it and we have:

$$m\ddot{y}' = 0 \quad m\ddot{y} = 0$$

This is the trivial case as nothing as such occurs here. The observer in the lift, after releasing the ball, finds it at rest too and infers that the frame is inertial.

Case 2: Suppose now that the lift is attached to a rocket which is producing a constant acceleration a upwards. Then we have $\ddot{y}_0 = a$. Note that, O' is no longer an inertial frame. The observer inside the lift, after releasing the ball, will find it moving. Hence they will infer that their frame is non-inertial and cannot apply NLM 😞.

The outside observer can still apply NLM and since no force acts on the ball inside the lift (force acts only on the lift by the rocket), we have:

$$m\ddot{y} = 0 \implies \ddot{y}' = 0 - a = -a$$


Hence we see that, from the outside frame, the ball will seem to move downwards with an acceleration a . This is what we call the *pseudo-force*. Note that, no force actually acts on the ball.

9.3 Lift under Gravity

Case 1: Now, consider that we turn on a gravitational field $\mathbf{g} = -g\hat{\mathbf{y}}$ and let the lift be hanging from some support. O' is non-inertial (as inside observer, after dropping the ball, will see it move without any force applied) and hence only O frame observer can use NLM. Since from frame O , a force $-mg$ acts on the ball, we have:

$$m\ddot{y} = -mg \implies \ddot{y} = -g$$

The lift is not moving so $\ddot{y}_0 = 0$. Thus, from this, we get $\ddot{y}' = -g$



Case 2: Consider that the thread holding the lift is cut and the lift tragically falls. Then, we have from frame O :

$$m\ddot{y} = -mg$$


$$m_l\ddot{y}_0 = -m_l g$$

Then, from the above two and the constraint equation, we get $\ddot{y}' = -g - (-g) = 0$. The inside observer will then see that the acceleration of the ball with respect to them is zero. When released, the ball remains where it was. The observer then infers that they are in a 'free space'.

Conclusions: For an observer inside the lift,

1. A lift in *constant gravity* is same as an *accelerating lift*.
2. A *freely-falling* lift is the same as an *inertial lift*.

We see that there exist atleast one observer (the freely-falling one) with respect to whom, motion of an object under gravity appears as if it is moving under no force. We can then conclude that the force of gravity cannot be a *vector* as, if it was indeed a vector, it would transform like a vector which would mean that force in any other frame would have been zero¹.



¹A vector is zero in all frames iff it is zero in one frame (direct consequence of transformation)

Note that, if the force of gravity was *inhomogenous*, that is, varying with distance (which we know it is), then there would be pronounced effect inside the lift. Two balls thrown at a separation d inside the lift, will have their distance change if d is comparable to say the diameter of the earth. Then, the inside observer will no longer think that this is a free-space.

Thus, to eliminate the effect of gravity in a freely-falling frame, one must consider the frame to be *local*, that is, everything should be happening around a point and its neighbourhood only. Only then, the gravitational field can be treated as homogenous and the free-space assumption will be correct.

Lecture 10: Connections are important

In the previous section, we had seen that the concept of being *local* was introduced in order to approximate a frame where the effect of gravity is eliminated. Using such a motivation, we have:

Principle 2 (Einstein's Equivalence Principle):

All freely falling frames under the influence of *gravity* are *locally inertial*, that is, Lorentz frames where special relativity is valid and there is absence of gravity

Consider the case of the falling lift as before. Then, if we write the equation of the ball from the O frame, we have:

$$\frac{d\mathbf{r}}{dt} = -\mathbf{g} \implies \mathbf{r} = \mathbf{r}_i - \frac{1}{2}\mathbf{g}t^2$$

where $\mathbf{r}_i$ is the initial position of the ball. Now, from the O' frame, as no force acts on the ball (note that the observer sees the ball floating in front of them) we have:

$$\frac{d^2\mathbf{r}'}{dt'^2} = 0 \implies \mathbf{r}' = \mathbf{r}'_i$$

From the above equations we have:

$$\mathbf{r}' = \mathbf{r}'_i = (\mathbf{r}_i - \mathbf{r}_0) = \left(\mathbf{r} + \frac{1}{2}\mathbf{g}t^2\right) - \mathbf{r}_0$$

This essentially describes a *coordinate transformation* where $\mathbf{r} \rightarrow \mathbf{r}'$ and $t \rightarrow t'$. However, note that a term t^2 appears in the expression and hence, the coordinate transformations are *non-linear*. In general, we can have the coordinate transformation matrices themselves to be dependent on the coordinates, that is, $\Lambda^\mu{}_\nu \equiv \Lambda^\mu{}_\nu(x^\alpha)$

If we allow non-linear coordinate transformations, we can go to a frame where *gravity is absent!*

Thus, we essentially have a recipe to tackle gravity. We first go to a frame where gravity is absent (the equivalence principle guarantees the existence of such a frame), do our calculations in that frame, then transform back to our original coordinate system.

Consider a particle moving freely under a purely gravitational force. Now, suppose O' is such a frame where gravity is absent. Then we have:

$$\frac{d^2x'^\mu}{d\tau^2} = 0$$

How do we obtain the equation of motion in frame O ? Well, we will just transform to that frame. Since x'^μ is dependent on the coordinates of the other frame x^ν , we can write,

$$\begin{aligned}
 0 &= \frac{d}{d\tau} \left(\frac{dx'^\mu}{d\tau} \right) \\
 &= \frac{d}{d\tau} \left(\frac{\partial x'^\mu}{\partial x^\nu} \frac{dx^\nu}{d\tau} \right) \\
 &= \frac{\partial x'^\mu}{\partial x^\nu} \frac{d^2 x^\nu}{d\tau^2} + \frac{dx^\nu}{d\tau} \frac{d}{d\tau} \left(\frac{\partial x'^\mu}{\partial x^\nu} \right) \\
 &= \frac{\partial x'^\mu}{\partial x^\nu} \frac{d^2 x^\nu}{d\tau^2} + \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} \frac{dx^\alpha}{d\tau} \frac{dx^\nu}{d\tau}
 \end{aligned}$$

The first term seems to contain the acceleration in the O frame and it would be nice to isolate this term. For that, let us multiply both sides by $\frac{\partial x^\beta}{\partial x'^\mu}$ and contract using the fact that $\frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial x'^\mu}{\partial x^\nu} = \delta^\beta_\nu$

$$\begin{aligned}
 0 &= \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial x'^\mu}{\partial x^\nu} \frac{d^2 x^\nu}{d\tau^2} + \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} \frac{dx^\alpha}{d\tau} \frac{dx^\nu}{d\tau} \\
 &= \delta^\beta_\nu \frac{d^2 x^\nu}{d\tau^2} + \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} \frac{dx^\alpha}{d\tau} \frac{dx^\nu}{d\tau} \\
 &= \frac{d^2 x^\beta}{d\tau^2} + \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} \frac{dx^\alpha}{d\tau} \frac{dx^\nu}{d\tau} \\
 &= \frac{d^2 x^\beta}{d\tau^2} + \Gamma^\beta_{\nu\alpha} u^\alpha u^\nu
 \end{aligned}$$

Here we have defined the following:

$$\Gamma^\beta_{\nu\alpha} := \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} = (\Lambda^{-1})^\beta_\mu \frac{\partial}{\partial x^\nu} (\Lambda^\mu_\alpha)$$

The above entities (for all indices) are termed as *connection coefficients*. The name connection coefficients comes from the fact that they connect neighboring points and tell us how to calculate the rate of change of a vector field from one point to another nearby, even though the coordinate system may itself be changing. The equation of motion thus obtained is:

$$\frac{d^2 x^\beta}{d\tau^2} = -\Gamma^\beta_{\nu\alpha} u^\alpha u^\nu \quad (2)$$

The above equation is called the *geodesic equation* and resembles some kind of *pseudo-force* or *non-inertial force*. Note that the connection coefficients in its present form, depends on both $\bar{x}$ and $\bar{x}'$ which is kind of problematic since we might not always have knowledge of both the frames.

Since the primed frame is locally inertial, the metric is *Minkowski*, that is,

$$\eta_{\mu\nu} = g'_{\mu\nu} = \frac{\partial x^\alpha}{\partial x'^\mu} \frac{\partial x^\beta}{\partial x'^\nu} g_{\alpha\beta}$$

Inverting the relation, we have:

$$g_{\mu\nu} = \frac{\partial x'^\alpha}{\partial x^\mu} \frac{\partial x'^\beta}{\partial x^\nu} \eta_{\alpha\beta}$$

From the definition of the affine connection, we can see that, since partial derivatives commute, the affine connections are *symmetric* in the lower indices. Thus,

$$\Gamma^\alpha_{\mu\nu} = \Gamma^\alpha_{\nu\mu}$$

Now, let us contract the upper index of the affine connection definition which will be used later:

$$\frac{\partial x'^\lambda}{\partial x^\beta} \Gamma^\beta_{\nu\alpha} = \frac{\partial x'^\lambda}{\partial x^\beta} \frac{\partial x^\beta}{\partial x'^\mu} \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} = \delta^\lambda_\mu \frac{\partial^2 x'^\mu}{\partial x^\nu \partial x^\alpha} = \frac{\partial^2 x'^\lambda}{\partial x^\nu \partial x^\alpha} = \partial_\nu \left(\frac{\partial x'^\lambda}{\partial x^\alpha} \right)$$

Then, let us differentiate both sides with x^ρ and using the above found relation, we obtain:

$$\begin{aligned} \partial_\rho g_{\mu\nu} &= \partial_\rho \left(\frac{\partial x'^\alpha}{\partial x^\mu} \right) \times \frac{\partial x'^\beta}{\partial x^\nu} \eta_{\alpha\beta} + \partial_\rho \left(\frac{\partial x'^\beta}{\partial x^\nu} \right) \times \frac{\partial x'^\alpha}{\partial x^\mu} \eta_{\alpha\beta} \\ &= \Gamma^\sigma_{\rho\mu} \frac{\partial x'^\alpha}{\partial x^\sigma} \frac{\partial x'^\beta}{\partial x^\nu} \eta_{\alpha\beta} + \Gamma^\sigma_{\rho\nu} \frac{\partial x'^\beta}{\partial x^\sigma} \frac{\partial x'^\alpha}{\partial x^\mu} \eta_{\alpha\beta} \\ &= \Gamma^\sigma_{\rho\mu} g_{\sigma\nu} + \Gamma^\sigma_{\rho\nu} g_{\mu\sigma} \end{aligned}$$

We thus obtain:

$$\partial_\rho g_{\mu\nu} = \Gamma^\sigma_{\rho\mu} g_{\sigma\nu} + \Gamma^\sigma_{\rho\nu} g_{\mu\sigma} \quad (3)$$

Since these are true for any indices, we obtain two more equations by the exchange $\rho \leftrightarrow \mu$ and $\rho \leftrightarrow \nu$:

$$\partial_\mu g_{\rho\nu} = \Gamma^\sigma_{\mu\rho} g_{\sigma\nu} + \Gamma^\sigma_{\mu\nu} g_{\rho\sigma} \quad (4)$$

$$\partial_\nu g_{\mu\rho} = \Gamma^\sigma_{\nu\mu} g_{\sigma\rho} + \Gamma^\sigma_{\nu\rho} g_{\mu\sigma} \quad (5)$$

Adding Eq. 4 and Eq. 5 and then subtracting Eq. 3, we obtain:

$$\begin{aligned} \partial_\mu g_{\rho\nu} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu} &= \Gamma^\sigma_{\mu\rho} g_{\sigma\nu} + \Gamma^\sigma_{\mu\nu} g_{\rho\sigma} + \Gamma^\sigma_{\nu\mu} g_{\sigma\rho} + \Gamma^\sigma_{\nu\rho} g_{\mu\sigma} - \Gamma^\sigma_{\rho\mu} g_{\sigma\nu} - \Gamma^\sigma_{\rho\nu} g_{\mu\sigma} \\ \Rightarrow \Gamma^\sigma_{\mu\nu} g_{\rho\sigma} &= \frac{1}{2} (\partial_\mu g_{\rho\nu} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}) \\ \Rightarrow g^{\rho\lambda} \Gamma^\sigma_{\mu\nu} g_{\rho\sigma} &= \frac{1}{2} g^{\rho\lambda} (\partial_\mu g_{\rho\nu} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}) \\ \Rightarrow \delta^\lambda_\sigma \Gamma^\sigma_{\mu\nu} &= \frac{1}{2} g^{\rho\lambda} (\partial_\mu g_{\rho\nu} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}) \end{aligned}$$

The red and the green terms cancel first and then we contract the metric on the left to obtain an extremely important identity for the affine connection:

$$\Gamma^\lambda_{\mu\nu} = \frac{1}{2} g^{\rho\lambda} (\partial_\mu g_{\rho\nu} + \partial_\nu g_{\mu\rho} - \partial_\rho g_{\mu\nu}) \quad (6)$$

We obtained an expression for the affine connection in terms of the metric for the space. In this form, these are called the *Christoffel symbols* and denoted by:

$$\left\{ \begin{matrix} \lambda \\ \mu\nu \end{matrix} \right\}$$

NOTE: We can remember the equation like this: In the left hand side, the upper index is λ which comes only in the metric in the right side. The derivatives with $+$ sign are with respect to the lower indices μ and ν and the index of the metric in the derivative is a dummy index ρ and the other lower index not appearing in the derivative. Finally, the derivative with $-$ sign is with respect to the dummy index and the metric contains both the lower indices of the left.

Hence, we see that the equation of motions are known provided the Christoffel symbols are known, which are known if metric is known!

10.1 Action tells a lot!

We will now consider one of the most fundamental quantities in physics, the *action*, albeit in the *relativistic* sense.

$$\mathcal{S} = \int dt L = \int dt (T - V)$$

Note that, action must be invariant under 3-rotation in Newtonian mechanics. Since relativistic case should also converge to the classical action in the Newtonian limit, we want the action to be also invariant.

Till now, we know three invariant (scalar) quantities: $m, c, d\tau$ (or ds). Let us construct a quantity with same dimension as of Lagrangian. We then define:

$$\mathcal{S}_{\text{rel}} = \int d\tau L := \mathcal{A} \int d\tau mc^2$$

where we replaced the Lagrangian by $\mathcal{A}mc^2$, $\mathcal{A}$ is just some constant. This is just our guess but let us see what happens in the Newtonian limit! Note that, using Eq. 1, we can write the action as:

$$\begin{aligned} \mathcal{S}_{\text{rel}} &= \mathcal{A}mc^2 \int dt \sqrt{1 - v^2/c^2} \\ &= \mathcal{A}mc^2 \int dt (1 - v^2/2c^2 + \mathcal{O}(v^4/c^4)) \\ &= -\mathcal{A} \int dt \left(\frac{1}{2}mv^2 - mc^2 \right) \end{aligned}$$

Well, this looks like kind of $T - V$ if we identify the rest energy as some 'potential' energy, provided $\mathcal{A} = -1$. Thus, in the Newtonian limit atleast, this definition of *relativistic action* makes sense kinda. Hence we have:

$$\mathcal{S}_{\text{rel}} := \int d\tau (-mc^2) \quad (7)$$

Note that, we had previously seen that, $\|\bar{\mathbf{u}}\|^2 = -c^2$ and hence we can replace $-c$ with this quantity in the action expression. Using this we obtain:

$$\mathcal{S}_{\text{rel}} = \int d\tau m \|\bar{\mathbf{u}}\|^2 = \int d\tau m g_{\mu\nu} u^\mu u^\nu$$

Now, we know that we can derive the ELEOM from the least action principle ($\delta\mathcal{S} = 0$). We have:

$$\frac{\partial L}{\partial x^\lambda} = m \partial_\lambda g_{\mu\nu} u^\mu u^\nu \quad \frac{\partial L}{\partial u^\lambda} = 2m g_{\lambda\nu} u^\nu$$

Now, the metric could be a function of the coordinates and hence we have:

$$\frac{d}{d\tau} \left(\frac{\partial L}{\partial u^\lambda} \right) = 2m \times \left(g_{\lambda\nu} \frac{du^\nu}{d\tau} + \partial_\sigma g_{\lambda\nu} u^\sigma u^\nu \right)$$

Plugging this into the ELEOM, we obtain:

$$2m \times \left(g_{\lambda\nu} \frac{du^\nu}{d\tau} + \partial_\sigma g_{\lambda\nu} u^\sigma u^\nu \right) = m \partial_\lambda g_{\mu\nu} u^\mu u^\nu \implies g_{\lambda\nu} \frac{du^\nu}{d\tau} = \frac{1}{2} (\partial_\lambda g_{\mu\nu} u^\mu u^\nu) - \partial_\sigma g_{\lambda\nu} u^\sigma u^\nu$$

Then, contracting the metric on the left side with $g^{\rho\lambda}$, we get:

$$\frac{du^\rho}{d\tau} = g^{\lambda\rho} \times \left(\frac{1}{2} \partial_\lambda g_{\sigma\nu} - \partial_\sigma g_{\lambda\nu} \right) u^\sigma u^\nu$$

Let us manipulate the term in the bracket:

$$\begin{aligned}
 & \left(\frac{1}{2} \partial_\lambda g_{\sigma\nu} - \partial_\sigma g_{\lambda\nu} \right) u^\sigma u^\nu \\
 &= \frac{1}{2} \partial_\lambda g_{\sigma\nu} u^\sigma u^\nu - \frac{1}{2} (\partial_\sigma g_{\lambda\nu} + \partial_\sigma g_{\nu\lambda}) u^\sigma u^\nu \\
 &= \frac{1}{2} \partial_\lambda g_{\sigma\nu} u^\sigma u^\nu - \frac{1}{2} \partial_\sigma g_{\lambda\nu} u^\sigma u^\nu - \frac{1}{2} \partial_\sigma g_{\nu\lambda} u^\sigma u^\nu \\
 &= \frac{1}{2} \partial_\lambda g_{\sigma\nu} u^\sigma u^\nu - \frac{1}{2} \partial_\sigma g_{\lambda\nu} u^\sigma u^\nu - \frac{1}{2} \partial_\nu g_{\sigma\lambda} u^\nu u^\sigma \\
 &= \frac{1}{2} (\partial_\lambda g_{\sigma\nu} - \partial_\sigma g_{\lambda\nu} - \partial_\nu g_{\sigma\lambda}) u^\sigma u^\nu
 \end{aligned}$$


Above, we had used the symmetry of the metric tensor and then we renamed the dummy indices in the sum. We substitute this in the previous equation and obtain:

$$\frac{du^\rho}{d\tau} + \frac{1}{2} g^{\lambda\rho} (\partial_\sigma g_{\lambda\nu} + \partial_\nu g_{\sigma\lambda} - \partial_\lambda g_{\sigma\nu}) u^\sigma u^\nu = 0 \implies \frac{du^\rho}{d\tau} + \Gamma^\rho_{\sigma\nu} u^\sigma u^\nu = 0$$

This is exactly the *geodesic* equation that we had obtained earlier. Hence, using this definition of the action, we can recover the geodesic equation. Thus, **geodesics are paths of extreme action!**

Lecture 11: Finding Geodesics

Since a particle following the geodesics have no *proper acceleration* in a local inertial frame, the geodesics generally represent the 'straightest' path between two points in a space. Let us take the example for the shortest distance between two cities *A* and *B* on the Earth. Since the Earth can be assumed to be a sphere, we just need to find the geodesic equation on $\mathbb{S}^2$ and then solve it.



Intuitively, out of the infinite ways of going between those cities, we think that moving along a line of latitude would be more economical. However, let us check if that is indeed so!

For that, note that the invariant distance on the sphere $\mathbb{S}^2$ of radius r_0 is:

$$ds^2 = r_0^2 d\theta^2 + r_0^2 \sin^2 \theta d\phi^2$$

From this, we can easily identify the metric and inverse metric for the space as:

$$[g] \equiv \begin{pmatrix} r_0^2 & 0 \\ 0 & r_0^2 \sin^2 \theta \end{pmatrix} \quad [g^{-1}] \equiv \begin{pmatrix} 1/r_0^2 & 0 \\ 0 & 1/r_0^2 \sin^2 \theta \end{pmatrix}$$

Thus we have,

$$g^{\theta\theta} = \frac{1}{r_0^2} \quad g_{\theta\theta} = r_0^2 \quad g^{\phi\phi} = \frac{1}{r_0^2 \sin^2 \theta} \quad g_{\phi\phi} = r_0^2 \sin^2 \theta$$

Now, the geodesic equation contains the Christoffel symbols and let us calculate them one by one. Note that the metric being diagonal is a blessing since then, the apparent summation in the Christoffel symbol expression is not needed.

$$\Gamma_{\theta\theta}^{\theta} = \frac{1}{2}g^{\theta\theta}(\partial_{\theta}g_{\theta\theta} + \partial_{\theta}g_{\theta\theta} - \partial_{\theta}g_{\theta\theta}) = 0$$

$$\Gamma_{\theta\phi}^{\theta} = \frac{1}{2}g^{\theta\theta}(\partial_{\theta}g_{\theta\phi} + \partial_{\phi}g_{\theta\theta} - \partial_{\theta}g_{\theta\phi}) = 0$$

$$\begin{aligned}\Gamma_{\phi\phi}^{\theta} &= \frac{1}{2}g^{\theta\theta}(\partial_{\phi}g_{\theta\phi} + \partial_{\phi}g_{\theta\phi} - \partial_{\theta}g_{\phi\phi}) \\ &= -\frac{1}{2}r_0^2 \sin \theta \cos \theta \cdot \frac{1}{r_0^2} \\ &= -\sin \theta \cos \theta\end{aligned}$$

$$\Gamma_{\theta\theta}^{\phi} = \frac{1}{2}g^{\phi\phi}(\partial_{\theta}g_{\theta\phi} + \partial_{\theta}g_{\theta\phi} - \partial_{\phi}g_{\theta\theta}) = 0$$

$$\begin{aligned}\Gamma_{\theta\phi}^{\phi} &= \frac{1}{2}g^{\phi\phi}(\partial_{\theta}g_{\phi\phi} + \partial_{\phi}g_{\theta\phi} - \partial_{\phi}g_{\theta\phi}) \\ &= \frac{1}{2} \cdot \frac{1}{r_0^2 \sin^2 \theta} \cdot (2r_0^2 \sin \theta \cos \theta) \\ &= \cot \theta\end{aligned}$$

$$\Gamma_{\phi\phi}^{\phi} = \frac{1}{2}g^{\phi\phi}(\partial_{\phi}g_{\phi\phi} + \partial_{\phi}g_{\phi\phi} - \partial_{\phi}g_{\phi\phi}) = 0$$

Thus, we get only two non-zero Christoffel symbols. Since there are two parameters θ and ϕ for the problem, we would obtain two geodesic equations which are:

$$\frac{d^2x^{\theta}}{d\tau^2} + \Gamma_{\phi\phi}^{\theta} \frac{dx^{\phi}}{d\tau} \frac{dx^{\phi}}{d\tau} = 0 \quad \implies \quad \ddot{\theta} - \sin \theta \cos \theta \dot{\phi}^2 = 0 \quad (8)$$

$$\frac{d^2x^{\phi}}{d\tau^2} + \Gamma_{\theta\phi}^{\phi} \frac{dx^{\phi}}{d\tau} \frac{dx^{\theta}}{d\tau} + \Gamma_{\phi\theta}^{\phi} \frac{dx^{\theta}}{d\tau} \frac{dx^{\phi}}{d\tau} = 0 \quad \implies \quad \ddot{\phi} + 2 \cot \theta \dot{\theta} \dot{\phi} = 0 \quad (9)$$

Multiplying Eq. 9 with $\sin^2 \theta$, we obtain:

$$\sin^2 \theta \ddot{\phi} + 2 \sin \theta \cos \theta \dot{\phi} \dot{\theta} \implies \frac{d}{d\tau}(\dot{\phi} \sin^2 \theta) = 0$$

The above equation implies that $\dot{\phi} \sin^2 \theta$ is a constant of motion and let us denote it by $\mathcal{A}$. Thus, we obtain,

$$\dot{\phi} = \frac{\mathcal{A}}{\sin^2 \theta}$$

Using the above in Eq. 8, we will get:

$$\ddot{\theta} = \frac{\cos \theta}{\sin^3 \theta} \mathcal{A}^2$$

Note that we had obtained the equations for θ and ϕ in terms of τ , however, we want to understand θ in terms of ϕ or vice-versa, which would give us some equation of the trajectory. For this, let us use our beloved chain rule!

$$\begin{aligned}\dot{\theta} &= \frac{d\theta}{d\phi} \frac{d\phi}{d\tau} = \frac{d\theta}{d\phi} \times \frac{\mathcal{A}}{\sin^2 \theta} \\ \ddot{\theta} &= \frac{d\phi}{d\tau} \frac{d}{d\phi} \left(\frac{d\theta}{d\phi} \frac{\mathcal{A}}{\sin^2 \theta} \right) = \frac{\mathcal{A}}{\sin^2 \theta} \frac{d}{d\phi} \left(\frac{d\theta}{d\phi} \frac{\mathcal{A}}{\sin^2 \theta} \right)\end{aligned}$$

Using this in Eq. 8 we have:

$$\begin{aligned} \frac{\mathcal{A}}{\sin^2 \theta} \frac{d}{d\phi} \left(\frac{\mathcal{A}}{\sin^2 \theta} \frac{d\theta}{d\phi} \right) - \frac{\mathcal{A}^2}{\sin^2 \theta} \frac{\sin \theta \cos \theta}{\sin^2 \theta} &= 0 \\ \Rightarrow \frac{d}{d\phi} \left(\frac{\mathcal{A}}{\sin^2 \theta} \frac{d\theta}{d\phi} \right) - \mathcal{A} \cot \theta &= 0 \\ \Rightarrow \frac{d}{d\phi} \left(\frac{1}{\sin^2 \theta} \frac{d\theta}{d\phi} \right) - \cot \theta &= 0 \end{aligned}$$

Now note that $\frac{d}{d\phi}(\cot \theta) = -\frac{1}{\sin^2 \theta} \frac{d\theta}{d\phi}$ and using this, our equation becomes:

$$-\frac{d}{d\phi} \left(\frac{d}{d\phi}(\cot \theta) \right) - \cot \theta = 0 \Rightarrow \frac{d^2 \cot \theta}{d\phi^2} = -\cot \theta$$

The solution of the above can be written as:

$$\cot \theta = c_1 \cos \phi + c_2 \sin \phi \quad \text{for some } c_1, c_2 \quad (10)$$

So, what does this equation represent? For that, let us consider a plane passing through the origin whose equation is given by:

$$ax + by + cz = 0$$

Now consider any sphere of radius r_0 and centred at origin. Any point on the sphere is parameterised by:

$$x = r_0 \sin \theta \cos \phi \quad y = r_0 \sin \theta \sin \phi \quad z = r_0 \cos \theta$$

The intersection of the plane passing through origin (centre of sphere) and the sphere essentially gives what is known as *great circles* whose equation can be given by simultaneously solving the two equations above. We have,

$$a(r_0 \sin \theta \cos \phi) + b(r_0 \sin \theta \sin \phi) + cr_0 \cos \theta = 0 \Rightarrow (a \cos \phi + b \sin \phi) + c \cot \phi = 0$$

This gives us the equation of the great circle as,

$$\cot \theta = d_1 \cos \phi + d_2 \sin \phi$$

which is the same form as Eq. 10. Thus, we conclude that the geodesics on the sphere $\mathbb{S}^2$ are the *great circles* and hence it is actually shorter to move along the great circle to reach from city A to city B, rather than moving along the latitudes.





Figure 6: Moving along the great circle is shorter than moving along the latitude.

Note however that, if we travel on the opposite arc of the great circle, it will be the longest route. This is because, geodesics are paths of *extreme* action, not only minimum. In general, we should thus never fly along the latitude (except if it is the equator, since equator is itself a great circle). Now, let us consider a cylinder.



The invariant displacement for the cylinder can be written as

$$ds^2 = r_0^2 d\theta^2 + dz^2$$

Note that the metric for this space is,

$$[g] \equiv \begin{pmatrix} r_0^2 & 0 \\ 0 & 1 \end{pmatrix}$$

This is a *constant metric* which does not depend on the parameters θ or z . Then, all the Christoffel symbols will be zero since Christoffel symbols are essentially written in terms of the derivatives of the metric. The geodesic equations thus simplify to:

$$\frac{d^2\theta}{d\tau^2} = 0 \quad \frac{d^2z}{d\tau^2} = 0$$

which have the solutions:

$$\theta = c_1\tau + c_2 \quad z = d_1\tau + d_2 \implies z = d_1\left(\frac{\theta - c_2}{c_1}\right) + d_2 \implies z = m\theta + k$$

Suitably taking $k = 0$, we see that the equation represents a helix with pitch $2\pi m$. Thus, the geodesics on the cylinder are helices and the shortest distance to move from one point to another on the cylinder is along a helix.

NOTE: Similar to the case of the cylinder, for the flat spacetime, the metric is $\eta_{\mu\nu} = \text{diag}(-1, +, +, +)$ which is constant too and hence, the geodesics for the Minkowski space will also satisfy the same equation as of the cylinder:

$$x^\mu = a^\mu + u^\mu\tau$$

where u^μ is the constant four-velocity and a^μ is the initial position (at $\tau = 0$). This shows that geodesics in flat spacetime are straight lines.

Lecture 12: Towards the action for Gravity: Invariant Volumes

Motion under gravity can be determined once we know the metric $g_{\mu\nu}$, however, we want to see how to obtain the equation for the metric. In other words, since equations of motion can be derived from *action*, we need to find an action for gravity!

12.1 Maxwell's action in spacetime

In free space, the *electromagnetic* action can be written as:

$$\mathcal{S}_{\text{EM}} = \int d^4\bar{\mathbf{x}} \left[-\frac{1}{4\mu_0} F_{\mu\nu} F^{\mu\nu} \right]$$

Here $F_{\mu\nu}$ is the anti-symmetric electromagnetic tensor and is defined by:

$$F_{\mu\nu} := \partial_\mu A_\nu - \partial_\nu A_\mu \quad F_{\mu\nu} = -F_{\nu\mu}$$

$A^\mu = (\phi, \mathbf{A})$ is the four-potential, which we will simply call the *electromagnetic field*. From the above defined action, we can obtain all of Maxwell's equations. We can manipulate the action as follows,

$$\begin{aligned} \mathcal{S}_{\text{EM}} &= \frac{1}{4\mu_0} \int d^4\bar{\mathbf{x}} - [\partial_\mu A_\nu - \partial_\nu A_\mu] F^{\mu\nu} \\ &= 2 \times \frac{1}{4\mu_0} \int d^4\bar{\mathbf{x}} [-\partial_\mu A_\nu F^{\mu\nu}] \quad (\text{using anti-symmetric property}) \\ &\stackrel{\text{IBP}}{=} \frac{1}{2\mu_0} \int d^4\bar{\mathbf{x}} [A_\nu \partial_\mu F^{\mu\nu} - \partial_\mu (A_\nu F^{\mu\nu})] \end{aligned}$$

The last integral is a total-derivative which can be converted to the surface integral using Stokes' theorem. We assume that at spatial and temporal infinities, the fields vanish and thus, the integral is zero. Then we obtain,

$$\mathcal{S}_{\text{EM}} = \frac{1}{2\mu_0} \int d^4\bar{\mathbf{x}} [A_\nu \partial_\mu F^{\mu\nu}] = \frac{1}{2\mu_0} \int d^4\bar{\mathbf{x}} [A_\nu \partial_\mu (\partial_\mu A_\nu - \partial_\nu A_\mu)]$$


Properties of the Maxwell action:

- $\mathcal{S}_{\text{EM}}$ is a functional of A_μ , that is, it takes a value of A_μ and returns a number.
- $\mathcal{S}_{\text{EM}}$ is a scalar under Lorentz transformation. We can see this since the indices in F are contracted and $d^4\bar{\mathbf{x}}$ can be shown to be Lorentz-invariant.
- $\mathcal{S}_{\text{EM}}$ has at most two derivatives of the field A_μ

Einstein took inspiration from this action, however, he tried to include the invariance of this action for general transformations too! He proposed that *action* for the metric should be of the following form:

$$\mathcal{S}_G = \int d^4\bar{\mathbf{x}} f(g_{\mu\nu}, \partial_\alpha \partial_\beta g_{\mu\nu})$$

The above mentioned form has many subtleties, let us look at them one by one. First note that if the function f has to be a scalar under general coordinate transformations, then the integral measure $d^4\bar{\mathbf{x}}$ must also be a scalar. However, this is not apriori obvious. As an example, note how the 'volume' element changes when we move from Cartesian to polar coordinates.



In Cartesian coordinates, the volume element (actually the area element since it is 2D) is $dV = dx dy$ while in the polar coordinates, the volume element is $dV' = dr (r d\theta) = r d\theta dr$. There is an extra factor r apart from the coordinate differential. We denote it by $J(r, \theta)$ which is the *Jacobian* of the transformation. Thus, to keep the volume element invariant, we need to add the effect of the Jacobian too!

12.2 Metric, Transformations and Jacobian

Consider the Leibniz formula for the determinant of a square $n \times n$ matrix. The formula says that for any matrix A , we have ¹:

$$\det(A) = \sum_{\tau \in S_n} \text{sgn}(\tau) \prod_{i=1}^n a_{\tau(i)}^i = \sum_{\sigma \in S_n} \text{sgn}(\sigma) \prod_{i=1}^n a_{\sigma(i)}^i$$

From this, we can derive another formula for the determinant using the Levi-Civita symbol:

Identity 1:

$$\det(M) = \tilde{\epsilon}^{\mu'_1 \dots \mu'_n} \tilde{\epsilon}_{\mu_1 \dots \mu_n} A^{\mu_1}_{\mu'_1} A^{\mu_2}_{\mu'_2} \dots A^{\mu_n}_{\mu'_n}$$

Let us keep this aside and calculate the volume element for a two-dimensional coordinate system, in primed and unprimed frames. Note that the volume element can be written in the following way:

$$\begin{aligned} dV &= J(\bar{x}) \frac{1}{2!} \epsilon_{\alpha\beta} (-1)^p dx^\alpha dx^\beta \\ dV &= J(\bar{x}') \frac{1}{2!} \epsilon_{\mu\nu} (-1)^p dx^\mu dx^\nu \end{aligned}$$

What just happened here! 😊. First note that p here is the sign of the permutation, that is, if the permutation α, β is odd then $p = -1$, else $p = 0$. This is done to ensure that the quantity is positive. Also note that, J as defined here should always be taken to be positive, to make dV non-negative. Subtleties aside, if we expand the first expression, we would get:

$$dV = J(\bar{x}) \frac{1}{2} (\epsilon_{12} (-1)^0 dx^1 dx^2 + \epsilon_{21} (-1)^1 dx^2 dx^1) = J(\bar{x}) dx^1 dx^2$$

This is indeed the expression for the volume element in the unprimed frame. The above formula is related to differential forms and wedge products which are defined in terms of *alternating-operator* and gives a generalisation of the volume element $dV = dx^1 \wedge dx^2 \wedge \dots$ which we will not explore.

Anyways, let us now calculate something. Since we want $dV = dV'$, then we demand that:

$$J(\bar{x}) \frac{1}{2!} \epsilon_{\alpha\beta} (-1)^p dx^\alpha dx^\beta = J(\bar{x}') \frac{1}{2!} \epsilon_{\mu\nu} (-1)^p dx'^\mu dx'^\nu$$

Let us take the primed coordinate to be Cartesian and hence, $J(\bar{x}') = 1$. Then, transforming the coordinates we obtain:

$$J(\bar{x}) \frac{1}{2!} \epsilon_{\alpha\beta} (-1)^p dx^\alpha dx^\beta = \frac{1}{2!} \epsilon_{\mu\nu} (-1)^p \Lambda^\mu_\alpha \Lambda^\nu_\beta dx^\alpha dx^\beta$$

Now, since the differentials are arbitrary, both quantities are equal. Let us contract both sides with $\epsilon^{\alpha\beta}$ which gives us:

$$J(\bar{x}) \frac{1}{2!} \epsilon_{\alpha\beta} \epsilon^{\alpha\beta} = \frac{1}{2!} \epsilon_{\mu\nu} \epsilon^{\alpha\beta} \Lambda^\mu_\alpha \Lambda^\nu_\beta = \det(\Lambda^\mu_\alpha)$$

In the RHS, we have used the formula that we had written for the determinant previously. In the LHS, note that $\epsilon_{\alpha\beta} \epsilon^{\alpha\beta}$ will be 2 and hence, the final result that we obtained is:

$$J(\bar{x}) = \det(\Lambda^\mu_\alpha)$$

Thus, the Jacobian of the transformation is basically the determinant of the transformation matrix. Using the above found expression, we can write the volume element (now, in arbitrary dimensions) as:

$$dV = \det(\Lambda^\mu_\nu) \prod_\alpha dx^\alpha$$

¹You may also look into Sec 6.5.3 [here](#)!

Now, let us take the metric. Since the primed frame is *flat*, the metric there is Minkowski, $\eta_{\mu\nu}$. Then, from the transformation rule, we will have:

$$g = \Lambda^T \eta \Lambda$$

Since the determinant of Λ and Λ^T are same and $\det(\eta) = -1$, we obtain:

$$\det(g) = -\det(\Lambda)^2 = -J^2 \implies J = \sqrt{|\mathbf{g}|}$$

where we have defined $\mathbf{g} := \det(g)$, the determinant of the metric tensor for the space. Generalising to n dimensions, we have the volume element as:

$$dV = d^n \vec{x} \sqrt{|\mathbf{g}|}$$

As an example, for 2D polar coordinates, we have:

$$g \equiv \text{diag}(1, r^2) \implies \mathbf{g} = r^2 \implies dV = dr d\theta \sqrt{r^2} = r dr d\theta$$

And for 3D spherical coordinates we have,

$$g \equiv \text{diag}(1, r^2, r^2 \sin^2 \theta) \implies \mathbf{g} = r^4 \sin^2 \theta \implies dV = dr d\theta d\phi \sqrt{r^4 \sin^2 \theta} = r^2 \sin \theta dr d\theta d\phi$$

which are indeed the case as we know from our experience! Thus, this gives a nice expression to deal with the volume elements in some arbitrary space with some arbitrary metric.

Lecture 13: Towards the action for Gravity: Covariant Derivatives

Previously, we focussed on making the integral measure invariant in the action expression. Now, we focus on the function $f(g_{\mu\nu}, \partial_\alpha \partial_\beta g_{\mu\nu})$. Since this contains derivatives, let us first see what happens to derivatives of tensors!

Consider a scalar field ϕ^1 . Then, under a coordinate transformation $\bar{x} \rightarrow \bar{x}'$, we have $\phi(\bar{x}) \rightarrow \phi'(\bar{x}')$. Since this is a scalar field, we expect $\phi'(\bar{x}') = \phi(\bar{x})$. An example of such a field can be the temperature at each point in space and at each time.

Let us see how the derivative of this scalar field changes under a general coordinate transformation.

$$\partial'_\mu \phi'(\bar{x}') = (\Lambda^{-1})^\alpha_\mu \partial_\alpha (\phi(\bar{x}))$$

This transforms as a tensor of rank $(0, 1)$ and is thus a valid tensor. Now, if we take a vector instead, let us see what changes.

$$\begin{aligned} \partial'_\mu V'^\rho &= (\Lambda^{-1})^\alpha_\mu \partial_\alpha (\Lambda^\rho_\beta V^\beta) \\ &= (\Lambda^{-1})^\alpha_\mu \Lambda^\rho_\beta (\partial_\alpha V^\beta) + (\Lambda^{-1})^\alpha_\mu V^\beta \partial_\alpha (\Lambda^\rho_\beta) \end{aligned}$$

If the second term had not been there, the quantity would have transformed as a rank $(1, 1)$ tensor but since under a general coordinate transformation, Λ is itself a function of the coordinates, the second term is non-zero in general, which spoils the transformation property.

Since this causes a problem, a potential solution could be to define a new kind of derivative operator, say ∇_μ , such that $\nabla_\mu V^\rho$ becomes a tensor of rank $(1, 1)$. Thus, we demand that,

$$\nabla'_\mu V'^\nu = (\Lambda^{-1})^\alpha_\mu \Lambda^\nu_\rho (\nabla_\alpha V^\rho)$$

¹It gives us a single value of some variable for every point in spacetime

Inverting this expression, we obtain:

$$\nabla_\alpha V^\rho = (\Lambda^{-1})^\rho{}_\nu \Lambda^\mu{}_\alpha (\nabla'_\mu V'^\nu) \quad (11)$$

Moving forward with this would be difficult and hence we make a comment on physical grounds, that, in a *local inertial frame*, where spacetime is flat, the new derivative is equivalent to the normal derivative ∂_μ . Hence, if the coordinate system $\{\bar{x}'\}$ is locally inertial, then

$$\nabla'_\mu = \partial'_\mu$$

Using this in Eq. 11, we have:

$$\begin{aligned} \nabla_\alpha V^\rho &= (\Lambda^{-1})^\rho{}_\nu \Lambda^\mu{}_\alpha [\partial'_\mu (\Lambda^\nu{}_\beta V^\beta)] \\ &= (\Lambda^{-1})^\rho{}_\nu \partial_\alpha [(\Lambda^\nu{}_\beta V^\beta)] \\ &= (\Lambda^{-1})^\rho{}_\nu \Lambda^\nu{}_\beta \partial_\alpha V^\beta + (\Lambda^{-1})^\rho{}_\nu V^\beta \partial_\alpha (\Lambda^\nu{}_\beta) \\ &= \delta^\rho{}_\beta \partial_\alpha V^\beta + \Gamma^\rho{}_{\alpha\beta} V^\beta \\ &= \partial_\alpha V^\rho + \Gamma^\rho{}_{\alpha\beta} V^\beta \end{aligned}$$

Thus, on physical grounds we obtained:

$$\nabla_\alpha V^\rho = \partial_\alpha V^\rho + \Gamma^\rho{}_{\alpha\beta} V^\beta \quad (12)$$

The above defines a new derivative, called the *covariant derivative*. If we use this definition, then we can obtain the correct transformation. Note that the extra term that spoiled the transformation, is already incorporated into the covariant derivative definition through the affine connection. Thus, *covariant derivatives behave nicely!*

NOTE: An easy way to remember this is that two of the indices of the affine connection are placed exactly similar as in the LHS while the other index is summed over. And we have a normal derivative on the RHS too.

If we instead start with a covariant vector, then:

$$\begin{aligned} \nabla_\alpha V_\beta &= \Lambda^\nu{}_\beta \Lambda^\mu{}_\alpha \nabla'_\mu V'_\nu \\ &= \Lambda^\nu{}_\beta \Lambda^\mu{}_\alpha \partial'_\mu V'_\nu \\ &= \Lambda^\nu{}_\beta \Lambda^\mu{}_\alpha [\partial'_\mu ((\Lambda^{-1})^\rho{}_\nu V_\rho)] \\ &= \Lambda^\nu{}_\beta \partial_\alpha [(\Lambda^{-1})^\rho{}_\nu V_\rho] \\ &= \underbrace{\Lambda^\nu{}_\beta (\Lambda^{-1})^\rho{}_\nu}_{\delta^\rho{}_\beta} \partial_\alpha V_\rho + \Lambda^\nu{}_\beta V_\rho \partial_\alpha (\Lambda^{-1})^\rho{}_\nu \\ &= \partial_\alpha V_\beta + \Lambda^\nu{}_\beta V_\rho \partial_\alpha (\Lambda^{-1})^\rho{}_\nu \end{aligned}$$

Now note the following:

$$\begin{aligned} (\Lambda^{-1})^\rho{}_\nu \Lambda^\nu{}_\theta &= \delta^\rho{}_\theta \\ \Rightarrow (\partial_\alpha (\Lambda^{-1})^\rho{}_\nu) \Lambda^\nu{}_\theta &= -(\partial_\alpha \Lambda^\nu{}_\theta) (\Lambda^{-1})^\rho{}_\nu \quad (\text{Liebniz rule}) \\ \Rightarrow (\partial_\alpha (\Lambda^{-1})^\rho{}_\nu) \Lambda^\nu{}_\theta (\Lambda^{-1})^\theta{}_\gamma &= -(\partial_\alpha \Lambda^\nu{}_\theta) (\Lambda^{-1})^\rho{}_\nu (\Lambda^{-1})^\theta{}_\gamma \\ \Rightarrow \partial_\alpha (\Lambda^{-1})^\rho{}_\gamma &= -(\partial_\alpha \Lambda^\nu{}_\theta) (\Lambda^{-1})^\rho{}_\nu (\Lambda^{-1})^\theta{}_\gamma \end{aligned}$$

Substituting this in the above equation, we obtain:

$$\begin{aligned}
 \nabla_\alpha V_\beta &= \partial_\alpha V_\beta - \Lambda^\nu{}_\beta \partial_\alpha (\Lambda^{-1})^\rho{}_\nu V_\rho \\
 &= \partial_\alpha V_\beta - \Lambda^\nu{}_\beta V_\rho (\partial_\alpha \Lambda^\gamma{}_\theta) (\Lambda^{-1})^\rho{}_\gamma (\Lambda^{-1})^\theta{}_\nu \\
 &= \partial_\alpha V_\beta - V_\rho (\partial_\alpha \Lambda^\gamma{}_\theta) (\Lambda^{-1})^\rho{}_\gamma \delta^\theta{}_\beta \\
 &= \partial_\alpha V_\beta - V_\rho (\partial_\alpha \Lambda^\gamma{}_\beta) (\Lambda^{-1})^\rho{}_\gamma \\
 &= \partial_\alpha V_\beta - \Gamma^\rho{}_{\alpha\beta} V_\rho
 \end{aligned}$$

Thus, we obtain:

$$\nabla_\alpha V_\beta = \partial_\alpha V_\beta - \Gamma^\rho{}_{\alpha\beta} V_\rho$$

For any general mixed tensor, for each contravariant index, we have ‘+’ and for every covariant index, we have ‘−’ in the expression for the covariant derivative, for example,

$$\nabla_\mu T^\alpha{}_\beta = \partial_\mu T^\alpha{}_\beta + \Gamma^\alpha{}_{\mu\rho} T^\rho{}_\beta - \Gamma^\rho{}_{\mu\beta} T^\alpha{}_\rho$$

This new definition of derivative gave us a way to compute quantities nicely, so that the transformation rule remains valid. Later, we would also see physical interpretation this quantity.

Lecture 14: Towards the action for Gravity: Riemann Tensor

Covariant derivatives transform a tensor of rank (p, q) to a tensor of rank $(p, q + 1)$ as seen before. Since in our proposed form of the action for gravity, there exists two derivatives of the metric, let us calculate and see what the covariant derivative of the metric is. We have,

$$\nabla_\mu (g_{\alpha\beta}) = \partial_\mu g_{\alpha\beta} - \Gamma^\rho{}_{\mu\alpha} g_{\rho\beta} - \Gamma^\sigma{}_{\mu\beta} g_{\alpha\sigma}$$

Now using Eqn. 6 and contracting metric on the right hand side, we get:

$$g_{\rho\beta} \Gamma^\rho{}_{\mu\alpha} = \frac{1}{2} (\partial_\mu g_{\alpha\beta} + \partial_\alpha g_{\mu\beta} - \partial_\beta g_{\alpha\mu})$$

Using this in the equation above, we get:

$$\nabla_\mu (g_{\alpha\beta}) = \partial_\mu g_{\alpha\beta} - \frac{1}{2} (\partial_\mu g_{\alpha\beta} + \partial_\alpha g_{\mu\beta} - \partial_\beta g_{\alpha\mu}) - \frac{1}{2} (\partial_\mu g_{\beta\alpha} + \partial_\beta g_{\mu\alpha} - \partial_\alpha g_{\beta\mu}) = 0$$

We see that the covariant derivative of the metric is zero, hence the metric tensor is said to be *covariantly conserved*.

Show the following things:

- $\Gamma^\alpha{}_{\mu\alpha} = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g})$
- $\nabla_\mu V^\mu = \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} V^\mu)$
- Leibniz rule is satisfied by covariant derivatives, that is,

$$\nabla_\nu (\phi V^\mu) = (\nabla_\nu \phi) V^\mu + \phi (\nabla_\nu V^\mu) \quad \nabla_\nu (W_\lambda V^\mu) = (\nabla_\nu W_\lambda) V^\mu + W_\lambda (\nabla_\nu V^\mu)$$

▷ For this we will prove the following identity first:

Identity 2:

For any diagonalizable matrix M ,

$$\text{Tr}(M^{-1}dM) = d(\ln \det M),$$

where dM is the differential of the matrix.

We first calculate the variation of the RHS:

$$\begin{aligned} d \ln(\det M) &= \ln \det(M + dM) - \ln \det(M) \\ &= \ln \left(\frac{\det(M + dM)}{\det M} \right) \\ &= \ln (\det(M^{-1}) \det(M + dM)) \\ &= \ln \det(\mathbb{1} + M^{-1}dM). \end{aligned}$$

Now suppose that M is diagonalizable, hence, M can be written as, $M = PDP^{-1}$, with D being the diagonal matrix of eigenvalues $\{\lambda_i\}$. Then we have:

$$\begin{aligned} \det M &= \prod_{i=1}^N \lambda_i \implies \ln \det M = \sum_i \ln \lambda_i = \text{Tr}[\mathbb{1} \times \text{diag}(\ln \lambda_1, \ln \lambda_2, \dots)] \\ &= \text{Tr}[P^{-1}P \text{diag}(\ln \lambda_1, \ln \lambda_2, \dots)] \\ &= \text{Tr}[P \text{diag}(\ln \lambda_1, \ln \lambda_2, \dots)P^{-1}] \\ &= \text{Tr} \ln M. \end{aligned}$$

Using the above two facts, we obtain:

$$\begin{aligned} d \ln(\det M) &= \ln \det(\mathbb{1} + M^{-1}dM) = \text{Tr} \ln(\mathbb{1} + M^{-1}dM) \\ &= \text{Tr}[M^{-1}dM + \mathcal{O}(M^{-1}dM M^{-1}dM)] \end{aligned}$$

Now, take the case when $M = g^{\mu\nu} \equiv g$ and $\det(g) = \mathfrak{g}$. Applying the above formula, we obtain:

$$\partial_\mu \ln(-\mathfrak{g}) = \text{Tr}(g \partial_\mu g) = (g \partial_\mu g)^{\rho\rho} = g^{\rho\sigma} \partial_\mu g_{\rho\sigma} \quad (13)$$

Now, from the definition of Christoffel symbol:

$$\Gamma^\alpha_{\mu\alpha} = \frac{1}{2} g^{\alpha\sigma} (\partial_\mu g_{\alpha\sigma} + \partial_\alpha g_{\mu\sigma} - \partial_\sigma g_{\mu\alpha}) = \frac{1}{2} g^{\alpha\sigma} \partial_\mu g_{\alpha\sigma}$$

Note that, if we rename $\sigma \rightarrow \alpha$, the last term cancels with the first term above and hence we get the simplified result. Substituting this expression in Eqn. 13, we get:

$$\begin{aligned} \Gamma^\alpha_{\mu\alpha} &= \partial_\mu \ln(-\mathfrak{g}) \\ &= \frac{1}{2} \frac{1}{-\mathfrak{g}} \partial_\mu (-\mathfrak{g}) \\ &= \frac{1}{2\sqrt{-\mathfrak{g}}\sqrt{-\mathfrak{g}}} \partial_\mu (-\mathfrak{g}) \\ &= \frac{1}{\sqrt{-\mathfrak{g}}} \partial_\mu (\sqrt{-\mathfrak{g}}) \end{aligned}$$

Therefore we finally proved the identity,

$$\Gamma^\alpha_{\mu\alpha} = \frac{1}{\sqrt{-\mathfrak{g}}} \partial_\mu \sqrt{-\mathfrak{g}}.$$

▷ For the second part, we use the first identity.

$$\begin{aligned}
 \nabla_\alpha V^\alpha &= \partial_\alpha V^\alpha + \Gamma^\alpha_{\mu\alpha} V^\mu \\
 &= \partial_\alpha V^\alpha + \frac{1}{\sqrt{-g}} \partial_\mu \sqrt{-g} V^\mu \\
 &= \partial_\alpha V^\alpha + \frac{1}{\sqrt{-g}} (\partial_\mu (\sqrt{-g} V^\mu) - \sqrt{-g} \partial_\mu V^\mu) \\
 &= \partial_\alpha V^\alpha - \cancel{\partial_\mu V^\mu} + \frac{1}{\sqrt{-g}} \partial_\mu (\sqrt{-g} V^\mu)
 \end{aligned}$$

Changing $\mu \rightarrow \alpha$, we have proved the identity.

▷ For the third part, we just try to match the LHS and RHS. First we do it for the scalar and vector.

$$\begin{aligned}
 \nabla_\nu (\phi V^\mu) &= \partial_\nu (\phi V^\mu) + \Gamma^\mu_{\nu\sigma} (\phi V^\sigma) \\
 &= (\partial_\nu \phi) V^\mu + \phi \partial_\nu V^\mu + \phi \Gamma^\mu_{\nu\sigma} V^\sigma \\
 &= (\nabla_\nu \phi) V^\mu + \phi \nabla_\nu V^\mu
 \end{aligned}$$

For scalars, partial and covariant derivatives are the same and hence we obtained the first term like that. Now, let us show for the two contravectors and covectors.

$$\begin{aligned}
 \nabla_\mu (V^\alpha W_\beta) &= \partial_\mu (V^\alpha W_\beta) + (\Gamma^\alpha_{\mu\sigma} V^\sigma) W_\beta - (\Gamma^\rho_{\beta\mu} V^\alpha) W_\rho \\
 &= \partial_\mu (V^\alpha) W_\beta + V^\alpha \partial_\mu (W_\beta) + (\Gamma^\alpha_{\mu\sigma} V^\sigma) W_\beta - (\Gamma^\rho_{\beta\mu} V^\alpha) W_\rho \\
 &= W_\beta \{ \partial_\mu V^\alpha + \Gamma^\alpha_{\mu\sigma} V^\sigma \} + V^\alpha \{ \partial_\mu W_\beta - (\Gamma^\rho_{\beta\mu} W_\rho) \} \\
 &= W_\beta (\nabla_\mu V^\alpha) + V^\alpha (\nabla_\mu W_\beta)
 \end{aligned}$$

Now, let us delve into an involved calculation. The covariant derivative of the metric is already zero, so in the expression for the action, we cannot put any covariant derivatives of the metric. However, we need to somehow obtain an expression with two normal derivatives of the metric. For that, let us just see what happens when we take two covariant derivatives of a contravector. ¹

$$\begin{aligned}
 \nabla_\mu (\nabla_\nu V^\rho) &= \partial_\mu (\nabla_\nu V^\rho) + \Gamma^\rho_{\mu\sigma} (\nabla_\nu V^\sigma) - \Gamma^\sigma_{\mu\nu} (\nabla_\sigma V^\rho) \\
 &= \partial_\mu (\partial_\nu V^\rho + \Gamma^\rho_{\nu\lambda} V^\lambda) + \Gamma^\rho_{\mu\sigma} (\partial_\nu V^\sigma + \Gamma^\sigma_{\nu\lambda} V^\lambda) - \Gamma^\sigma_{\mu\nu} (\nabla_\sigma V^\rho)
 \end{aligned}$$

Let us now calculate the commutator of this, that is, $[\nabla_\mu, \nabla_\nu] V^\rho$. To write in short, we will denote the other symmetric term with $\mu \leftrightarrow \nu$.

$$[\nabla_\mu, \nabla_\nu] V^\rho = \partial_\mu (\partial_\nu V^\rho + \Gamma^\rho_{\nu\lambda} V^\lambda) + \Gamma^\rho_{\mu\sigma} (\partial_\nu V^\sigma + \Gamma^\sigma_{\nu\lambda} V^\lambda) - \Gamma^\sigma_{\mu\nu} (\nabla_\sigma V^\rho) - (\mu \leftrightarrow \nu)$$

Note that the red term is itself symmetric in μ and ν and hence will be cancelled at once from the above expression. Same goes with the blue term. Thus, we are left with:

$$\begin{aligned}
 [\nabla_\mu, \nabla_\nu] V^\rho &= \partial_\mu (\Gamma^\rho_{\nu\lambda} V^\lambda) + \Gamma^\rho_{\mu\sigma} \partial_\nu V^\sigma + \Gamma^\rho_{\mu\sigma} \Gamma^\sigma_{\nu\lambda} V^\lambda - (\mu \leftrightarrow \nu) \\
 &= \partial_\mu (\Gamma^\rho_{\nu\lambda}) V^\lambda + \{ \Gamma^\rho_{\nu\lambda} \partial_\mu (V^\lambda) + \Gamma^\rho_{\mu\sigma} \partial_\nu V^\sigma \} + \Gamma^\rho_{\mu\sigma} \Gamma^\sigma_{\nu\lambda} V^\lambda - (\mu \leftrightarrow \nu)
 \end{aligned}$$

Note the term in the bracket. This term, as a whole, is symmetric under the exchange of μ and ν and hence will be cancelled from the commutator. This significantly reduces the number of terms and we obtain,

$$\begin{aligned}
 [\nabla_\mu, \nabla_\nu] V^\rho &= \partial_\mu (\Gamma^\rho_{\nu\lambda}) V^\lambda + \Gamma^\rho_{\mu\sigma} \Gamma^\sigma_{\nu\lambda} V^\lambda - \partial_\nu (\Gamma^\rho_{\mu\lambda}) V^\lambda - \Gamma^\rho_{\nu\sigma} \Gamma^\sigma_{\mu\lambda} V^\lambda \\
 &= \{ \partial_\mu (\Gamma^\rho_{\nu\lambda}) + \Gamma^\rho_{\mu\sigma} \Gamma^\sigma_{\nu\lambda} - \partial_\nu (\Gamma^\rho_{\mu\lambda}) - \Gamma^\rho_{\nu\sigma} \Gamma^\sigma_{\mu\lambda} \} V^\lambda
 \end{aligned}$$

¹Note that the quantity in the bracket below, on the LHS, is a $(1, 1)$ tensor.

The commutator is a proper tensor since this is difference of two tensors and thus, the right hand side must also be a tensor which we define as:

$$R^\rho{}_{\lambda\mu\nu} := \partial_\mu \Gamma^\rho{}_{\nu\lambda} + \Gamma^\rho{}_{\mu\sigma} \Gamma^\sigma{}_{\nu\lambda} - \partial_\nu \Gamma^\rho{}_{\mu\lambda} - \Gamma^\rho{}_{\nu\sigma} \Gamma^\sigma{}_{\mu\lambda}$$

The above object is a tensor of rank $(1, 3)$ and is called the *Riemann tensor*. The tensor represents the curvature of the space, which we will discuss later. We can lower all the indices to obtain a $(0, 4)$ tensor as,

$$R_{\rho\lambda\mu\nu} = g_{\rho\sigma} R^\sigma{}_{\lambda\mu\nu}$$

NOTE: How to remember this horrendous term? Well, what works for me is, note that the first lower index always goes as the last lower index in the RHS. There are two terms, one containing a derivative and another one containing the product of the Christoffel symbols.

Start putting all the indices on the RHS in the same order as in LHS (and in the same way, upwards or downwards). Then, ∂ will get second lower index, μ Christoffel gets the other two indices. In product of Christoffels, the first Christoffel gets ρ and μ and then all the up indices gets exhausted, so put the other index down in the next Christoffel. Fill the remaining terms with some dummy index (indicating summing over). And in the next other terms, interchange $\mu \leftrightarrow \nu$. As simple as that...

Some fun facts...

- The Riemann tensor has two normal derivatives of the metric (as the Christoffel symbol contains one derivative and we are taking derivative of the symbols).
- For Minkowski space, all Christoffel symbols vanish and thus $R^\rho{}_{\lambda\mu\nu}$ is identically zero. This gives a hint of the tensor's connection with curvatures, since Minkowski space is 'flat' with no curvature.
- If we start with a covariant vector, we have:

$$\begin{aligned} [\nabla_\mu, \nabla_\nu] V_\rho &= [\nabla_\mu, \nabla_\nu] (g_{\rho\lambda} V^\lambda) = g_{\rho\lambda} [\nabla_\mu, \nabla_\nu] V^\lambda = g_{\rho\lambda} R^\lambda{}_{\sigma\mu\nu} V^\sigma \\ &= R_{\rho\sigma\mu\nu} V^\sigma \\ &= R_{\rho\sigma\mu\nu} g^{\sigma\theta} V_\theta \\ &= R_{\rho}{}^{\theta}{}_{\mu\nu} V_\theta \end{aligned}$$

- Since the commutator is anti-symmetric with respect to μ and ν , the Riemann tensor is also anti-symmetric with respect to the last two indices by construction, that is,

$$R_{\rho\sigma\mu\nu} = -R_{\rho\sigma\nu\mu}$$

- R is also anti-symmetric with respect to the first two indices (which is not too obvious). For that, we use the commutator on the metric and use the property of commutator for covariant vectors:

$$0 = [\nabla_\mu, \nabla_\nu] g_{\alpha\beta} = R_{\alpha}{}^{\sigma}{}_{\mu\nu} g_{\sigma\beta} + R_{\beta}{}^{\sigma}{}_{\mu\nu} g_{\alpha\sigma} = R_{\alpha\beta\mu\nu} + R_{\beta\alpha\mu\nu} \implies R_{\alpha\beta\mu\nu} = -R_{\beta\alpha\mu\nu}$$

- $R_{\rho\sigma\mu\nu}$ is perhaps a very big ranked tensor with four indices. We can try to contract it perhaps. Note that, if we contract with the first two or last two, it will give zero, since metric is *symmetric* and Riemann tensor is *anti-symmetric* with respect to the first or last two indices. Thus, let us contract with the first and third index. We obtain a tensor called the *Ricci tensor*, that is,

$$R_{\mu\nu} = g^{\alpha\beta} R_{\alpha\mu\beta\nu}$$

- We can contract the Ricci tensor further to get a *Ricci scalar*:

$$R = g^{\mu\nu} R_{\mu\nu}$$

Note that the Ricci scalar has two derivatives of the metric and is also a scalar, thus satisfying the condition to be a part of the action.

Thus, we are finally in a position to sew together all the components of the action. The function f can be a function of the Ricci scalar, that is, $f \equiv f(R)$. We take the simplest possible case where f is linear in R , that is,

$$f \equiv \alpha(R - 2\Lambda)$$

Here α and Λ ¹ are some constants. With this, the action becomes:

$$S_G = \alpha \int d^4x \sqrt{-g} [R - 2\Lambda]$$

This is almost fine, we just need to find something more about α . Note that action has dimensions of the angular momentum and the metric is dimensionless. Thus, R will have dimensions L^{-2} since it has two derivatives of a dimensionless quantity (metric). Thus,


$$\dim \alpha = \frac{\dim S_G}{[L^4] \dim R} = \frac{ML^2T^{-1}}{L^4L^{-2}} = \frac{(LT^{-1})^3}{\dim G} = \frac{c^3}{G}$$

So, upto some numerical factor, $\alpha \sim \frac{c^3}{G}$. The final action, which we refer to as the *Einstein-Hilbert action*, is given by:

$$S_{EH} = \frac{c^3}{16\pi G} \int d^4x \sqrt{-g} [R - 2\Lambda] = \frac{c^4}{16\pi G} \int dt d^3x \sqrt{-g} [R - 2\Lambda]$$

Lecture 15: Curvatures

We try to develop a physical interpretation of the Ricci tensor. Note that this contains a *double derivative* of the metric, that is $\partial_\alpha \partial_\beta g_{\mu\nu}$.



Suppose we have some arbitrary smooth curve. Then, at each point of the curve, we can associate a circle whose first and second derivatives match with that of the curve. And there is always a radius associated with such a circle. Thus, to each point on the curve, we can associate a radius r , referred to as the *radius of curvature* of the curve at that point.

Now, for simplicity, let us assume that at a point P on the curve, we have drawn the circle and we shifted our origin to P . Thus, the equation of the circle becomes:

$$x^2 + y^2 = r^2$$

From the equation of the circle, we obtain,

$$y \frac{dy}{dx} + x = 0 \implies y \frac{d^2y}{dx^2} + \left(\frac{dy}{dx} \right)^2 + 1 = 0 \implies y = \frac{1 + \left(\frac{dy}{dx} \right)^2}{-\frac{d^2y}{dx^2}}$$

¹ Λ is called the *cosmological constant*. While deriving the equation, Einstein added this since addition of this constant led to a solution which conveyed that the Universe was *static*. However, after Hubble's discovery that the Universe is expanding, this constant was dropped. Then again, the constant was brought back since it was discovered that the Universe was apparently accelerating.

Also, from above, we have

$$\frac{dy}{dx} = -\frac{x}{y}$$

Then we obtain an expression for the radius of the circle:


$$\begin{aligned} r &= \left| \sqrt{x^2 + y^2} \right| = \left| y \sqrt{1 + \left(\frac{x}{y} \right)^2} \right| \\ &= \left| \frac{1 + \left(\frac{dy}{dx} \right)^2}{-\frac{d^2y}{dx^2}} \times \sqrt{1 + \left(\frac{x}{y} \right)^2} \right| \\ &= \left| \frac{\left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{3/2}}{\frac{d^2y}{dx^2}} \right| \end{aligned}$$

By choosing an appropriate coordinate system, we can always set the first derivative to zero ¹ and we have a result that,

$$r = \left| \frac{d^2y}{dx^2} \right|^{-1}$$

Thus, the second derivative represents the curvature of the curve, since it is intrinsically related to the radius of curvature. And since the Ricci/Riemann tensor contains double derivative of the metric, it somehow represents the curvature of the space. Hence, it is sometimes also called *Riemann curvature tensor*!

Now, for the sake of getting our hands dirty, let us calculate the Ricci scalar of $\mathbb{S}^2$ (sphere), of radius r_0 .



For this, we first need to calculate the Riemann tensor, then contract it to get Ricci tensor and then again contract it to get Ricci scalar. Thus, this seems a very involved task and we have to do that carefully!

As calculate before, the metric for the sphere is written as:

$$ds^2 = r_0^2 d\theta^2 + r_0^2 \sin^2 \theta d\phi^2$$

From there we get the metric tensor and inverse metric tensor as:

$$g_{\mu\nu} = \text{diag}(r_0^2, r_0^2 \sin^2 \theta) \quad g^{\mu\nu} = \text{diag}(1/r_0^2, 1/r_0^2 \sin^2 \theta)$$

Also, we had earlier found that only two of the Christoffel symbols are non-zero, namely,

$$\Gamma_{\phi\phi}^{\theta} = -\sin \theta \cos \theta \quad \Gamma_{\theta\phi}^{\phi} = \cot \theta$$

Calculating the Riemann Tensor

$$R^{\rho}_{\sigma\mu\nu} = \partial_{\mu}\Gamma^{\rho}_{\nu\sigma} + \Gamma^{\rho}_{\mu\lambda}\Gamma^{\lambda}_{\nu\sigma} - \partial_{\nu}\Gamma^{\rho}_{\mu\sigma} - \Gamma^{\rho}_{\nu\lambda}\Gamma^{\lambda}_{\mu\sigma}$$

¹Note that since first derivative roughly gives us the slope of the tangent, with proper rotation and shifting of the coordinates, we can indeed align our coordinate system along the tangent so that it's slope is zero

Let us take $\rho = \theta, \sigma = \phi, \mu = \theta, \nu = \phi$ first. Then we have:

$$\begin{aligned}
 R^\theta_{\phi\theta\phi} &= \partial_\theta \Gamma^\theta_{\phi\phi} + \Gamma^\theta_{\theta\lambda} \Gamma^\lambda_{\phi\phi} - \partial_\phi \Gamma^\theta_{\theta\phi} - \Gamma^\theta_{\phi\lambda} \Gamma^\lambda_{\theta\phi} \\
 &= -\partial_\theta \left(\frac{1}{2} \sin 2\theta \right) + \sin \theta \cos \theta \times \cos \theta \\
 &= -\frac{1}{2} \cos 2\theta + \sin \theta \cos \theta \times \frac{\cos \theta}{\sin \theta} \\
 &= \sin^2 \theta - \cos^2 \theta + \cos^2 \theta \\
 &= \sin^2 \theta
 \end{aligned}$$

In fact, this is the only non-zero Riemann tensor component. Other components will be zero owing to the anti-symmetric and other nice properties of the Riemann tensor.

Calculating the Ricci Tensor

$$R_{\mu\nu} = R^\lambda_{\mu\lambda\nu}$$

We calculate the components one by one.

$$R_{\theta\theta} = R^\lambda_{\theta\lambda\theta}$$

Note that we only found out one component of the Riemann tensor. We have to transform all quantities to that form only. Here, the only non-zero index will be $\lambda = \phi$, thus we get,

$$\begin{aligned}
 R_{\theta\theta} &= R^\phi_{\theta\phi\theta} \\
 &= g^{\phi\alpha} R_{\alpha\theta\phi\theta} \\
 &= g^{\phi\alpha} (-1)(-1) R_{\theta\alpha\theta\phi} \\
 &= g^{\phi\phi} R_{\theta\phi\theta\phi} \quad (\because g \text{ diagonal, so } \alpha = \phi) \\
 &= \frac{1}{r_0^2 \sin^2 \theta} g_{\theta\sigma} R^\sigma_{\phi\theta\phi} \\
 &= \frac{1}{r_0^2 \sin^2 \theta} r_0^2 R^\theta_{\phi\theta\phi} \quad (\because g \text{ diagonal, so } \sigma = \theta) \\
 &= 1
 \end{aligned}$$

$R_{\theta\phi} = 0$ due to the anti-symmetric property and $R_{\phi\phi} = R^\lambda_{\phi\lambda\phi} = R^\theta_{\phi\theta\phi} = \sin^2 \theta$. Now, we have obtained all the components of the Ricci tensor and hence, let us move on to the Ricci scalar.

$$R = R^\mu_{\mu} = g^{\mu\nu} R_{\mu\nu} = \frac{1}{r_0^2} \times 1 + \frac{1}{r_0^2 \sin^2 \theta} \times \sin^2 \theta = \frac{2}{r_0^2}$$

Thus we obtain that the Ricci scalar for the sphere is $\frac{2}{r_0^2}$. Thus, the Ricci scalar indeed refers to the usual curvature and its inverse represents the radius of curvature!

$$R^{-1} \sim r_0^2$$

Lecture 16: Bianchi Identities

16.1 Natural Units

Instead of human-defined units, like *metre* and *seconds* are somewhat problematic due since these are not fundamental. We could have easily defined '1 m' to be something else and worked with that. It is better to

use some fundamental quantities, appearing in Nature, to define a system of measurement. In the *natural units*, we use the speed of light and the Planck's constant and put them to be dimensionless and numerically equal to 1, that is,

$$c = 1 \implies 1\text{sec} = 3 \times 10^8 \text{ m}$$

$$\hbar = 1 \implies \frac{10^{-34} \text{ kg} \cdot \text{m}^2}{3 \times 10^8 \text{ m}} = 1 \implies 1 \text{ m} \approx 3 \times 10^{42} \text{ kg}^{-1}$$

Note that, since action has the same dimension as $\hbar$, action is actually dimensionless in natural units (which is cool!)

We will see two identities which will be very useful while deriving the Einstein tensor for the field equations. The starting of the derivations of both is the Jacobi identity, which is a general property satisfied by an *binary operation*.

Identity 3 (Jacobi Identity):

Let $\times$ be a binary operation and x, y, z be arbitrary three elements of a particular mathematical structure. Then the binary operation satisfies the Jacobi Identity if,

$$x \times (y \times z) + z \times (x \times y) + y \times (z \times x) = 0 \iff x \times (y \times z) + \text{cyc.}(x, y, z)$$

The Jacobi identity is satisfied by many binary operations, like the cross-product, the commutator bracket, the Poisson bracket, etc¹. We will consider the Jacobi identity for the commutator of the covariant derivatives here!

16.2 First Bianchi Identity

Let us apply the Jacobi identity on a scalar function ϕ .

$$[\nabla_\rho, [\nabla_\mu, \nabla_\nu]] \phi + \text{cyc.}(\rho, \mu, \nu) = 0$$

Now note the following,

$$[\nabla_\mu, \nabla_\nu] \phi = \nabla_\mu(\nabla_\nu \phi) - \mu \leftrightarrow \nu = [\partial_\mu \partial_\nu \phi - \Gamma_{\mu\nu}^\lambda \partial_\lambda \phi] - \mu \leftrightarrow \nu$$

Since the term in the bracket is symmetric in μ and ν , it cancels with the second term ² and hence the commutator acting on a scalar field is zero. With this, the initial equation simplifies as,

$$\begin{aligned} 0 &= -[\nabla_\mu, \nabla_\nu] (\nabla_\rho \phi) + \text{cyc.}(\rho, \mu, \nu) \\ &= -R_{\rho\sigma\mu\nu} \partial^\sigma \phi + \text{cyc.}(\rho, \mu, \nu) \\ &= (R_{\sigma\rho\mu\nu} + \text{cyc.}(\rho, \mu, \nu)) \partial^\sigma \phi \end{aligned}$$

Since the scalar field is arbitrary, we have the first identity as:

$$R_{\sigma\rho\mu\nu} + R_{\sigma\nu\rho\mu} + R_{\sigma\mu\nu\rho} = 0$$

Contracting the above with $g^{\lambda\sigma}$, we have:

$$R^\lambda_{\rho\mu\nu} + R^\lambda_{\nu\rho\mu} + R^\lambda_{\mu\nu\rho} = 0$$

¹All these are particular cases of a broadened thing called *Lie Bracket*

²Note that in spaces where torsion is there, partial derivatives do not commute and we do not get this result!

With the first Bianchi identity, we can derive a nice property of the Riemann tensor. From the identity, we have:

$$\begin{aligned}
 R_{\rho\sigma\mu\nu} &= -R_{\rho\nu\sigma\mu} - R_{\rho\mu\nu\sigma} \\
 &= R_{\nu\rho\sigma\mu} + R_{\mu\rho\nu\sigma} \\
 &= -R_{\nu\mu\rho\sigma} - R_{\nu\sigma\mu\rho} - R_{\mu\sigma\rho\nu} - R_{\mu\nu\sigma\rho} \quad (\text{using Bianchi identity}) \\
 &= 2R_{\mu\nu\rho\sigma} + (R_{\sigma\nu\mu\rho} + R_{\sigma\mu\rho\nu}) \\
 &= 2R_{\mu\nu\rho\sigma} + (R_{\sigma\nu\mu\rho} + R_{\sigma\mu\rho\nu} + R_{\sigma\rho\nu\mu}) - R_{\sigma\rho\nu\mu} \\
 &= 2R_{\mu\nu\rho\sigma} + 0 - R_{\sigma\rho\mu\nu}
 \end{aligned}$$

From this, we obtain that the Riemann tensor is also symmetric under exchange of two pair of indices, that is,

$$R_{(\rho\sigma)(\mu\nu)} = R_{(\mu\nu)(\rho\sigma)}$$

Recall that the Ricci tensor was defined as the contraction of first and third index of the Riemann tensor.

$$R_{\mu\nu} = g^{\alpha\beta} R_{\alpha\mu\beta\nu} = g^{\beta\alpha} R_{\beta\nu\alpha\mu} = R_{\nu\mu}$$

Hence we obtain that the Ricci tensor is a symmetric tensor.

16.3 Second Bianchi Identity

For this, we apply the Jacobi identity on a vector field V^ρ .

$$\begin{aligned}
 0 &= \nabla_\sigma([\nabla_\mu, \nabla_\nu] V^\rho) - [\nabla_\mu, \nabla_\nu](\nabla_\sigma V^\rho) + \text{cyc.}(\sigma, \mu, \nu) \\
 &= \nabla_\sigma(R^\rho_{\lambda\mu\nu} V^\lambda) - R^\rho_{\lambda\mu\nu} \nabla_\sigma V^\lambda - R_{\sigma\lambda\mu\nu} \nabla_\lambda V^\rho + \text{cyc.}(\sigma, \mu, \nu) \quad (\text{applying commutator on } (1, 1) \text{ tensor}) \\
 &= [\nabla_\sigma(R^\rho_{\lambda\mu\nu}) V^\lambda + \cancel{R^\rho_{\lambda\mu\nu} \nabla_\sigma(V^\lambda)}] - \cancel{R^\rho_{\lambda\mu\nu} \nabla_\sigma V^\lambda} - g^{\beta\lambda} R_{\sigma\beta\mu\nu} \nabla_\lambda V^\rho + \text{cyc.}(\sigma, \mu, \nu) \quad (\because \text{Liebniz rule}) \\
 &= \nabla_\sigma(R^\rho_{\lambda\mu\nu}) V^\lambda + g^{\beta\lambda} R_{\sigma\beta\mu\nu} \nabla_\lambda V^\rho + \text{cyc.}(\sigma, \mu, \nu) \\
 &= [\nabla_\sigma(R^\rho_{\lambda\mu\nu}) + \text{cyc.}(\sigma, \mu, \nu)] V^\lambda + \left[\cancel{R^\lambda_{\sigma\mu\nu} + \text{cyc.}(\sigma, \mu, \nu)} \right] \nabla_\lambda V^\rho \quad (\text{first Bianchi identity}) \\
 &= (\nabla_\sigma(R_{\rho\lambda\mu\nu}) + \nabla_\nu(R_{\rho\lambda\sigma\mu}) + \nabla_\mu(R_{\rho\lambda\nu\sigma})) V^\lambda
 \end{aligned}$$

Thus, we obtain the second Bianchi identity,

$$\nabla_\sigma(R_{\rho\lambda\mu\nu}) + \nabla_\nu(R_{\rho\lambda\sigma\mu}) + \nabla_\mu(R_{\rho\lambda\nu\sigma}) = 0$$

If we contract the equation with $g^{\rho\mu}$,

$$\begin{aligned}
 0 &= g^{\rho\mu}(\nabla_\sigma(R_{\rho\lambda\mu\nu}) + \nabla_\nu(R_{\rho\lambda\sigma\mu}) + \nabla_\mu(R_{\rho\lambda\nu\sigma})) \\
 &= \nabla_\sigma(R^\mu_{\lambda\mu\nu}) + \nabla_\nu(R^\mu_{\lambda\sigma\mu}) + \nabla^\rho(R_{\rho\lambda\nu\sigma}) \\
 &= \nabla_\sigma(R_{\lambda\nu}) - \nabla_\nu(R_{\lambda\sigma}) + \nabla^\rho(R_{\rho\lambda\nu\sigma})
 \end{aligned}$$

Contracting again with $g^{\lambda\sigma}$, we obtain:

$$\begin{aligned}
 0 &= g^{\lambda\sigma}(\nabla_\sigma(R_{\lambda\nu}) - \nabla_\nu(R_{\lambda\sigma}) + \nabla^\rho(R_{\rho\lambda\nu\sigma})) \\
 &= \nabla^\lambda R_{\lambda\nu} - \nabla_\nu R + \nabla^\rho R_{\rho\nu} \\
 &= \nabla^\lambda R_{\lambda\nu} - \nabla_\nu R + \nabla^\lambda R_{\lambda\nu} \\
 &= 2\nabla^\lambda R_{\lambda\nu} - g_{\lambda\nu} \nabla^\lambda R \\
 &= \nabla^\lambda \left(R_{\lambda\nu} - \frac{1}{2} g_{\lambda\nu} R \right)
 \end{aligned}$$

If we define a new tensor as follows,

$$G_{\lambda\nu} := R_{\lambda\nu} - \frac{1}{2}g_{\lambda\nu}R$$

then, we get the fact that:

$$\nabla^\lambda G_{\lambda\nu} = 0$$

Thus, the tensor $G_{\lambda\nu}$ is covariantly conserved, similar to the metric, as seen before. The tensor is called the *Einstein tensor* and will appear in the field equations to be seen later.