


General Theory Of Relativity and Cosmology

LECTURE NOTES

Instructor: Prof. Golam Mortuza Hossain



Sagnik Seth

**Dept of Physical Sciences
IISER Kolkata**

Contents

Lecture 01: Introduction	3
Lecture 02: Galilean Stuffs	4
2.1 Galilean Transformation	4
2.2 A Big Blow to Galileo	6
Lecture 04: Vectors and Stuffs	6
Lecture 05: Some more Vectors and Stuffs	8
5.1 Gradient Vector	9
5.2 Tensors	9

Lecture 01: Introduction

In the physical sense, the word 'relativity' can be used to relate the measurement of one observer with that of another, for the same object/quantity. So using relativity, we can characterise the difference in the observations of two observers in two different frame of reference.

Note that, physical laws should not matter based on who is observing (that is, physical laws are *universal*), hence relativity becomes very important when dealing with the resolution of conflicts.

Let us consider two reference frames, and let v be the relative velocity between the frames.





Figure 1: Two frames of reference with a relative velocity v

Fig. 1 shows two reference frames. The moving frame is denoted by O' (henceforth, primes will generally denote moving frames). If the relative velocity v between the two frames remain constant, then it falls under the domain of *Special Relativity* and if unfortunately (or fortunately) it doesn't, then it comes under *General Relativity*.


The theory of relativity dates back to the inception of Newtonian mechanics and hence, let us review the laws in brief (without being much technical 😊):

- **The First Law** : $\approx$ velocity of a body remains constant unless acted by an external, unbalanced force.
- **The Second Law** : $\approx \vec{F} = m\vec{a}$ where $\vec{F}$ is the external force and $\vec{a}$ is the acceleration.
- **The Third Law** : $\approx$ Every action has an equal and opposite reaction. Note that for this, two bodies are needed.



From the figure above, we will have $\vec{F}_{21} = -\vec{F}_{12}$

- **The Gravitational Law** : $\approx$



In the above situation, if we consider for mass m_1 , we have:

$$\vec{F}_{12} = G \frac{m_1 m_2}{r_{21}^2} \hat{r}_{21}$$

An important observation: in the second law, if we take the force to be zero, then apparently $\vec{a} = \frac{d\vec{v}}{dt} = 0 \implies \vec{v} = \text{const.}$ which is the statement of the first law. However, the second law does not imply the first law, since the first law defines what an *inertial frame* is and the statement of the second law applies only in case of inertial frames. So, a better formulation of the second and third law can be like, "In a frame where the first law is valid, blah blah blah..."

Lecture 02: Galilean Stuffs

In the previous lecture, we saw how Newton's laws define an inertial frame and how we can rephrase the second and third law in terms of inertial frames, to avoid confusion. It happens that, there are reference frames where the first law doesn't hold true.

Imagine a person inside a darkened car, isolated from the outside world, with a ball hanging from the roof of the car. The car suddenly starts accelerating and (as intuition speaks) the ball starts moving. The person claims "The ball has moved!" 🧠.

Note that from the perspective of the person, no force has been applied to the ball, yet it moved, which is a direct violation of the second law. Thus, we can call this frame *non-inertial*.

Now, consider a person inside an ill-fated lift, tragically falling 😊. Everything in the lift frame is accelerating together under gravity and hence 'weightlessness' (free-fall) occurs. If somehow a coin is also there in the lift, it simply floats. And if the person pushes the coin, it moves with an almost constant velocity. Thus, it is a very good approximation to an *inertial frame* (although, extremely tragic for the person!).




Figure 2: A sad guy floating with a coin in a falling lift

We had always wanted the physical laws to stay the same. This is actually the statement of Galilean relativity:

"Laws of physics (nature) must be the same in all inertial frames of reference"

By physical laws, we imply Newton's laws of mechanics and the law of gravitation. To have a more mathematical (and less philosophical) aspect to the above statement, we define the *Galilean transformation*.

2.1 Galilean Transformation

Let an object be at point P and consider two reference frames O and O' , which coincided at $t = 0$ but have moved apart since, along the common x-axis. (Henceforth, reference frames will simply be termed *observers*).




Figure 3: Observing same point from two different frames

Now, the distance between the origins of two frames, at time t , will be $R = vt$. Then accordingly, we can write the relation between the coordinates of the two observers as follows:

$$\begin{aligned} x' &= x - vt \\ y' &= y \\ z' &= z \\ t' &= t \end{aligned} \quad \begin{pmatrix} t' \\ x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ -v & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} t \\ x \\ y \\ z \end{pmatrix}$$

An important assumption: The Galilean transformation assumes a *universal time*, that is, time elapsed on clocks in both frames are the same ¹.

Now, it is not necessary for the observers to move along the x axis only. Hence, generalising the transformation to arbitrary direction, we obtain:

$$t' = t \quad \mathbf{r}' = \mathbf{r} - \mathbf{v}t$$

Now, let us look how the physical laws react under the Galilean transformation.

▪ **Law of Gravitation:**

Consider the situation in the diagram below:




Figure 4: Gravitational law from two different frames

According to Galilean transformation, we have:

$$\begin{aligned} \mathbf{z} &= \mathbf{r}_2 - \mathbf{r}_1 \\ &= (\mathbf{r}'_2 + \mathbf{v}t) - (\mathbf{r}'_1 + \mathbf{v}t) \\ &= \mathbf{r}'_2 - \mathbf{r}'_1 \\ &= \mathbf{z}' \end{aligned}$$

¹This was perhaps a fair assumption since time, atleast in the philosophical aspect, has always seemed to be *superior*.

Thus, we see that the vector $\hat{\mathbf{z}}$ is unchanged and since this is the only vector appearing in Newton's law of gravitation, we have:

$$\mathbf{F}_{12} = G \frac{m_1 m_2}{r^2} \hat{\mathbf{z}} = G \frac{m_1 m_2}{r'^2} \hat{\mathbf{z}}' = \mathbf{F}'_{12}$$

We see that the expression of the force does not change between frames, thus indicating a *universality*.

▪ Second Law:

Using the second law, we can write

$$\mathbf{F} = m \frac{d^2 \mathbf{r}}{dt^2}$$

Now, using Galilean transformation, we obtain:

$$\mathbf{r}(t) = \mathbf{r}'(t') + \mathbf{v}t \implies \frac{d\mathbf{r}}{dt} = \frac{d\mathbf{r}'}{dt} \times \frac{dt'}{dt} + \mathbf{v} \implies m \frac{d^2 \mathbf{r}}{dt^2} = m \frac{d^2 \mathbf{r}'}{dt'^2}$$

Thus, we see that in general, force will be the same for observers in different inertial frames. Note that we assumed the *time universality*, that is, $\frac{dt'}{dt} = 1$

▪ Third Law:

Note that since the force expression for gravity is symmetric in m_1 and m_2 , the law of gravitation automatically incorporates the third law. Newton knew only about gravity and perhaps, third law would not have been necessary if he had relied only on gravity, however, he postulated that for all other forces, third law holds.

2.2 A Big Blow to Galileo

Let us consider that in Fig. 3, point P is also moving with some velocity $\mathbf{u}'$ and $\mathbf{u}$ in primed and unprimed frame respectively. Then, we can write:

$$\frac{d\mathbf{u}}{dt} = \frac{d}{dt}(\mathbf{r}' + \mathbf{v}t) = \frac{d\mathbf{r}'}{dt} + \mathbf{v} = \mathbf{u}' + \mathbf{v} \implies \mathbf{u} = \mathbf{u}' + \mathbf{v} \quad : \text{velocity addition formula}$$

However, observations, specially the Michelson-Morley experiment, were made in different frames but found the speed of light (henceforth denoted as c) to be a constant, which contradicted the velocity addition formula when applied to speed of light.

Since the velocity addition formula depended entirely on the Galilean transformation, to resolve this conflict, we have to change the transformation rule in its entirety.

Lecture 04: Vectors and Stuffs

Let us consider rotation about the z -axis by an angle θ , in which case, the x and y components transform as:

$$\begin{aligned} x' &= x \cos \theta + y \sin \theta \\ y' &= -x \sin \theta + y \cos \theta \end{aligned}$$

Then, we can write the infinitesimal change in the components in the following form:

$$\begin{pmatrix} dx' \\ dy' \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

Note that, this can also be written as:

$$\begin{pmatrix} dx' \\ dy' \end{pmatrix} = \begin{pmatrix} \frac{\partial x'}{\partial x} & \frac{\partial x'}{\partial y} \\ \frac{\partial y'}{\partial x} & \frac{\partial y'}{\partial y} \end{pmatrix} \begin{pmatrix} dx \\ dy \end{pmatrix}$$

The matrix in this case is the *Jacobian matrix* and this is a more general structure of coordinate transformations and hence any transformation can be written like this, even the Galilean or the Lorentz transformations.

This shows that transformations are just multiplication of a matrix with a vector. To this extent, we define a *vector* ${}^1\bar{x}$ with the following components:

$$\bar{x} = (x^0, x^1, x^2, x^3)^T$$

Here $x^0 = ct$ and $x^1 = x, x^2 = y, x^3 = z$. Note that since t and x are not dimensionally same, we took $x^0 = ct$ to make it in terms of distance. We multiplied by c (and not any other velocity) because c is a universal constant.

We can express any infinitesimal transformation as:

$$\begin{pmatrix} dx'^0 \\ dx'^1 \\ dx'^2 \\ dx'^3 \end{pmatrix} = \begin{pmatrix} \frac{\partial x'^0}{\partial x^0} & \cdots & \frac{\partial x'^0}{\partial x^3} \\ \vdots & \ddots & \vdots \\ \frac{\partial x'^3}{\partial x^0} & \cdots & \frac{\partial x'^3}{\partial x^3} \end{pmatrix} \begin{pmatrix} dx^0 \\ dx^1 \\ dx^2 \\ dx^3 \end{pmatrix}$$

In a more compact way, we can write in the vector-matrix form or in the component form using Einstein's summation convention:

$$d\bar{x}' = \Lambda d\bar{x} \iff dx'^\mu = \Lambda^\mu{}_\nu dx^\nu$$

where $\Lambda^\mu{}_\nu = \frac{\partial x'^\mu}{\partial x^\nu}$ is an element of the *matrix* Λ .

Summation convention: We drop the summation sign if twice repeated indices are there on the same side of the equation.

Now, let us define what a vector is:

Definition 1 (Vector):

If a mathematical object transforms under a coordinate transformation like a differential $d\bar{x}$, that is, $d\bar{x} \rightarrow d\bar{x}' = \Lambda d\bar{x}$, then it is called a *contravariant vector* or a *contravector* or a *tensor of rank (1, 0)*

Let us show that under the Galilean transformation, velocity, as defined by $\mathbf{u} = \frac{d\mathbf{r}}{dt}$ is a vector. We will define everything in contravariant and four-vector notation. For this, we consider the vector defined as before:

$$d\bar{x} = (ct, x, y, z)^T \equiv (x^0, x^1, x^2, x^3)^T$$

The Galilean transformation is defined by:

$$\lambda \equiv \begin{pmatrix} 1 & 0 & 0 & 0 \\ -v^1/c & 1 & 0 & 0 \\ -v^2/c & 0 & 1 & 0 \\ -v^3/c & 0 & 0 & 1 \end{pmatrix} \iff \begin{aligned} x'^0 &= x^0 \\ x'^1 &= x^1 - (v^1/c)x^0 \\ x'^2 &= x^2 - (v^2/c)x^0 \\ x'^3 &= x^3 - (v^3/c)x^0 \end{aligned}$$

¹This is often called a *four vector*

We can then define the four velocity as¹:

$$u^\mu \equiv \frac{dx^\mu}{dt} = (c, u^1, u^2, u^3)^T$$

Now, we have:

$$\begin{aligned} dx'^\mu &= \frac{\partial x'^\mu}{\partial x^\nu} dx^\nu \\ \Rightarrow u'^\mu dt' &= \frac{\partial x'^\mu}{\partial x^\nu} u^\nu dt \quad (\text{as } dt = dt') \\ \Rightarrow u'^\mu &= \frac{\partial x'^\mu}{\partial x^\nu} u^\nu \end{aligned}$$

Thus, we see that the components of the velocity vector transform similar to the differential displacement and hence, velocity is indeed a vector under Galilean transformation. In a similar way, we can show that acceleration $\mathbf{a} = \frac{d\mathbf{v}}{dt}$ is also a vector under Galilean transformation.

Now, let us turn to Lorentz transformation. Things become bad here! For example, consider the same definition of velocity $\mathbf{u} = \frac{d\mathbf{x}}{dt}$. However, here t does not remain invariant under coordinate transformation. Then we will have:

$$u'^\mu = \frac{dx'^\mu}{dt'} = \frac{\Lambda^\mu{}_\nu x^\nu}{\Lambda^0{}_\alpha dx^\alpha} \neq \frac{dx^\mu}{dt}$$

Thus, this definition fails to transform similar to a differential displacement and hence, the way it is defined presently, velocity is not a vector 😞.

Lecture 05: Some more Vectors and Stuffs

Previously, we saw how a vector was defined in terms of the transformation of the differential displacements. In here, we will consider another situation.

First, note that, using a given transformation,

$$dx'^\mu = \Lambda^\mu{}_\nu dx^\nu$$

we can define the inverse transformation,

$$dx^\nu = (\Lambda^{-1})^\nu{}_\mu dx'^\mu$$

The above statement is valid only if the transformation matrix is invertible (in most case, transformations like translations, rotations, etc. are invertible).

Let us now see what the inverse transformation matrix looks like. For that, consider:

$$\begin{aligned} dx'^\mu &= \frac{\partial x'^\mu}{\partial x^\nu} dx^\nu \\ \Rightarrow \frac{\partial x^\beta}{\partial x'^\mu} dx'^\mu &= \frac{\partial x^\beta}{\partial x'^\mu} \underbrace{\frac{\partial x'^\mu}{\partial x^\nu}}_{\delta^\beta{}_\nu} dx^\nu \\ \Rightarrow dx^\beta &= \frac{\partial x^\beta}{\partial x'^\mu} dx'^\mu \end{aligned}$$

Thus we see that:

$$(\Lambda^{-1})^\nu{}_\mu \equiv \frac{\partial x^\nu}{\partial x'^\mu}$$

¹Note that four velocity is defined as derivative with respect to something called as *proper time*, however, for Galilean transformations, proper time coincides with the coordinate time and hence this definition is okay.

5.1 Gradient Vector

In the Cartesian coordinates, we all know:

$$\nabla = \hat{\mathbf{i}} \frac{\partial}{\partial x} + \hat{\mathbf{j}} \frac{\partial}{\partial y} + \hat{\mathbf{k}} \frac{\partial}{\partial z}$$

In line with this, we will define another quantity (often called *four gradient*) such that:

$$\nabla_\mu := \frac{\partial}{\partial x^\mu}$$

A few points on this:

1. Note that we wrote the index in the down position for ∇_μ . This is done since we will see that the four gradient does not follow the transformation rule of a vector, hence it is a different entity. Since in vectors we put the indices upwards, to differentiate it from a vector, we put it down.
2. In spherical or cylindrical or any other coordinates, the *ordinary* gradient is not just the partial derivatives of the coordinates. There are some obnoxious prefactors involved also. Irrespective of that, we define the four gradient as just the derivative of the coordinates ('coz later we will see that those prefactors are just some jugglery with the *metric*).

The four gradient is better denoted by ∂_μ and thus, we have:

$$\bar{\partial} \equiv (\partial_0 \quad \partial_1 \quad \partial_2 \quad \partial_3) = \left(\frac{1}{c} \frac{\partial}{\partial t} \quad \frac{\partial}{\partial x} \quad \frac{\partial}{\partial y} \quad \frac{\partial}{\partial z} \right)$$

Under a coordinate transformation $x^\mu \rightarrow x'^\mu = \Lambda^\mu{}_\nu x^\nu$, we have using chain rule:

$$\partial'_\mu = \frac{\partial x^\nu}{\partial x'^\mu} \frac{\partial}{\partial x^\nu} \implies \partial'_\mu = (\Lambda^{-1})^\nu{}_\mu \frac{\partial}{\partial x^\nu}$$

The four gradient apparently transform accordingly as $\bar{\partial}' = \Lambda^{-1} \bar{\partial}$ which is contrary to that of the vector.

Definition 2:

Covector If a mathematical object transforms under a coordinate transformation like the derivative $\bar{\partial}$, then we call it a *covariant vector* or *covector* or a *tensor of rank (0,1)*

Actually vectors and covectors are defined based how the basis vectors of the space transform. Vectors transform in a way *contrary* to the basis vectors and hence are called *contravariant* while covectors transform in a way similar to how the basis transforms and hence called *covariant*.¹

5.2 Tensors

Consider the product,

$$\omega = \bar{\mathbf{x}} \bar{\mathbf{x}}^T \equiv \begin{pmatrix} dx^0 \\ dx^1 \\ dx^2 \\ dx^3 \end{pmatrix} (dx^0 \quad dx^1 \quad dx^2 \quad dx^3) = \begin{pmatrix} dx^0 dx^0 & \dots & dx^0 dx^3 \\ \vdots & \ddots & \vdots \\ dx^3 dx^0 & \dots & dx^3 dx^3 \end{pmatrix}$$

Note that, defined like this, ω represents a square matrix and thus, has two indices. This operation is often called *outer product* and is denoted as

$$\omega = \bar{\mathbf{x}} \otimes \bar{\mathbf{x}} \quad \omega^{\mu\nu} = dx^\mu dx^\nu$$

¹In the elegant language of differential geometry, vectors are actually the *tangent vectors* belonging to the *tangent space* of the space (*manifold*) while covectors belong to the *dual space* of the *tangent space*.

Now, let us see how it transforms:

$$\begin{aligned}\omega^{\mu\nu} &= dx'^{\mu} dx'^{\nu} \\ &= \left(\frac{\partial x'^{\mu}}{\partial x^{\alpha}} dx^{\alpha} \right) \left(\frac{\partial x'^{\nu}}{\partial x^{\beta}} dx^{\beta} \right) \\ &= \left(\frac{\partial x'^{\mu}}{\partial x^{\alpha}} \right) \left(\frac{\partial x'^{\nu}}{\partial x^{\beta}} \right) dx^{\alpha} dx^{\beta}\end{aligned}$$

This is yet another kind of transformation, different from either vector and covector.

Definition 3:

Tensor If a mathematical object transforms, under a coordinate transformation, like $\underbrace{d\bar{x} \otimes \dots \otimes d\bar{x}}_m$ and like $\underbrace{\bar{\partial} \otimes \dots \otimes \bar{\partial}}_n$, then it is called a *tensor of rank (m,n)*