


Quantum Field Theory

LECTURE NOTES

Instructor: Prof. Ritesh Kumar Singh



Sagnik Seth

Dept of Physical Sciences
IISER Kolkata


Contents

Lecture 01: Intro and Quantum SHO	3
Lecture 02: Dimensional Deep Dive	4
Lecture 03: From Classical to Quantum	7
3.1 Time Evolution	7
3.1.1 Free Particle	8
3.1.2 Continuity Equation	9
Lecture 04: Classical Fields	9
4.1 Our actions tell a lot!	11
Lecture 05 & 06: Looking into Hamiltonians	12
6.1 Functional Derivatives	13
6.2 Hamilton's Equations of Motion	14
6.3 Poisson Bracket	14
Lecture 07 & 08: Representations of Rotation and Lorentz group	15
Lecture 09: Noether's Theorem	15
9.1 Examples: Spacetime Symmetries	18
9.1.1 Translational Invariance	18
9.1.2 Lorentz Invariance	19
9.2 Examples: Internal Symmetries	20
9.2.1 Global $U(1)$ Symmetry of Complex Scalar Field	20
Lecture 10 & 11: Field Quantisation	21
11.1 Conserved Charges	23
11.2 Normal Ordering	24
11.3 Constructing the Hilbert Space	24
Lecture 12: Making things invariant	25
12.1 Invariant Inner Product	25
12.2 Invariant Measure	26
12.3 Position states?	27
Lecture 13: Heisenberg Picture	28
Lecture 14 & 15: Propagators and Causality	28

Lecture 01: Intro and Quantum SHO

*People do 'weird' stuffs for earning,
You can do the same for learning!*

Many physical theories have been proposed and it is imperative to see how each theory is linked with the previous in some way.



From the proposed theory of everything, we can transform to a *relativistic* quantum field theory which will be the topic of discussion for most part. When subjected to *decoherence*, a relativistic QFT collapse to a classical QFT as noise hides the ‘quantumness’ from consideration. Hence we will start our discussion from classical fields and then move to quantum fields.

Since the foundational aspect of QFT (and many other topics in Physics) is the Harmonic Oscillator, let us discuss the quantum Harmonic oscillator for introduction. For that, note that the classical Hamiltonian for the harmonic oscillator is given by:

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega^2 x^2$$

We do not like the things like m and ω which prevent us from seeing things clearly 😊. Hence, we invoke the holy action of making things dimensionless. For that, we define:

$$\begin{aligned} X &= \sqrt{\frac{m\omega}{\hbar}} x & \longrightarrow & \quad x^2 = \frac{\hbar}{m\omega} X^2 \\ P &= \frac{1}{\sqrt{m\omega\hbar}} p & \longrightarrow & \quad p^2 = m\omega\hbar P^2 \end{aligned}$$

Substituting these in the Hamiltonian, we have:

$$H = \frac{\hbar\omega}{2} (X^2 + P^2)$$

Note that, we are still within the classical domain. Now, let us elevate x and p to operators and we define:

$$[x, p] = i\hbar \mathbb{1} \implies [X, P] = \sqrt{\frac{m\omega}{\hbar}} \frac{i\hbar \mathbb{1}}{\sqrt{m\omega\hbar}} = i$$

Introducing the commutator bracket brings us to the quantum world. Now, we invoke our very own ladder operators:

$$\begin{aligned} \hat{a} &= \frac{1}{\sqrt{2}} (\hat{X} + i\hat{P}) & : \text{annihilation operator} \\ \hat{a}^\dagger &= \frac{1}{\sqrt{2}} (\hat{X} - i\hat{P}) & : \text{creation operator} \end{aligned}$$

Then we will have ¹:

$$[a, a^\dagger] = \frac{1}{2} [X + iP, X - iP] = \frac{-i}{2} ([X, P] - [P, X]) = -i \times i = 1$$

¹henceforth, forsaking the hat symbol and identity operator $\mathbb{1}$, since they cause nothing but trouble, when the context is clear

Also, note that:

$$a^\dagger a = \frac{1}{2}(X^2 + P^2 + i \underbrace{(XP - PX)}_i) = \frac{1}{2}(X^2 + P^2) - \frac{1}{2} \implies H = \frac{\hbar\omega}{2}(a^\dagger a + \frac{1}{2})$$

Let us consider the *complete set of commuting observables* (CSCO) for this problem. Evidently, the set $\{H\}$ itself satisfies the condition since the eigenvalues are all non-degenerate (hence, we can label each state with only one index). To understand why, let us consider the action of the annihilation operator on a (normalised) state $|\psi\rangle$. For that we note the following:

$$\begin{aligned} [a^\dagger a, a] &= -a \implies [H, a] = -\hbar\omega a \\ [a^\dagger a, a^\dagger] &= a^\dagger \implies [H, a^\dagger] = \hbar\omega a^\dagger \end{aligned}$$

Now, we have:

$$Ha|\psi\rangle - aH|\psi\rangle = [H, a]|\psi\rangle = -\hbar\omega a|\psi\rangle \implies H(a|\psi\rangle) = (E - \hbar\omega)(a|\psi\rangle)$$

Thus, if $|\psi\rangle$ has an energy eigenvalue E , then $a|\psi\rangle$ will have an energy eigenvalue $E - \hbar\omega$. Thus, starting from any energy state, we can change to another state with energy reduced by one unit of $\hbar\omega$, using the annihilation operator. Similarly, we will have:

$$H(a^\dagger|\psi\rangle) = (E + \hbar\omega)(a^\dagger|\psi\rangle)$$

Let us denote the states $a^\dagger|\psi\rangle$ and $a|\psi\rangle$ by $|\psi_+\rangle$ and $|\psi_-\rangle$ respectively. Then, we will have

$$\langle\psi_-|\psi_-\rangle = \langle\psi|a^\dagger a|\psi\rangle = \langle\psi|\left(\frac{H}{\hbar\omega} - \frac{1}{2}\right)|\psi\rangle$$

Now, since $|\psi_-\rangle$ is a valid vector of the Hilbert space, its norm must be non-negative and finite. Hence, we obtain the condition:

$$0 \leq \frac{E}{\hbar\omega} - \frac{1}{2} < \infty \implies \frac{\hbar\omega}{2} \leq E$$

Hence, we get a lower bound on the energy eigenvalue, that is, there exists a state $|\psi_{\min}\rangle$ such that $H|\psi_{\min}\rangle = E_{\min}|\psi_{\min}\rangle$ where $E_{\min} = \frac{\hbar\omega}{2}$. Then, starting from this state, if we apply the creation operator, we will get successively increasing energies (and hence the system is non-degenerate). We thus obtain:

$$E_n = \left(n + \frac{1}{2}\right)\hbar\omega \quad n = 0, 1, \dots$$

Lecture 02: Dimensional Deep Dive

For the quantum harmonic oscillator, we had seen that how making things dimensionless eases calculations a bit. We consider three fundamental constants, each occupying their own special place in their field of action.

$$\begin{aligned} c &\equiv [LT^{-1}] && : \text{relativity} \\ \hbar &\equiv [L^2MT^{-1}] && : \text{quantum} \\ G &\equiv [L^3M^{-1}T^{-2}] && : \text{gravity} \end{aligned}$$

Lengths are tractable for us, since we can see the ‘length’, same goes for mass, atleast we can ‘feel’ it. However, time is an *enigma*. *We can't hold two ends of time at the same time* unlike holding two ends of a rod to measure its length. The absolute truth is: *Time passes!*

Note that, in physics, we are mostly concerned with equations like $E = mc^2$ and $E = \hbar\omega$. To that extend, we define the natural units:

$$c = 1$$

$$\hbar = 1$$

We want these quantities to be numerically equal to 1 and dimensionless. Note that, we had also done this kind of things before. When writing Newton's law, we had said that

$$F \propto m, F \propto a \implies F \propto ma \implies F = kma \quad \text{for some } k$$

Now, we chose unit and dimension of force in a way such that $k = 1$ (dimensionless and unit value) which gave us the celebrated law.

Note that:

- Making c dimensionless:

$$[LT^{-1}] \equiv 1 \implies [L] = [T]$$

- Making $\hbar$ dimensionless:

$$[L^2MT^{-1}] = [L^2ML^{-1}] = [LM] \equiv 1 \implies [L] = [M^{-1}]$$

Hence, we see that length and time are equivalent while mass and length have inverse relation. In this natural units, we have that energy is equivalent to mass and any other unit can be represented in terms of mass. Thus, by our convention, we choose mass or energy as the only important dimension.

Note that

$$\hbar c \equiv Jm = 1 \quad (\text{in natural units})$$

From this, we can say heuristically that increasing length scale is decreases the energy (mass) scale. We mainly use the length scale in context of *de Broglie wavelength*.

This is perhaps not the only way to define natural units. In cosmology, G plays a more important role and hence it is better to set $G = 1$, leaving aside $\hbar$. Using this *natural units*, we find:

$$[L] = [M]$$

$$[L] = [T]$$

Here, we see that length and mass scale are directly related. Well, in this regard, we treat the length scale to be that of the Schwarzschild radius of a blackhole ¹ which intuitively grows with mass. Since the above two natural units are widely distinct, there is as such no problem, however, in the unfortunate case where we have to consider both de Broglie wavelength and the Schwarzschild radius (that is, in the infamous domain of quantum gravity 🤖), one needs to be very careful.

Theorem 1 (π -Theorem):

Let $q_1, \dots, q_n$ be n variables which are physically relevant to a problem and which are related by an expression, that is,

$$F(q_1, \dots, q_n) = 0 \iff q_i = \tilde{F}(q_1, \dots, \hat{q}_i, \dots, q_n)$$

If k is the number of fundamental dimensions required to describe the n variables, then we can group these in $(n - k)$ groups of dimensionless variables $\Pi_1, \dots, \Pi_{n-k}$ such that for some f , we have:

$$f(\Pi_1, \dots, \Pi_{n-k}) = 0 \iff \Pi_i = \tilde{f}(\Pi_1, \dots, \hat{\Pi}_i, \dots, \Pi_{n-k})$$

.

¹This crap is the radius of an object such that if the body is squeezed to a radius lesser than the Schwarzschild radius, the gravitational attraction between the constituents of the body causes its irreversible collapse, turning it to a black hole



The theorem seems a bit vague (and pointless too). Let us take a physical example. Consider a spherical ball in a viscous fluid. The variable in the problem are:

$$\begin{aligned} \text{Drag force: } q_1 &\rightarrow F \quad [MLT^{-2}] \\ \text{Sphere diameter: } q_2 &\rightarrow d \quad [L] \\ \text{Fluid density: } q_3 &\rightarrow \rho \quad [ML^{-3}] \\ \text{Fluid velocity: } q_4 &\rightarrow v \quad [LT^{-1}] \\ \text{Fluid viscosity: } q_5 &\rightarrow \eta \quad [ML^{-1}T^{-1}] \end{aligned}$$

So, there are 5 such parameters and only three units viz. M, L, T are needed to describe them. Hence we will have two Π groups. It is a good thing to choose the repeating variables (variables which will be in both groups) that relate to mass, geometry and the kinematics of the problem. Also, note that since the Π groups are dimensionless, we can take the non-repeating variable's power to be 1. Hence, in this problem we choose them to be ρ, d, v . Thus, we will have:

$$\begin{aligned} \Pi_1 &= \rho^{a_1} d^{a_2} v^{a_3} F \equiv [ML^{-3}]^{a_1} [L]^{a_2} [LT^{-1}]^{a_3} [MLT^{-2}] = [M^{a_1+1} L^{-3a_1+a_2+a_3+1} T^{-a_3-2}] \\ \Pi_2 &= \rho^{b_1} d^{b_2} v^{b_3} \eta \equiv [ML^{-3}]^{b_1} [L]^{b_2} [LT^{-1}]^{b_3} [ML^{-1}T^{-1}] = [M^{b_1+1} L^{-3b_1+b_2+b_3-1} T^{-b_3-1}] \end{aligned}$$

Hence we obtain two sets of equations:

$$\begin{aligned} a_1 + 1 &= 0 \implies a_1 = -1 \\ -a_3 - 2 &= 0 \implies a_3 = -2 \\ -3a_1 + a_2 + a_3 + 1 &= 0 \implies 3 + a_2 - 2 + 1 = 0 \implies a_2 = -2 \\ b_1 &= -1 \\ b_3 &= -1 \\ -3b_1 + b_2 + b_3 - 1 &= 0 \implies 3 + b_2 - 2 = 0 \implies b_2 = -1 \end{aligned}$$

Then we obtain the Π groups as:

$$\Pi_1 = \frac{F}{\rho d^2 v^2} \quad \Pi_2 = \frac{\eta}{\rho d v} = \frac{1}{\frac{\rho d v}{\eta}}$$

We identify $\frac{\rho d v}{\eta}$ to be the *Reynold's numer* $\mathcal{R}$. Then we can say, for some ϕ ,

$$\frac{F}{\rho d^2 v^2} = \phi(\mathcal{R})$$

To some extraterrestrial being, the physical laws will (hopefully) be valid, however, they might not understand the human-made units (like metres and seconds) in which we measure these quantities.

We need something natural, based on Nature and hence we used the natural units. For this purpose, we will use things like $c, \hbar, G, k_b \dots$ and we set all of them to 1. Doing this will lead to change in all scales. For that, we define the Planck units, made of fundamental constants of Nature:

- **Planck mass:** $E_p = \sqrt{\frac{\hbar c}{G}}$
- **Planck length:** $l_p = \sqrt{\frac{\hbar G}{c^3}}$
- **Planck time:** $t_p = \sqrt{\frac{\hbar c}{G^5}}$

Let us now analyse the physical regimes in which the fundamental constants become important. For that, we will use a tuple $(G, \frac{1}{c}, \hbar)$ (note that all the elements of the tuple are written in terms of very small quantities of the SI scale).

$(0, 0, 0) \rightarrow$	Classical mechanics
$(G, 0, 0) \rightarrow$	Newtonian gravity
$(0, \frac{1}{c}, 0) \rightarrow$	Special relativity
$(0, 0, \hbar) \rightarrow$	Basic quantum mechanics
$(G, \frac{1}{c}, 0) \rightarrow$	General relativity
$(0, \frac{1}{c}, \hbar) \rightarrow$	QFT and relativistic QM
$(G, 0, \hbar) \rightarrow$	Non-relativistic gravity
$(G, \frac{1}{c}, \hbar) \rightarrow$	Quantum gravity


Lecture 03: From Classical to Quantum

Inspecting the transition from a classical to quantum description is necessary for understanding QFT, which is just a *sophisticated mechanics*.

We know that the position $\mathbf{q} \equiv \{q_1, \dots, q_n\}$ and momentum $\mathbf{p} \equiv \{p_1, \dots, p_n\}$ uniquely define the state of a *classical* particle.

$$(\mathbf{q}, \mathbf{p}) \in \Pi \text{ (} 2n \text{ dimensional phase space)} \quad q_i, p_i \in \mathbb{R}$$

The Hamiltonian $H(\mathbf{q}, \mathbf{p})$ is just a way of performing time evolution on the system. Using the Hamiltonian, we can ‘fibrate’ the phase space, that is, starting from one point, we can go to some other point. Also, note that, changing the parameters in the Hamiltonian (eg. ω in SHO Hamiltonian), we change the fibration patterns(eg. *Lissajou’s* figures).



In the quantum world, the phase space changes to the Hilbert space $\mathcal{H}$ while a point in the phase space (q, p) changes to a vector $|\psi\rangle$ in the Hilbert space.

Also, these real variables become Hermitian (self-adjoint) operators ¹and the classical Poisson bracket now transforms to the commutator.

$$\begin{aligned} \text{CM:} \quad & \{q, p\} = 1 \\ \text{QM:} \quad & [\hat{q}, \hat{p}] = i \end{aligned}$$

3.1 Time Evolution

From classical Hamilton’s equations of motion, we have:

$$\begin{aligned} \dot{q}_i &= \{q_i, H\} = \frac{\partial H}{\partial p_i} \\ \dot{p}_i &= \{p_i, H\} = -\frac{\partial H}{\partial q_i} \end{aligned}$$

¹A self-adjoint operator $\mathcal{O}$ is such that $\mathcal{O} = \mathcal{O}^\dagger$ in all respect, that is, $\mathcal{O}$ and $\mathcal{O}^\dagger$ have the same domain and action. In general, $\mathbb{D}(\mathcal{O}) \subseteq \mathbb{D}(\mathcal{O}^\dagger)$ where $\mathbb{D}(\cdot)$ represents the domain of some operator. If the domains are not equal but action is same on a restricted domain (mainly occurs in infinite-dimensional spaces), then those operators are not self-adjoint but called *symmetric/Hermitian* (in some places Hermiticity also requires a symmetric operator to be bounded).

If $\mathbf{z} = \begin{pmatrix} \mathbf{p} \\ \mathbf{q} \end{pmatrix}$ is a $2n$ dimensional vector, then $\dot{\mathbf{z}} = \{\mathbf{z}, H\}$. Now, moving to quantum mechanics, we know the celebrated Schrödinger equation, which specifies the time evolution of a state vector $|\psi(t)\rangle \in \mathcal{H}$:

$$i \frac{\partial |\psi(t)\rangle}{\partial t} = H |\psi(t)\rangle$$

Thus, in both classical and quantum case, the time derivative of a quantity is equal to some action of the Hamiltonian on that quantity (classical $\rightarrow$ Poisson bracket, quantum $\rightarrow$ Multiplication with Hamiltonian)¹.

The Hamiltonian in the position basis becomes:

$$H = \frac{-\hbar^2}{2m} \nabla^2 + V(\mathbf{x})$$

Thus, the Schrödinger equation now acts on the wavefunction $\psi(\mathbf{x}, t) = \langle \mathbf{x} | \psi \rangle$:

$$i \frac{\partial \psi(\mathbf{x}, t)}{\partial t} = -\frac{1}{2m} \nabla^2 \psi(\mathbf{x}, t) + V(\mathbf{x}) \psi(\mathbf{x}, t)$$

And the energy and momentum operators become $E \rightarrow i \frac{\partial}{\partial t}$ and $p \rightarrow -i \nabla$.

3.1.1 Free Particle

The dispersion relation for a non-relativistic free particle is:

$$E = \frac{\mathbf{p}^2}{2m}$$

The typical solution can be written as plane-waves, $e^{i(\mathbf{k} \cdot \mathbf{x} - \omega t)} = e^{i(\mathbf{p} \cdot \mathbf{x} - Et)} \equiv \exp(-ip^\mu x_\mu)$ where we have used the repeated summation notation and taken the metric $\eta_{\mu\nu} = (+, -, -, -)$.

For a relativistic free particle, the dispersion relation becomes:

$$E^2 = m^2 + p^2$$

Instead of the wavefunction $\psi(\mathbf{x}, t)$, we will use $\phi(\mathbf{x}, t)$, since the resulting equation actually acts on a scalar-field and ϕ is common notation for a scalar-field. Substituting the energy and momentum operators here, we get:

$$\begin{aligned} i^2 \frac{\partial^2 \phi(\mathbf{x}, t)}{\partial t^2} &= (-i)^2 \nabla^2 \phi(\mathbf{x}, t) + m^2 \phi(\mathbf{x}, t) \\ \Rightarrow \frac{\partial^2 \phi(\mathbf{x}, t)}{\partial t^2} - \nabla^2 \phi(\mathbf{x}, t) &= -m^2 \phi(\mathbf{x}, t) \\ \Rightarrow \left(\frac{\partial^2}{\partial t^2} - \nabla^2 + m^2 \right) \phi(\mathbf{x}, t) &= 0 \end{aligned}$$

The above equation is called the *Klein-Gordon equation*. In covariant notation, we can compactly write it as:

$$(\partial_\mu \partial^\mu + m^2) \phi(x^\nu) = 0$$

The KG equation is Lorentz invariant, that is, under a Lorentz transformation, $x^\mu \rightarrow x'^\mu$, $\phi'(x'^\mu) = \phi(x^\mu)$

¹ A better analogy would be to use density matrices which gives the von Neumann equation which uses commutator bracket

3.1.2 Continuity Equation

Taking the complex conjugate of the Schrödinger's equation and then after some algebraic manipulation, we obtain:

$$\frac{\partial(\psi^*\psi)}{\partial t} = \frac{i}{2m} \nabla \cdot [\psi^*(\nabla\psi) - \psi(\nabla\psi^*)]$$

Here, we identify:

$$\rho := \psi^*\psi \quad \mathbf{J} = \frac{-i}{2m} [\psi^*(\nabla\psi) - \psi(\nabla\psi^*)]$$

which yields the well-known form of the *continuity equation*:

$$\frac{\partial\rho}{\partial t} + \nabla \cdot \mathbf{J} = 0 \quad \longrightarrow \quad \partial_\mu J^\mu = 0$$

where $J^\mu = (\rho, \mathbf{J})$ is the four-current. Note that in this case, ρ is a positive-definite quantity and can indeed have the interpretation of probability. Also, note that if ψ somehow becomes real-valued, then $\mathbf{J} = 0$ which implies that ρ is constant in time (though it can change in space).

Doing the same thing to Klein-Gordon equation yields:

$$\frac{\partial}{\partial t} \left(\phi^* \frac{\partial\phi}{\partial t} - \phi \frac{\partial\phi^*}{\partial t} \right) - \nabla \cdot (\phi^* \nabla\phi - \phi \nabla\phi^*) = 0$$

From this equation, we can identify:

$$\rho := \left(\phi^* \frac{\partial\phi}{\partial t} - \phi \frac{\partial\phi^*}{\partial t} \right) \quad \mathbf{J} := -(\phi^* \nabla\phi - \phi \nabla\phi^*)$$

Thus here, the 4-current becomes $J^\mu = (\phi^* \partial^\mu \phi - \phi \partial^\mu \phi^*)$. Note the apparent problems with this identification:

- It is not apriori obvious that ρ is positive-definite and hence has problem with probability interpretation.
- The equation is second-order in time and hence ρ seems to have a term evolving forward and one term evolving backward in time.
- The dispersion relation does not have a single solution; the solutions are $\pm E$

The thing is, KG equation treats time and space on equal footing (unlike Schrödinger equation where time was in first order and space was in second order). Hence we have to consider all possibilities of moving back and forth in both space and time.

Lecture 04: Classical Fields

Let us start with a familiar setting, the spring-mass system in one-dimensions. So, each mass is connected with isotropic springs of natural length a and spring constant k , and at some instance, i^{th} spring is displaced from its equilibrium position by an amount η_i as shown below:




Figure 1: A spring-mass system; each mass is displaced by some amount from its equilibrium position

Then the kinetic and potential energies can be written as:

$$T = \frac{1}{2} \sum_i \dot{\eta}_i^2 \quad V = \frac{1}{2} \sum_i k(\eta_{i+1} - \eta_i)^2$$

From this, the Langrangian of the system is obtained as:

$$L = T - V = \frac{1}{2} \sum_i a \left[(m/a) \dot{\eta}_i^2 - ka \left(\frac{\eta_{i+1} - \eta_i}{a} \right)^2 \right] = a \sum_i L_i$$

The resulting ELEOM are:

$$\frac{d}{dt} \left(\frac{\partial L}{\partial \dot{\eta}_i} \right) = \frac{\partial L}{\partial \eta_i} \implies (m/a) \ddot{\eta}_i - ka \left(\frac{\eta_{i+1} - \eta_i}{a^2} \right) + ka \left(\frac{\eta_i - \eta_{i-1}}{a^2} \right) = 0$$

In the *continuum* limit, that is, $a \rightarrow 0$ (so, the system kind of becomes like a rod), we have:

$$\begin{aligned} \frac{m}{a} &\longrightarrow \mu && : \text{mass per unit length of the rod} \\ ka &\longrightarrow Y && : \text{Young's modulus of the rod} \\ i &\longrightarrow x && : \text{position label} \\ \eta_i &\longrightarrow \eta(x) && : \text{field} \\ \frac{\eta_{i+1} - \eta_i}{a} &\longrightarrow \frac{d\eta}{dx} && : \text{derivative} \\ \sum_i &\longrightarrow \int dx && : \text{integral} \end{aligned}$$

Hence in the continuum limit, the ELEOM becomes:

$$\lim_{a \rightarrow 0} \left(\mu \frac{d^2 \eta}{dt^2} - \frac{Y}{a} \frac{d\eta}{dx} \Big|_x + \frac{Y}{a} \frac{d\eta}{dx} \Big|_{x-a} \right) = 0 \implies \mu \frac{d^2 \eta}{dt^2} - Y \frac{d^2 \eta}{dx^2} = 0$$

Defining the velocity as $v := \sqrt{\frac{Y}{\mu}}$, we obtain the wave-equation:

$$\frac{1}{v^2} \frac{d^2 \eta}{dt^2} - \frac{d^2 \eta}{dx^2} = 0$$

Note that, in this case, position becomes a mere label like in the discrete case, i just labelled the index of the i^{th} mass. Also, technically, η can also depend on time, so it is better to write it as $\eta(x, t)$.

Position and time, both are kind of labels and thus are independent. Hence, total and partial derivatives coincide in this case. Note that in continuum, the Langrangian itself becomes an integral of the form:

$$L = \frac{1}{2} \int \left[\mu \dot{\eta}^2 - Y \left(\frac{d\eta}{dx} \right)^2 \right] dx$$

The bracketted term is the Lagrangian density, denoted by $\mathcal{L}$ and hence, the Lagrangian is the space integral of the Langrangian density:

$$L = \int \mathcal{L} dx$$

Henceforth, we allow ourselves to be *sloppy* enough and refer to Lagrangian density as simply the Langrangian when context is clear.

4.1 Our actions tell a lot!

Let us now consider a general setting. So, the Lagrangian density can be a function of the field, the spatial and temporal derivatives and space and time themselves. In compact notation, we write:

$$\mathcal{L} \equiv \mathcal{L}(\phi, \partial_\mu \phi, x^\mu)$$

Remember, that we define the classical action S as:

$$S = \int_{t_1}^{t_2} dt L = \int_{t_1}^{t_2} dt \int d^3\mathbf{x} \mathcal{L}(\phi, \partial_\mu \phi, x^\mu) = \int_{\mathcal{R} \subset \mathbb{R}^4} d^4\mathbf{x} \mathcal{L}(\phi, \partial_\mu \phi, x^\mu)$$

The integration can indeed be over a region $\mathcal{R}$ which is a subset of $\mathbb{R}^4$ in general. Mostly, we take $\mathcal{R} = [t_1, t_2] \times \mathbb{R}^3$.

We now want to use Hamilton's principle, which states that, under fixed boundary conditions, the system takes a path in configuration space for which the action is stationary, that is,

$$\delta S = 0$$

Note that the action is a functional which takes a field ϕ as input and returns a real number¹. Let us consider a variation in the field itself: $\phi \rightarrow \phi' = \phi + \delta\phi$ such that $\delta\phi$ vanishes at the boundary of the integration domain. Then we have the following:

$$\delta S = S[\phi'] - S[\phi] = \int d^4\mathbf{x} (\mathcal{L}(\phi', \partial_\mu \phi', x^\mu) - \mathcal{L}(\phi, \partial_\mu \phi, x^\mu))$$

Now, we try to find what the above quantity is:

$$\begin{aligned} \mathcal{L}(\phi', \partial_\mu \phi', x^\mu) &= \mathcal{L}(\phi + \delta\phi, \partial_\mu(\phi + \delta\phi), x^\mu) \\ &= \mathcal{L}(\phi, \partial_\mu \phi, x^\mu) + \frac{\partial \mathcal{L}}{\partial \phi} \delta\phi + \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \delta(\partial_\mu \phi) \end{aligned}$$

Also note that $\delta(\partial_\mu \phi) = \partial_\mu(\delta\phi)$. Hence, from the above expression, we obtain that:

$$\delta S = \int d^4\mathbf{x} \left(\frac{\partial \mathcal{L}}{\partial \phi} \delta\phi + \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial_\mu(\delta\phi) \right)$$

Using *integration by parts* on the second term, we obtain:

$$\int d^4\mathbf{x} \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial_\mu(\delta\phi) = - \int d^4\mathbf{x} \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \right) \delta\phi + \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \delta\phi \right) \Big|_{\text{boundary}}$$

The variation in the field vanishes at the boundary by our assumption. Then we obtain, for the variation of the action:

$$\delta S = \int d^4\mathbf{x} \left(\frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \right) \right) \delta\phi = 0$$

Since the variation in the field was arbitrary, we obtain the Euler-Lagrange's equations for the field as:

$$\frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \right) = 0$$

In general, we want the action to be *Lorentz* invariant, in order to maintain the universality of physical laws. Since the volume element of Minkowski space $d^4\mathbf{x}$ is invariant under Lorentz transformations, this imply that $\mathcal{L}$ should also be constructed of Lorentz-invariant quantities.

¹For functionals, a common notation for the input is to use square brackets, that is, if A is a linear functional taking function f as input, $A[f] \in \mathbb{R}$

The field ϕ (and all its powers) is a Lorentz *scalar* and is hence invariant. The derivative $\partial_\mu \phi$ is not invariant, however, $\partial^\mu \partial_\mu \phi$ is. Bilinears like $(\partial^\mu \phi)(\partial_\mu \phi)$ are also invariant. Hence, a simple Lagrangian should be constructed of these quantities. Let us consider the Lagrangian:

$$\mathcal{L} = \frac{1}{2}(\partial^\mu \phi)(\partial_\mu \phi) - \frac{1}{2}m^2 \phi^2 = \frac{1}{2}\eta^{\mu\nu}(\partial_\nu \phi)(\partial_\mu \phi) - \frac{1}{2}m^2 \phi^2$$

Then, from the Lagrange's equations,

$$\begin{aligned}\frac{\partial \mathcal{L}}{\partial \phi} &= -m^2 \phi \\ \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} &= \frac{1}{2}\eta^{\mu\nu}(\partial_\nu \phi + \partial_\mu \phi \delta^\mu_\nu) \\ &= \frac{1}{2}\eta^{\mu\nu}(2 \times \partial_\nu \phi) \\ &= \eta^{\mu\nu} \partial_\nu \phi \\ &= \partial^\mu \phi\end{aligned}$$

Then we finally obtain:

$$-m^2 \phi - \partial_\mu \partial^\mu \phi = 0 \implies (\partial_\mu \partial^\mu + m^2)\phi = 0$$

This is the KG equation that we had seen earlier. Thus, the above Lagrangian models the KG equation.

When we have multiple fields $\{\phi_i\}$, then for each field, we will obtain:

$$\frac{\partial \mathcal{L}}{\partial \phi_i} - \partial_\mu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) = 0$$

NOTE: In terms of the total Lagrangian, the ELEOM for the field is:

$$\frac{\delta L}{\delta \phi} - \partial_t \left(\frac{\delta L}{\delta \dot{\phi}} \right) = 0$$

Lecture 05 & 06: Looking into Hamiltonians

Previously, we saw a ton of crap about the Lagrangians and how the equations of motions are obtained from the action being stationary. Now, we will look into the *Hamiltonian* formalism which will allow us to quantise the field.

In basic classical mechanics, we had seen that given a Lagrangian $L(q_i, \dot{q}_i, t)$, we obtained the Hamiltonian by the Legendre transformation as:

$$H(q_i, p_i, t) = p_i \dot{q}_i - L(q_i, \dot{q}_i, t)$$

where $p_i := \frac{\partial L}{\partial \dot{q}_i}$ was defined to be the generalised momenta, conjugate to the generalised coordinate q_i . Similarly, for classical fields, we can define:

$$\pi^i(x^\mu) := \frac{\partial \mathcal{L}}{\partial(\partial_0 \phi_i)} = \frac{\partial \mathcal{L}}{\partial \dot{\phi}_i}$$

This is the *canonical momentum* corresponding to the field. From the ELEOM, we obtain $\dot{\pi}^i = \frac{\delta L}{\delta \phi}$ Using this, the *Hamiltonian density* can be defined as:

$$\mathcal{H}(\phi^a, \pi_a, \partial_i \phi_a, x^\mu) := \pi^a \dot{\phi}_a - \mathcal{L}(\phi_a, \partial_\mu \phi_a, x^\mu)$$

In the final expression on the LHS, we have to replace $\dot{\phi}_a$ in terms of π^a to get the Hamiltonian density (Replacing this will get rid of all the time derivative and hence, in the Hamiltonian, we used ∂_i , the spatial derivatives only). Using this then, we get the total Hamiltonian as:

$$H[\pi^a, \phi_a] = \int d^3 \mathbf{x} \mathcal{H}(\phi_a, \pi_a, \partial_i \phi_a, x^\mu)$$

6.1 Functional Derivatives

Since action, Lagrangian, etc. are functionals, an apt way to describe their variation is through the *functional derivative*. These come under the domain of functional analysis (which is Math 🧐), however, a short introduction might be necessary.

Let us take a functional $\mathcal{F}[f]$ and consider its variation $\delta\mathcal{F}$ when $f \rightarrow f + \delta f$, where $\delta f = \epsilon\eta$. Here ϵ is a small parameter and η is a generic function, often called the *test function*. Now,

$$\mathcal{F}[f + \epsilon\eta] = \mathcal{F}[f] + \left. \frac{d\mathcal{F}[f + \epsilon\eta]}{d\epsilon} \right|_{\epsilon=0} \epsilon + \mathcal{O}(\epsilon^2)$$

We define the functional derivative as the first-order coefficient of the expansion, that is,

$$\frac{\delta\mathcal{F}}{\delta f} = \lim_{\epsilon \rightarrow 0} \frac{\mathcal{F}[f + \epsilon\eta] - \mathcal{F}[f]}{\epsilon}$$

Functional derivatives seem to follow similar rules as in ordinary differential calculus (viz. linearity, Liebnitz rule, chain rule) and hence, can be treated to be operationally same as the ordinary derivative. However, exceptions are always there.

The differential (variation) of a functional is given as:

$$\delta\mathcal{F}[\phi] := \int \frac{\delta\mathcal{F}[\phi]}{\delta\phi} \delta\phi$$

Now let us try to relate functional derivative with normal derivative. For that, let us suppose that the space is discretised into cells, each of volume ΔV^i and each cell is assigned the *average value* of ϕ within the volume, that is,

$$\phi_i(t) = \frac{1}{\Delta V^i} \int_{\Delta V^i} d^3\mathbf{x} \phi(\mathbf{x}, t)$$

Then, L now depends on the discrete components, ϕ_i and $\dot{\phi}_i$ and the variation can be written as in terms of partial derivatives (since ϕ_i are now discrete):

$$\delta L = \sum_i \left(\frac{\partial L}{\partial \phi_i} \delta\phi_i + \frac{\partial L}{\partial \dot{\phi}_i} \delta\dot{\phi}_i \right) = \sum_i \frac{1}{\Delta V^i} \left(\frac{\partial L}{\partial \phi_i} \delta\phi_i + \frac{\partial L}{\partial \dot{\phi}_i} \delta\dot{\phi}_i \right) \Delta V^i$$

Now, consider the Lagrangian variation using the definition of functional derivative:

$$\delta L = \int d^3\mathbf{x} \left(\frac{\delta L}{\delta\phi} \delta\phi + \frac{\delta L}{\delta\dot{\phi}} \delta\dot{\phi} \right)$$

From the two above equations, we can make the following identifications:

$$\begin{aligned} \frac{\delta L}{\delta\phi} &= \lim_{\Delta V^i \rightarrow 0} \frac{1}{\Delta V^i} \frac{\partial L}{\partial \phi_i} \\ \frac{\delta L}{\delta\dot{\phi}} &= \lim_{\Delta V^i \rightarrow 0} \frac{1}{\Delta V^i} \frac{\partial L}{\partial \dot{\phi}_i} \end{aligned}$$

In the LHS, while taking functional derivative, $\mathbf{x}$ belongs to the i^{th} cell. Thus, we see that the functional derivative is proportional to the partial derivative. Now, given a Lagrangian, we can write the variation under $\phi \rightarrow \phi + \delta\phi$ using the Lagrangian density:

$$\delta L = \int d^3\mathbf{x} \left(\frac{\partial \mathcal{L}}{\partial \phi} \delta\phi + \frac{\partial \mathcal{L}}{\partial(\partial_i \phi)} \delta(\partial_i \phi) + \frac{\partial \mathcal{L}}{\partial \dot{\phi}} \delta\dot{\phi} \right)$$

Integration by parts and ignoring boundaries on the second term, we obtain:

$$\delta L = \int d^3\mathbf{x} \left(\left(\frac{\partial \mathcal{L}}{\partial \phi} - \partial_i \left(\frac{\partial \mathcal{L}}{\partial(\partial_i \phi)} \right) \right) \delta\phi + \frac{\partial \mathcal{L}}{\partial \dot{\phi}} \delta\dot{\phi} \right)$$

Comparing this with the definition of functional derivative, we obtain:

$$\frac{\delta L}{\delta\phi} = \frac{\partial \mathcal{L}}{\partial \phi} - \partial_i \left(\frac{\partial \mathcal{L}}{\partial(\partial_i \phi)} \right) \quad \frac{\delta L}{\delta\dot{\phi}} = \frac{\partial \mathcal{L}}{\partial \dot{\phi}}$$

6.2 Hamilton's Equations of Motion

To obtain HEOM, we start with the variation of the total Hamiltonian,

$$\delta H = \int d^3\mathbf{x} \delta(\pi^\alpha \dot{\phi}_\alpha - \mathcal{L}) = \int d^3\mathbf{x} (\pi^\alpha \delta \dot{\phi}_\alpha + \dot{\phi}_\alpha \delta \pi^\alpha) - \delta L$$

Now, from ELEOM:

$$\delta L = \int \frac{\delta L}{\delta \phi_\alpha} \delta \phi_\alpha + \frac{\delta L}{\delta \dot{\phi}_\alpha} \delta \dot{\phi}_\alpha = \int \partial_t \left(\frac{\delta L}{\delta \dot{\phi}_\alpha} \right) \delta \phi_\alpha + \pi^\alpha \delta \dot{\phi}_\alpha = \int \dot{\pi}^\alpha \delta \phi_\alpha + \pi^\alpha \delta \dot{\phi}_\alpha$$

Using this, the variation in Hamiltonian becomes:

$$\delta H = \int d^3\mathbf{x} (\dot{\phi}_\alpha \delta \pi^\alpha - \dot{\pi}^\alpha \delta \phi_\alpha)$$

Thus, from above, we can identify:

$$\frac{\delta H}{\delta \phi_\alpha} = -\dot{\pi}^\alpha \quad \frac{\delta H}{\delta \pi^\alpha} = \dot{\phi}_\alpha$$

Also, previously we saw the relation between functional derivative of the Lagrangian and partial derivative of the Lagrangian density. That could well be extended to the Hamiltonian and we obtain:

$$\dot{\pi}^\alpha = -\frac{\delta H}{\delta \phi_\alpha} = -\frac{\partial \mathcal{H}}{\partial \phi_\alpha} + \partial_i \left(\frac{\partial \mathcal{H}}{\partial (\partial_i \phi_\alpha)} \right) \quad \dot{\phi}_\alpha = \frac{\delta H}{\delta \pi^\alpha} = \frac{\partial \mathcal{H}}{\partial \pi^\alpha} - \partial_i \left(\frac{\partial \mathcal{H}}{\partial (\partial_i \pi^\alpha)} \right)$$

In general, the Hamiltonian density does not depend on $\partial_i \pi^\alpha$, so we have $\dot{\phi}_\alpha = \frac{\partial \mathcal{H}}{\partial \pi^\alpha}$

6.3 Poisson Bracket

In the Hamiltonian formulation of classical mechanics, we defined the Poisson bracket of two quantities A, B by:

$$\{A, B\} = \frac{\partial A}{\partial q_i} \frac{\partial B}{\partial p_i} - \frac{\partial A}{\partial p_i} \frac{\partial B}{\partial q_i}$$

Similarly, for fields also, given two functionals $\mathcal{F}[\phi, \pi]$ and $\mathcal{G}[\phi, \pi]$, we define the Poisson bracket by:

$$\{\mathcal{F}, \mathcal{G}\} := \int d^3\mathbf{x} \frac{\partial \mathcal{F}}{\partial \phi_\alpha} \frac{\partial \mathcal{G}}{\partial \pi^\alpha} - \frac{\partial \mathcal{F}}{\partial \pi^\alpha} \frac{\partial \mathcal{G}}{\partial \phi_\alpha}$$

Given a function $\mathcal{F}[\phi^\alpha, \pi^\alpha, t]$, note:

$$\begin{aligned} \dot{\mathcal{F}} &= \partial_t \mathcal{F} + \int d^3\mathbf{x} \left(\frac{\delta \mathcal{F}}{\delta \phi_\alpha} \frac{\partial \phi_\alpha}{\partial t} + \frac{\delta \mathcal{F}}{\delta \pi_\alpha} \frac{\partial \pi_\alpha}{\partial t} \right) \\ &= \partial_t \mathcal{F} + \int d^3\mathbf{x} \left(\frac{\delta \mathcal{F}}{\delta \phi_\alpha} \frac{\delta H}{\delta \pi^\alpha} - \frac{\delta \mathcal{F}}{\delta \pi_\alpha} \frac{\delta H}{\delta \phi_\alpha} \right) \quad (\text{from HEOM}) \\ &= \partial_t \mathcal{F} + \{\mathcal{F}, H\} \end{aligned}$$

This specifies the *time-evolution* of the functionals using the Poisson bracket. Now, let us consider the Poisson bracket of the fields ϕ_α and π^α . However, since the Poisson bracket was defined for functionals only, there is some problem. This can be resolved by noting that any function can be written as a functional depending on itself as:

$$\phi_\alpha(\mathbf{x}, t) = \int d^3\mathbf{x}' \mathbf{x}' \phi_\alpha(\mathbf{x}', t) \delta^3(\mathbf{x} - \mathbf{x}')$$

From the definition of the functional derivative, it is easy to note that:

$$\frac{\delta \phi_\alpha(\mathbf{x}, t)}{\delta \phi_\beta(\mathbf{x}', t)} = \delta_{\alpha\beta} \delta^3(\mathbf{x} - \mathbf{x}')$$

Also, ϕ and π are *independent functionals* and hence,

$$\frac{\delta\pi(\mathbf{x}, t)}{\delta\phi(\mathbf{x}', t)} = 0 \quad \frac{\delta\phi(\mathbf{x}, t)}{\delta\pi(\mathbf{x}', t)} = 0 \quad \forall \mathbf{x}, \mathbf{x}', t$$

Now, we can compute the Poisson bracket of the fields themselves and we obtain:

$$\begin{aligned} \{\phi_\alpha(\mathbf{x}, t), \pi^\beta(\mathbf{x}', t)\} &= \int d^3 \left(\mathbf{x}'' \frac{\delta\phi_\alpha(\mathbf{x}, t)}{\delta\phi_\mu(\mathbf{x}'', t)} \frac{\delta\pi^\beta(\mathbf{x}', t)}{\delta\pi^\mu(\mathbf{x}'', t)} - 0 \right) \\ &= \int d^3 (\mathbf{x}'' \delta_{\alpha\mu} \delta_{\beta\mu} \delta^3(\mathbf{x} - \mathbf{x}'') \delta^3(\mathbf{x}' - \mathbf{x}'')) \\ &= \delta_{\alpha\beta} \delta^3(\mathbf{x} - \mathbf{x}') \end{aligned}$$

We also have the other two Poisson brackets which are trivial:

$$\{\phi_\alpha, \phi_\beta\} = 0 \quad \{\pi^\alpha, \pi^\beta\} = 0$$


Lecture 07 & 08: Representations of Rotation and Lorentz group

A *group* is a tuple $(G, *)$ which follows certain rules like associativity, existence of identity, inverses, etc. A *Lie Group* ¹ is some continuous group with properties so special that it becomes very important (and cool) to study them.

I think a short, vague introduction cannot do justice to these elegant topics which are subjects on their own right and require proper analysis. Hence, I might skip them here and instead try to include them in the link ² below as and when I read about them!

Lecture 09: Noether's Theorem

To understand Noether's Theorem, let us first understand what symmetry actually is. Symmetry of a system can be thought of as some transformation under which the properties of the system remain invariant.



Discrete symmetries describes non-continuous changes in the system, like *parity* (kind of spatial reflection) and *time-reversal*. *Internal* symmetries include cases where the fields themselves vary while *space-time* symmetry describes the system when the space-time points vary. *Local* symmetries are transformations allowing us to change the system differently at different spacetime points while *global* symmetries act in the same way at every point.

¹A Lie Group is a group that is also a manifold.

²A very basic and nasty introduction can be found [here!](#)

Theorem 2 (Noether's Theorem):

For every continuous global symmetry there exists a conserved current J^μ , that is,

$$\partial_\mu J^\mu = 0$$

Proof. Note that, under any symmetry transformation, we required our equations of motion to be invariant and hence require our action $\mathcal{S}$ to be 'invariant' since action is a fundamental quantity to our system from which we derived our EOM. Note that while deriving the equations of motions, we ignored the boundary term of the action. Thus, the action can vary by some arbitrary boundary term. Using this, we define a *Noether symmetry* to be a transformation which keeps the action invariant upto an integral of a total divergence, that is,

$$\delta\mathcal{S} = \int_{\partial\mathcal{R}} d^4\mathbf{x} \partial_\mu K^\mu \implies \delta\mathcal{L} \equiv \epsilon \partial_\mu K^\mu$$

The implication comes from the fact that divergence should be in first order with respect to some infinitesimal variation ϵ . Let us consider the case of *internal symmetries* first.

Consider a variation in the field $\phi = \phi_0 + \delta\phi$. Then,

$$\delta\mathcal{L} = \left. \frac{\partial\mathcal{L}}{\partial\phi} \right|_{\phi_0} \delta\phi + \left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\phi_0} \delta(\partial_\mu\phi) = \left. \frac{\partial\mathcal{L}}{\partial\phi} \right|_{\phi_0} \delta\phi + \left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\phi_0} \partial_\mu(\delta\phi)$$

Now, suppose there exists a solution $\tilde{\phi}$ to the equation of motion of the system. Since the above equation is valid for any ϕ_0 , it is also valid for $\tilde{\phi}$ and hence, we get:

$$\begin{aligned} \delta\mathcal{L} &= \left. \frac{\partial\mathcal{L}}{\partial\phi} \right|_{\tilde{\phi}} \delta\phi + \left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \partial_\mu(\delta\phi) \\ &= \partial_\mu \left(\left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \delta\phi + \left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \partial_\mu(\delta\phi) \right) \quad (\text{using ELEOM}) \\ &= \partial_\mu \left(\left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \delta\phi \right) \quad (\text{using product rule}) \end{aligned}$$

Let the variation of the Lagrangian be a global symmetry, denoted by, $\delta_\alpha\phi$. Under this symmetry, we want our action $\mathcal{S}$ to change atmost by a boundary term and hence we have,

$$\partial_\mu \left(\left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \delta\phi \right) = \partial_\mu K^\mu \implies \partial_\mu \left(\left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \delta\phi - K^\mu \right) = 0$$

Let us define the quantity in the bracket by J^μ which is called the *Noether current*. Thus,

$$J^\mu := \left. \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)} \right|_{\tilde{\phi}} \delta\phi - K^\mu$$

from where we get $\partial_\mu J^\mu = 0 \implies$ Noether current is conserved.

The above proof worked for internal symmetries where the field itself was varied. Now, let us look into the *spacetime* symmetry, where the spacetime points are varied. For simplicity, we will deal with scalar fields only. Let us consider an infinitesimal coordinate transformation,

$$x'^\mu = x^\mu + \delta x^\mu$$

The variation in the field can be given by:

$$\delta\phi = \phi'(\mathbf{x}') - \phi(\mathbf{x}) = \phi'(x'^\mu) - \phi'(x^\mu) + \phi'(x^\mu) - \phi(x^\mu) = \delta_0\phi(x^\mu) + \phi'(x'^\mu) - \phi'(x^\mu)$$

where we have defined $\delta_0\phi(x^\mu) := \phi'(x^\mu) - \phi(x^\mu)$. We did this to distinguish between variation of the dependent and independent quantities. Using Taylor expansion, we have:

$$\phi'(x'^\mu) = \phi'(x^\mu + \delta x^\mu) = \phi'(x^\mu) + \frac{\partial\phi'}{\partial x^\mu}\delta x^\mu \implies \phi'(x'^\mu) - \phi'(x^\mu) = \partial_\mu\phi'\delta x^\mu$$

Now, $\partial_\mu\phi'\delta x^\mu \approx \partial_\mu\phi\delta x^\mu$, since if we do the Taylor expansion of $\delta_0\phi(x'^\mu)$, we would obtain a term $\partial_\nu\delta_0$ which is a variation of second order and hence, can be neglected. Thus, we obtain:

$$\phi'(x'^\mu) - \phi'(x^\mu) = \partial_\mu\phi\delta x^\mu \implies \delta\phi = \delta_0\phi(x^\mu) + \partial_\mu\phi\delta x^\mu$$

The action is not in general invariant under both changes in coordinates and fields. Indeed, the volume element (integral measure) $d^4\mathbf{x}$ transforms as:

$$d^4\mathbf{x}' = J d^4\mathbf{x}$$

where J is the determinant of the Jacobian matrix. Now, we find some expression for the Jacobian:

$$x'^\mu = x^\mu + \delta x^\mu \implies \partial_\nu x'^\mu = \delta^\mu_\nu + \partial_\nu(\delta x^\mu)$$

Now, we use an identity $\det(\mathbb{1} + \epsilon A) = 1 + \epsilon \text{Tr } A + \mathcal{O}(\epsilon^2)$ on the above equation to get an approximation for the Jacobian as:

$$J = \det\left(\frac{\partial x'^\mu}{\partial x^\nu}\right) = \det(\delta^\mu_\nu + \partial_\nu(\delta x^\mu)) \approx (1 + \partial_\nu(\delta x^\mu)) \implies \delta J = \partial_\mu(\delta x^\mu)$$

Now, we consider the variation of the action itself and we see:

$$\begin{aligned} \delta\mathcal{S} &= \int \delta(d^4\mathbf{x}) \mathcal{L} + \int d^4\mathbf{x} \delta\mathcal{L} \\ &= \int d^4\mathbf{x} \partial_\mu\delta x^\mu \mathcal{L} + \int d^4\mathbf{x} \delta\mathcal{L} \\ &= \int d^4\mathbf{x} \left(\partial_\mu\delta x^\mu \mathcal{L} + \frac{\partial\mathcal{L}}{\partial\phi}\delta_0\phi + \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\delta_0(\partial_\mu\phi) + \frac{\partial\mathcal{L}}{\partial x^\mu}\delta x^\mu \right) \end{aligned}$$

We had written the variation of the Lagrangian with respect to both dependent and independent variables. Now the first and last term can be written together using product rule and hence we have:

$$\begin{aligned} \delta\mathcal{S} &= \int d^4\mathbf{x} \left(\partial_\mu(\delta x^\mu \mathcal{L}) + \frac{\partial\mathcal{L}}{\partial\phi}\delta_0\phi + \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\partial_\mu\delta_0\phi \right) \\ &= \int d^4\mathbf{x} \left(\partial_\mu(\delta x^\mu \mathcal{L}) + \frac{\partial\mathcal{L}}{\partial\phi}\delta_0\phi + \partial_\mu\left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\delta_0\phi\right) - \partial_\mu\left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\right)\delta_0\phi \right) \\ &= \int d^4\mathbf{x} \left(\partial_\mu\left(\delta x^\mu \mathcal{L} + \left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\delta_0\phi\right)\right) + \delta_0\phi\left(\frac{\partial\mathcal{L}}{\partial\phi} - \partial_\mu\left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\right)\right) \right) \end{aligned}$$

For our field satisfying the equation of motion, the last term drops and we have our variation of the action and this should be equal to the boundary term, hence we have:

$$\partial_\mu\left(\delta x^\mu \mathcal{L} + \left(\frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi)}\delta_0\phi\right)\right) = \partial_\mu K^\mu$$

In the case for multiple fields, the *Noether current* turns out to be:

$$J^\mu := \frac{\partial\mathcal{L}}{\partial(\partial_\mu\phi_i)}\delta_0\phi_i + \delta x^\mu \mathcal{L} - K^\mu \quad \partial_\mu J^\mu = 0 \quad \square$$

The above condition is called *current conservation* which we had also seen earlier for internal symmetries. However, the second term, which is purely an artifact of the spacetime symmetry, did not occur there in the conserved current for internal symmetries. Associated with each current, we can define the charge enclosed in a volume V as:

$$Q_V := \int_V d^3\mathbf{x} J^0(\mathbf{x}, t) = \int_V d^3\mathbf{x} \rho$$

Then, we can write the conservation equation as:

$$\frac{d\rho}{dt} + \nabla \cdot \mathbf{J} = 0 \implies \frac{dQ}{dt} = - \int_V d^3\mathbf{x} \nabla \cdot \mathbf{J} = - \int_{\partial V} \mathbf{J} \cdot d\mathbf{S}$$

The last equality comes from Gauss' law and ∂V is the boundary of the region. If we extend the volume V to cover the entire space and assume that at spatial infinities, current is zero, then we have $\frac{dQ}{dt} = 0$ and hence, current is conserved. In most cases, we would have K^μ to be zero (and thus, let us take it to be zero). Let us see some examples for Noether's theorem:

9.1 Examples: Spacetime Symmetries

Let us start with some examples of Noether's theorem where spacetime symmetry is explicitly there. We will consider translations and rotations.

9.1.1 Translational Invariance

Classically, we know that translational invariance leads to *energy conservation*. Let us see what happens in case of fields. We thus consider an infinitesimal global translation:

$$x^\mu \rightarrow x'^\mu = x^\mu + \epsilon a^\mu$$

Then $\delta x^\mu = \epsilon a^\mu$. As calculated before, the variation in the Lagrangian is given by:

$$\delta \mathcal{L} = \partial_\mu \left(\delta x^\mu \mathcal{L} + \delta_0 \phi_i \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \right)$$

Note that $\delta x^\mu = \epsilon a^\mu$ and $\delta_0 \phi_i = \phi'_i(x^\mu) - \phi_i(x^\mu)$. Note that the new field at x^μ is just the previous field at $x^\mu - \epsilon a^\mu$ and hence we have from a first-order expansion that $\delta_0 \phi_i = -\epsilon a^\nu \partial_\nu \phi_i$. Substituting this above, we obtain:

$$\begin{aligned} \partial_\mu \left(\delta x^\mu \mathcal{L} + \delta_0 \phi_i \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \right) &= \partial_\mu \left(\epsilon a^\mu \mathcal{L} - \epsilon a^\nu \partial_\nu \phi_i \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \right) \\ &= \partial_\mu \left(\epsilon a^\nu \delta^\mu_\nu \mathcal{L} - \epsilon a^\nu \partial_\nu \phi_i \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \right) \\ &= -\epsilon a^\nu \partial_\mu \left(\partial_\nu \phi_i \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} - \delta^\mu_\nu \mathcal{L} \right) \end{aligned}$$

From the above, we can thus define the conserved current as:

$$T^\mu_\nu := \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \partial_\nu \phi_i - \delta^\mu_\nu \mathcal{L}$$

Now, note the following:

$$T^{\mu\nu} = g^{\alpha\nu} T^\mu_\alpha = \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} g^{\alpha\nu} \partial_\alpha \phi_i - g^{\alpha\nu} \delta^\mu_\alpha \mathcal{L} = \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi_i)} \partial^\nu \phi_i - g^{\mu\nu} \mathcal{L}$$

The above quantity can be shown to be a tensor and is called the *energy-momentum stress tensor*. Why?

Note that for each value of ν we have a current four vector. Let us calculate the charge for the time-translation invariance part, that is, $\nu = 0$. From this, we have:

$$P^0 = \int d^3\mathbf{x} T^{00} = \int d^3\mathbf{x} \left(\frac{\partial \mathcal{L}}{\partial(\partial_0 \phi_i)} \partial^0 \phi_i - g^{00} \mathcal{L} \right) = \int d^3\mathbf{x} (\pi^i \dot{\phi}_i - \mathcal{L}) = \int d^3\mathbf{x} \mathcal{H} = H$$

We see that corresponding to the conserved current from time translation invariance, we have a conserved charge $P^0 = H$. Thus, the total Hamiltonian is conserved under time translation and hence our classical intuition worked here as well!

Now, let us see what happens for space translations, for which we consider $\nu \equiv j = 1, 2, 3$. Then we have:

$$P^j = \int d^3\mathbf{x} T^{0j} = \int d^3\mathbf{x} \left(\frac{\partial \mathcal{L}}{\partial(\partial_0 \phi_i)} \partial^j \phi_i - g^{0j} \mathcal{L} \right) = \int d^3\mathbf{x} (\pi^i \partial^j \phi_i)$$

The last thing is the j^{th} component of the total momentum which is conserved in this case. Thus, we see that for spatial translation invariance, the total momentum is conserved, which once again matches with the classical analog.

9.1.2 Lorentz Invariance

An infinitesimal Lorentz transformation (which covers both *boosts* and *rotations*) can be written as:

$$\Lambda^\mu{}_\nu = \delta^\mu{}_\nu + \omega^\mu{}_\nu$$

where $\omega^\mu{}_\nu$ is an infinitesimal deviation from the identity. Now, we know for any Lorentz transformation,

$$\Lambda^\mu{}_\tau \eta^{\tau\sigma} \Lambda^\nu{}_\sigma = \eta^{\mu\nu} \iff \Lambda^T \eta \Lambda = \eta$$

If we put the infinitesimal transformation in the above equation and retain only terms upto first order in ω , we would have $\omega^{\mu\nu} = \omega^{\nu\mu}$, that is, the infinitesimal deviation is an anti-symmetric matrix. The spacetime coordinates transforms as:

$$x^\mu \rightarrow x'^\mu = \Lambda^\mu{}_\nu x^\nu = \delta^\mu{}_\nu x^\nu + \omega^\mu{}_\nu x^\nu = x^\mu + \omega^\mu{}_\nu x^\nu \implies \delta x^\mu = \omega^\mu{}_\nu x^\nu$$

And the variation in ϕ is given as:

$$\delta_0 \phi_i = \phi'(x^\mu) - \phi(x^\mu) = \phi(x^\mu + \omega^\mu{}_\nu x^\nu) - \phi(x^\mu) = -\omega^\mu{}_\nu x^\nu \partial_\mu \phi_i$$

Now, we are ready to put in the variation of the Lagrangian:

$$\begin{aligned} \partial_\mu \left(\delta x^\mu \mathcal{L} + \delta_0 \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) &= \partial_\mu \left(\omega^\mu{}_\nu x^\nu \mathcal{L} - \omega^\sigma{}_\nu x^\nu \partial_\sigma \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) \\ &= \partial_\mu \left(\eta^{\mu\rho} \omega_{\rho\nu} x^\nu \mathcal{L} - \eta^{\sigma\lambda} \omega_{\lambda\nu} x^\nu \partial_\sigma \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) \\ &= \partial_\mu \left(\eta^{\mu\rho} \omega_{\rho\nu} x^\nu \mathcal{L} - \omega_{\lambda\nu} x^\nu \partial^\lambda \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) \\ &= \partial_\mu \left(\eta^{\mu\rho} \omega_{\rho\nu} x^\nu \mathcal{L} - \omega_{\rho\nu} x^\nu \partial^\rho \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) \\ &= \partial_\mu \left(-\omega_{\rho\nu} x^\nu \times \left(-\eta^{\mu\rho} \mathcal{L} + \partial^\rho \phi_i \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi_i)} \right) \right) \\ &= \partial_\mu (-\omega_{\rho\nu} x^\nu T^{\rho\mu}) \\ &= \partial_\mu (\omega_{\nu\rho} x^\nu T^{\rho\mu}) \end{aligned}$$

Since the current must be conserved for all the six elements of the anti-symmetric matrix (since anti-symmetric so there are total $\frac{4 \times (4-1)}{2} = 6$ independent entries), the part of $x^\nu T^{\rho\mu}$ which is anti-symmetric in ρ and μ must be conserved. After antisymmetrising and considering current for each component, we have:

$$M^{\nu\mu\rho} := x^\nu T^{\mu\rho} - x^\rho T^{\mu\nu} \quad \partial_\mu M^{\nu\mu\rho} = 0$$

Let us now see the conserved charges for this case. Corresponding to spatial rotations, we have:

$$\mathcal{J}^{ij} = \int d^3\mathbf{x} M^{i0j} = \int d^3\mathbf{x} (x^i T^{0j} - x^j T^{0i})$$

This can be related to the angular-momentum since T^{0i} was related to P^i , the i^{th} component of the momentum. Hence, rotational invariance leads to the *conservation of angular momentum*. Let us see what happens when we give *boosts*.

$$\mathcal{J}^{0j} = \int d^3\mathbf{x} M^{00j} = \int d^3\mathbf{x} (x^0 T^{0j} - x^j T^{00})$$

As $t = x^0$ and we know that the current $\mathcal{J}^{0j}$ must be conserved, we have:

$$\frac{d\mathcal{J}^{0j}}{dt} = \frac{d}{dt} \int d^3\mathbf{x} (x^0 T^{0j} - x^j T^{00}) = \int d^3\mathbf{x} T^{0j} + \int d^3\mathbf{x} t \partial_t T^{0j} - \frac{d}{dt} \int d^3\mathbf{x} x^j T^{00} = 0$$

In terms of P^j , the above equation can be written as:

$$P^j + t \frac{dP^j}{dt} - \frac{d}{dt} \int d^3\mathbf{x} x^j T^{00} = 0$$

Previously we had seen that P^j was the conserved charge corresponding to spatial translation and hence P^j is a constant, which gives us:

$$\frac{d}{dt} \int d^3\mathbf{x} x^j T^{00} = \mathcal{C} \text{ (const.)}$$

The above relation can be seen as the centre of energy (or mass) of the field travels with constant velocity as T^{00} can be treated as the Hamiltonian ($\approx$ energy) density.

9.2 Examples: Internal Symmetries

Let us now look at internal symmetries, where the spacetime coordinates themselves do not change but the fields change due to some global transformations.

9.2.1 Global U(1) Symmetry of Complex Scalar Field

Consider a complex scalar field ϕ with the following Lagrangian:

$$\mathcal{L} = (\partial_\mu \phi)(\partial^\mu \phi^*) - m^2 \phi \phi^*$$

Let us consider a $U(1)$ transformation¹. Note that elements of the group $U(1)$ are represented by $e^{i\alpha}$, that is, they lie on the unit circle. Then, for an infinitesimal change $\psi \rightarrow e^{i\alpha}\psi$, $e^{i\alpha} = 1 + i\alpha + \mathcal{O}(\alpha^2)$ ² and we have:

$$\begin{aligned} \phi' &= (1 + i\alpha)\phi \implies \delta_\alpha \phi = i\alpha \phi \\ \phi^{*'} &= (1 - i\alpha)\phi^* \implies \delta_\alpha \phi^* = -i\alpha \phi^* \end{aligned}$$

Under this transformation, we see that the Lagrangian is invariant, that is,

$$\mathcal{L}' = (\partial_\mu \phi')(\partial^\mu \phi'^*) - m^2 \phi' \phi'^* = e^{i\alpha} e^{-i\alpha} (\partial_\mu \phi)(\partial^\mu \phi^*) - e^{i\alpha} e^{-i\alpha} m^2 \phi \phi^* = \mathcal{L}$$

From this, we can take $K^\mu = 0$. Using the formula for the conserved current, we have:

$$J^\mu = i\alpha \phi (\partial^\mu \phi^*) - i\alpha \phi^* (\partial^\mu \phi) \sim i(\partial^\mu \phi^* \phi - \partial^\mu \phi \phi^*)$$

The conserved charge is then given by the volume integral:

$$Q = \int d^3\mathbf{x} J^0 = -i \int d^3\mathbf{x} (\phi \dot{\phi}^* - \dot{\phi} \phi^*)$$

¹This is an *internal* symmetry as spacetime points are not varied.

²This is a *global* symmetry as α does not depend on the spacetime coordinates

Lecture 10 & 11: Field Quantisation

So far we had worked with classical fields. In the introduction, we saw that, quantum theories due to decoherence, transforms to classical theories, with an inevitable loss of information. Thus, finding the quantum theory leading to the classical model (working backwards) is not *unique*. More than one quantum field theory might have the same classical limit. Thus, while doing the quantisation process, we always have to make an *educated guess*!

Now, let us make the KG field *quantum* by quantising the canonical coordinates, similar to as we did for the harmonic oscillator.

We have our classical field $\phi(\mathbf{x}, t)$ and its canonical conjugate momentum $\pi(\mathbf{x}, t)$ which become operators $\hat{\phi}(x)$ and $\hat{\pi}(x)$ ¹. The classical Poisson bracket changes to the commutator, that is, $\{\cdot, \cdot\} \rightarrow i[\cdot, \cdot]$.

A quantum treatment of a classical Hamiltonian is sometimes messy since we have to maintain the Hermiticity condition and there can be multiple ways to do that. However, almost always, we just put the hats on appropriate places and wish we are correct!

Classically, we had $\{\phi(\mathbf{x}), \pi(\mathbf{y})\} = \delta^{(3)}(\mathbf{x} - \mathbf{y})$ and by the same logic:

$$[\hat{\phi}(\mathbf{x}), \hat{\pi}(\mathbf{y})] = i\delta^{(3)}(\mathbf{x} - \mathbf{y})$$

Let us take their Fourier transforms,

$$\begin{aligned}\tilde{\phi}(\mathbf{k}, t) &= \int d^3\mathbf{x} e^{-i\mathbf{k}\cdot\mathbf{x}} \phi(\mathbf{x}, t) \\ \tilde{\pi}(\mathbf{k}, t) &= \int d^3\mathbf{x} e^{-i\mathbf{k}\cdot\mathbf{x}} \pi(\mathbf{x}, t)\end{aligned}$$

Let us now consider the Klein-Gordon equation,

$$\mathcal{L} = \frac{1}{2}(\partial_\mu \phi)(\partial^\mu \phi) - \frac{1}{2}m^2 \phi^2 \iff (\partial_\mu \partial^\mu + m^2)\phi = 0$$

Expanding the field in Fourier space, we would obtain the following:

$$\left(\frac{\partial^2}{\partial t^2} + (|\mathbf{p}|^2 + m^2)\right) \tilde{\phi}(\mathbf{p}, t) = 0$$

Note that this is the equation for a classical harmonic oscillator with frequency $\omega_{\mathbf{p}} = \sqrt{|\mathbf{p}|^2 + m^2}$ and the general solution $\tilde{\phi}(\mathbf{x}, t)$ will be a linear combination of $e^{i\omega_{\mathbf{p}}t}$ and $e^{-i\omega_{\mathbf{p}}t}$. Thus we have,

$$\tilde{\phi}(\mathbf{p}, t) = A(\mathbf{p})e^{i\omega_{\mathbf{p}}t} + B(\mathbf{p})e^{-i\omega_{\mathbf{p}}t} \quad \omega_{\mathbf{p}} = \sqrt{|\mathbf{p}|^2 + m^2}$$

Now, in the quantum domain, $A(\mathbf{p})$ and $B(\mathbf{p})$ will become the creation and annihilation operators $a_{\mathbf{p}}$ and $a_{\mathbf{p}}^\dagger$ and in that spirit, let us define the field operators in terms of the annihilation and creation operators for each Fourier mode. Note that, for simplicity we define the field operators at $t = 0$.

$$\begin{aligned}\hat{\phi}(\mathbf{x}) &:= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} + \hat{a}_{\mathbf{p}}^\dagger e^{-i\mathbf{p}\cdot\mathbf{x}}) \\ \hat{\pi}(\mathbf{x}) &:= \int \frac{d^3\mathbf{p}}{(2\pi)^3} (-i) \sqrt{\frac{\omega_{\mathbf{p}}}{2}} (\hat{a}_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} - \hat{a}_{\mathbf{p}}^\dagger e^{-i\mathbf{p}\cdot\mathbf{x}})\end{aligned}$$

where $\omega_{\mathbf{p}} = \sqrt{|\mathbf{p}|^2 + m^2}$ is the frequency of the mode with momentum $\mathbf{p}$. Replacing $\mathbf{p}$ with $-\mathbf{p}$ in the second term, the above can also be written in a more compact way as:

$$\begin{aligned}\hat{\phi}(\mathbf{x}) &:= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger) e^{i\mathbf{p}\cdot\mathbf{x}} \\ \hat{\pi}(\mathbf{x}) &:= \int \frac{d^3\mathbf{p}}{(2\pi)^3} (-i) \sqrt{\frac{\omega_{\mathbf{p}}}{2}} (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger) e^{i\mathbf{p}\cdot\mathbf{x}}\end{aligned}$$

¹Note that we will write four vectors with non-bold letter, as x will denote the four components of the four-vector. To denote the three vectors, we will write in bold-face, that is, $\mathbf{x}$

Let us check the commutation relation:

$$[\hat{\phi}(\mathbf{x}), \hat{\pi}(\mathbf{y})] = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{(-i)}{2} \sqrt{\frac{\omega_{\mathbf{p}}}{\omega_{\mathbf{p}'}}} ([\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger, \hat{a}_{\mathbf{p}'} - \hat{a}_{-\mathbf{p}'}^\dagger]) e^{i(\mathbf{p}\cdot\mathbf{x} + \mathbf{p}'\cdot\mathbf{y})} = i\delta^{(3)}(\mathbf{x} - \mathbf{y})$$

Note that the above relation can be satisfied if we take:

$$[\hat{a}_{\mathbf{p}}, \hat{a}_{\mathbf{p}'}^\dagger] = (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}') \mathbb{1}$$

Now, let us do a horrendous calculation, which will then lead to a horrendous result and after some reasoning based on physical grounds, we will obtain a cute result. From the Hamiltonian density, we have the total Hamiltonian as:

$$\hat{H} = \int d^3\mathbf{x} \mathcal{H} = \int d^3\mathbf{x} \left(\frac{1}{2} \hat{\pi}^2 + \frac{1}{2} (\nabla \hat{\phi})^2 + \frac{1}{2} m^2 \hat{\phi}^2 \right)$$

Let us calculate this term by term. The first term gives us:

$$\begin{aligned} \int d^3\mathbf{x} \pi^2(\mathbf{x}) &= \int d^3\mathbf{x} \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}}{-2} (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} - \hat{a}_{-\mathbf{p}'}^\dagger) e^{i(\mathbf{p}+\mathbf{p}')\cdot\mathbf{x}} \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}}{-2} (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} - \hat{a}_{-\mathbf{p}'}^\dagger) \delta^{(3)}(\mathbf{p} + \mathbf{p}') \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[-\frac{\omega_{\mathbf{p}}}{2} \right] (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{-\mathbf{p}} - \hat{a}_{\mathbf{p}}^\dagger) \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[-\frac{\omega_{\mathbf{p}}}{2} \right] (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} - \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \end{aligned}$$

From the last term, we have:

$$\begin{aligned} \int d^3\mathbf{x} \phi^2(\mathbf{x}) &= \int d^3\mathbf{x} \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{1}{2\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}} (\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger) e^{i(\mathbf{p}+\mathbf{p}')\cdot\mathbf{x}} \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{1}{2\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}} (\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger) \delta^{(3)}(\mathbf{p} + \mathbf{p}') \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[\frac{1}{2\omega_{\mathbf{p}}} \right] (\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}^\dagger) \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left[\frac{1}{2\omega_{\mathbf{p}}} \right] (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \end{aligned}$$

Now comes the second term. For this, note that we have $\nabla(e^{i\mathbf{k}\cdot\mathbf{x}}) = i\mathbf{k}(e^{i\mathbf{k}\cdot\mathbf{x}})$. Using this, we obtain:

$$\begin{aligned} \int d^3\mathbf{x} (\nabla \phi) \cdot (\nabla \phi) &= \int d^3\mathbf{x} \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{1}{2\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}} [i\mathbf{p}(\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)e^{i\mathbf{p}\cdot\mathbf{x}}] \cdot [i\mathbf{p}'(\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger)e^{i\mathbf{p}'\cdot\mathbf{x}}] \\ &= - \int d^3\mathbf{x} \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{1}{2\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}} [(\mathbf{p} \cdot \mathbf{p}')(\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger)] e^{i(\mathbf{p}+\mathbf{p}')\cdot\mathbf{x}} \\ &= - \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} \frac{1}{2\sqrt{\omega_{\mathbf{p}}\omega_{\mathbf{p}'}}} [(\mathbf{p} \cdot \mathbf{p}')(\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger)] \delta^{(3)}(\mathbf{p} + \mathbf{p}') \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} [(\mathbf{p}^2)(\hat{a}_{\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger)(\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}^\dagger)] \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\mathbf{p}^2}{2\omega_{\mathbf{p}}} (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \end{aligned}$$

Adding all the terms with correct pre-factors, we obtain the following:

$$\begin{aligned}
\hat{H} &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \left\{ \left[-\frac{\omega_{\mathbf{p}}}{4} \right] (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} - \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \right\} + \left\{ \left[\frac{m^2}{4\omega_{\mathbf{p}}} \right] (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \right\} \\
&\quad + \left\{ \frac{\omega_{\mathbf{p}}^2 - m^2}{4\omega_{\mathbf{p}}} (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \right\} \\
&= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\omega_{\mathbf{p}}}{4} (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \\
&\quad + \frac{m^2}{4\omega_{\mathbf{p}}} (\hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{\mathbf{p}}\hat{a}_{-\mathbf{p}} - \hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}^\dagger) \\
&= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\omega_{\mathbf{p}}}{2} (\hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{-\mathbf{p}}^\dagger\hat{a}_{-\mathbf{p}}) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\omega_{\mathbf{p}}}{2} (\hat{a}_{\mathbf{p}}\hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}})
\end{aligned}$$

Now, using the commutator as obtained before, we get:

$$\hat{H} = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\omega_{\mathbf{p}}}{2} ((2\pi)^3 \delta^{(3)}(0) + 2\hat{a}_{\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \omega_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}} + \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{\omega_{\mathbf{p}}}{2} \delta^{(3)}(0)$$

This is the end of the horrendous calculation and now we discuss the horrendous term (the second term) which was obtained as a result of applying the commutator. Since this contains a delta-function (distribution?) and the integral is over all momentum, this definitely cannot be a finite quantity, hence it diverges. This term basically represents the zero-point energies of the harmonic oscillators (since there are infinite number of harmonic oscillators, we have a divergent zero-point energy). The fortunate thing is that this is a constant.

When physically measuring energies, only differences can be calculated and hence this term gets cancelled and thus, no divergence actually happens.¹

Thus, in all practicality, we can rescale our energy and we obtain the Hamiltonian as:

$$\hat{H} = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \omega_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger\hat{a}_{\mathbf{p}}$$

Now, let us see the commutator of the ladder operators with the Hamiltonian. We have:

$$[\hat{H}, \hat{a}_{\mathbf{p}}^\dagger] = \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} [\hat{a}_{\mathbf{p}'}^\dagger\hat{a}_{\mathbf{p}'}^\dagger, \hat{a}_{\mathbf{p}}^\dagger] = \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} \hat{a}_{\mathbf{p}'}^\dagger [\hat{a}_{\mathbf{p}'}^\dagger, \hat{a}_{\mathbf{p}}^\dagger] = \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} \hat{a}_{\mathbf{p}'}^\dagger (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}') = \omega_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger \quad (1)$$

$$[\hat{H}, \hat{a}_{\mathbf{p}}] = \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} [\hat{a}_{\mathbf{p}'}^\dagger\hat{a}_{\mathbf{p}'}^\dagger, \hat{a}_{\mathbf{p}}] = \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} [\hat{a}_{\mathbf{p}'}^\dagger, \hat{a}_{\mathbf{p}}] \hat{a}_{\mathbf{p}'} = - \int \frac{d^3\mathbf{p}'}{(2\pi)^3} \omega_{\mathbf{p}'} \hat{a}_{\mathbf{p}'} (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}') = -\omega_{\mathbf{p}} \hat{a}_{\mathbf{p}} \quad (2)$$

11.1 Conserved Charges

Note that we already found the Hamiltonian which is one of the conserved charges corresponding to T^{00} . Let us now focus on the momentum, that is, $\hat{P}^j$ corresponding to T^{0j}

$$\hat{P}^j = \int d^3\mathbf{x} \hat{T}^{0j} = - \int d^3\mathbf{x} \hat{\pi} \partial_j \hat{\phi}$$

Note that, while changing from the classical expression to quantum by ‘putting hats’, we had several choices for the ordering of $\hat{\pi}$ and $\hat{\phi}$ but here, we specify the one which works!

We now have two options, either to do the calculation again or just specify the result of this calculation

¹This infinite term is sometimes attributed to the *cosmological constant*. It also has some role in the *Casimir effect*.

since this had already been done by stalwarts before. We choose the former and thus, we have:

$$\begin{aligned} \int d^3\mathbf{x} \hat{\pi} \partial_j \hat{\phi} &= \int d^3\mathbf{x} \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{d^3\mathbf{p}'}{(2\pi)^3} (-i) \sqrt{\frac{\omega_{\mathbf{p}}}{2}} \frac{1}{\sqrt{2\omega_{\mathbf{p}'}}} (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger) i p'_j (\hat{a}_{\mathbf{p}'} + \hat{a}_{-\mathbf{p}'}^\dagger) e^{i(\mathbf{p}+\mathbf{p}')\cdot\mathbf{x}} \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \sqrt{\frac{\omega_{\mathbf{p}}}{2}} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger) p_j (\hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}}^\dagger) \\ &= \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} p_j (\hat{a}_{\mathbf{p}} \hat{a}_{-\mathbf{p}} + \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{-\mathbf{p}}^\dagger \hat{a}_{-\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}^\dagger) \end{aligned}$$

Now, note the following:

$$\mathcal{I} = \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} p_j (\hat{a}_{\mathbf{p}} \hat{a}_{-\mathbf{p}} - \hat{a}_{-\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}^\dagger) \quad \mathbf{p} \rightarrow -\mathbf{p} \quad \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} (-p_j) (\hat{a}_{-\mathbf{p}} \hat{a}_{\mathbf{p}} - \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{-\mathbf{p}}^\dagger) = -\mathcal{I}$$

The last equality comes since the creation and annihilation operators commute for different values of the momentum. Thus, this integral is zero and the expression for the total momentum becomes:

$$\hat{P}^j = \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} p_j (\hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger - \hat{a}_{-\mathbf{p}}^\dagger \hat{a}_{-\mathbf{p}}) = \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} p_j (\hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger + \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}) = \frac{1}{2} \int \frac{d^3\mathbf{p}}{(2\pi)^3} p_j \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}$$

In the last term, we introduced the commutator and removed the constant infinity term.

11.2 Normal Ordering

The way we are doing, like removing the infinities, is a common thing in QFT. This purely results from the fact the things commute in the classical case and hence there is always an order ambiguity when we move to quantum.

For example, if in classical scenario, we have some term like xp , when elevating (idk if its really elevating) these things to operators, we can have multiple possibilities, viz. $\hat{x}\hat{p}$, $\hat{p}\hat{x}$ or even $\frac{1}{2}(\hat{x}\hat{p} + \hat{p}\hat{x})$ (placing things symmetrically). Each one would present some additional constant on the final result, since the commutator of $\hat{p}$ and $\hat{x}$ is just a constant.

To be consistent, we generally use *normal ordering*, that is, keeping all the creation operators to the left. For that we define the *normal ordering operator*, $\hat{\mathcal{N}}$. For example,

$$\hat{\mathcal{N}}[\hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}] = \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}} \quad \hat{\mathcal{N}}[\hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger] = \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger \quad \hat{\mathcal{N}}[\hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger] = \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}} \quad \dots$$

The $\hat{\mathcal{N}}$ is not an operator in the sense that it doesn't act on a state to provide some new information. Instead, normal ordering is an interpretation that we use to eliminate the meaningless infinities that occur in field theories. So the rule goes that 'if you want to tell someone about a string of operators in quantum field theory, you have to normal order them first'. Thus, in operators like $\hat{P}$ and $\hat{H}$, in the final expression, we don't use the commutator but the normal operator operator, which will then completely subdue the issue of the appearance of Dirac-delta.

11.3 Constructing the Hilbert Space

The entire basis for this is the harmonic oscillator. In quantum SHO, we started with the vacuum and then acted creation operator successively on the vacuum to create excited states. Here, we will also do the same. Let us define the vacuum state $|\Omega\rangle$ such that,

$$\hat{a}_{\mathbf{p}} |\Omega\rangle = 0 \quad \forall \mathbf{p}$$

With such a definition, on physical grounds also, we obtain,

$$\hat{H} |\Omega\rangle = 0 \implies \text{Ground state has zero energy}$$

The vacuum can be written explicitly as:

$$|\Omega\rangle = \bigotimes_{\mathbf{p}} |\Omega\rangle_{\mathbf{p}} \quad |\Omega\rangle_{\mathbf{p}} \rightarrow \text{vac. of SHO with mom. } \mathbf{p}$$

The Hilbert space is then built as the space generated by all finite linear combinations of vectors of the form:

$$|\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle \equiv \hat{a}_{\mathbf{p}_1}^\dagger \hat{a}_{\mathbf{p}_2}^\dagger \dots \hat{a}_{\mathbf{p}_n}^\dagger |\Omega\rangle$$

Till now, this is not exactly the Hilbert space, since we did not account for the norm of the vectors, where a huge subtlety lies.

Note that $|\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle$ is an eigenstate of both $\hat{H}$ and $\hat{P}$ since,

$$\begin{aligned} \hat{H} |\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle &= (\omega_{\mathbf{p}_1} + \dots + \omega_{\mathbf{p}_n}) |\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle \\ \hat{P} |\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle &= (\mathbf{p}_1 + \dots + \mathbf{p}_n) |\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n\rangle \end{aligned}$$

Since the creation operators commute, the multi-particle states are symmetric under exchange of particles, that is, the particles follow *Bose-Einstein* statistics.

For a general many-particle system, this ‘pre’-Hilbert space is termed as *Fock space* $\mathfrak{F}$ and is formally defined with respect to direct-sums,

$$\mathfrak{F} = \bigoplus_{n=0}^{\infty} \mathcal{H}^{\otimes n}$$

where $\mathcal{H}^{\otimes n}$ is the n-times tensor product of the Hilbert space. In simple terms, $\mathcal{H}^{\otimes 0}$ is the space spanned by the vacuum state, $\mathcal{H}^{\otimes 1}$ is the space spanned by $|\mathbf{p}\rangle$. For $n \geq 2$, we have to consider the fermionic and bosonic properties too, that is, in our case, since these are bosons, $\mathcal{H}^{\otimes 2} = \text{Sym.}(\mathcal{H} \otimes \mathcal{H})$ is the space spanned by $|\mathbf{p}_1, \mathbf{p}_2\rangle$ such that Bose statistics is followed and so on...

Lecture 12: Making things invariant

The states that we had defined earlier, $|\mathbf{p}_1, \mathbf{p}_2, \dots\rangle$ are *improper*, in the sense that they cannot be normalised. The obvious choice, $\langle \mathbf{p} | \mathbf{p}' \rangle = (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}')$ is not physical due to the presence of the Dirac-delta. We can form physical states by ‘smearing out’ these deltas, by defining:

$$|\psi\rangle = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \psi(\mathbf{p}) |\mathbf{p}\rangle = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \psi(\mathbf{p}) \hat{a}_{\mathbf{p}}^\dagger |\Omega\rangle$$

where $\psi(\mathbf{p})$ is an $\mathbb{L}^2$ function. However, this smearing process is not a priori Lorentz invariant. Thus, we can force the process to follow Lorentz invariance since the inception only, so that we do not run into trouble later.

12.1 Invariant Inner Product

For simplicity, consider a Lorentz boost along the x direction. Then,

$$p'^1 = \gamma(p^1 + \beta E) \quad E' = \gamma(E + \beta p^1)$$

where $\gamma = 1/\sqrt{1-(v/c)^2}$ and $\beta = v/c$ are defined as usual. Now, note that $p^\mu p_\mu = m^2 \implies E^2 = (p^0)^2 + (p^1)^2 + (p^2)^2 + m^2 \implies 0 < E = \sqrt{(p^0)^2 + (p^1)^2 + (p^2)^2 + m^2}$ under the Minkowski metric. As the transformation is only along x direction, the momentum in x and y do not change. Thus, we have:

$$\frac{\partial E}{\partial p^j} = \frac{1}{2E} \times (2p^j) \implies \frac{\partial E}{\partial p^1} = \frac{p^1}{E}$$

Now, note the transformation of the Dirac-delta:

$$\int d^3\mathbf{y} \delta^{(3)}(\mathbf{x} - \mathbf{y}) = \int d^3\mathbf{y}' \delta^{(3)}(\mathbf{x}' - \mathbf{y}') = \int \left| \frac{\partial \mathbf{y}'}{\partial \mathbf{y}} \right| d^3\mathbf{y} \delta^{(3)}(\mathbf{x}' - \mathbf{y}')$$

Comparing, we see that the Dirac-delta transforms as $\delta^{(3)}(\mathbf{x} - \mathbf{y}) = \left| \frac{\partial \mathbf{y}'}{\partial \mathbf{y}} \right| \delta^{(3)}(\mathbf{x}' - \mathbf{y}')$ where $\left| \frac{\partial \mathbf{y}'}{\partial \mathbf{y}} \right|$ is the determinant of the Jacobian matrix J of the transformation. Hence, for the Lorentz transformation, we have:

$$\begin{aligned} \delta^{(3)}(\mathbf{p} - \mathbf{k}) &= \delta(p^1 - k^1) \delta(p^2 - k^2) \delta(p^3 - k^3) \\ &= \mathfrak{J} \delta(p'^1 - k'^1) \delta(p'^2 - k'^2) \delta(p'^3 - k'^3) \end{aligned}$$

Now, let us calculate the Jacobian matrix, whose elements will be given by $J^{ij} = \frac{\partial p'^i}{\partial p^j}$ (note that since we are concerned only with 3-momentum delta function, only spatial indices will appear). Note that $\frac{\partial p'^2}{\partial p^j} = \delta_{2j}$ and $\frac{\partial p'^3}{\partial p^j} = \delta_{3j}$. Only for $j = 1$, we have:

$$\frac{\partial p'^1}{\partial p^j} = \gamma \delta_{1j} + \gamma \beta \frac{\partial E}{\partial p^j}$$

and from the dispersion relation, we have $\frac{\partial E}{\partial p^j} = \frac{p^j}{E}$. The matrix is then:

$$J \equiv \begin{pmatrix} \gamma \left(1 + \beta \frac{p^1}{E}\right) & \gamma \beta \frac{p^2}{E} & \gamma \beta \frac{p^3}{E} \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \implies \mathfrak{J} = \det J = \gamma \left(1 + \beta \frac{p^1}{E}\right) = \frac{\gamma}{E} (E + \beta p^1) = \frac{E'}{E}$$

Thus, we obtain the transformation of the delta function as:

$$\delta^{(3)}(\mathbf{p} - \mathbf{k}) = \frac{E'}{E} \delta^{(3)}(\mathbf{p}' - \mathbf{k}') \implies E \delta^{(3)}(\mathbf{p} - \mathbf{k}) \text{ is invariant under L.T}$$

Then, we can now define the Lorentz-invariant inner product such that:

$$\langle \mathbf{p} | \mathbf{k} \rangle = (2E_{\mathbf{p}})(2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{k}) \equiv (2\omega_{\mathbf{p}})(2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{k})$$

In line with this, we can therefore define a single-particle state as:

$$|\mathbf{p}\rangle = \sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}}^\dagger |\Omega\rangle$$

Technically, we should have made some distinct notation to differentiate between the states with and without proper normalisation, however, henceforth we will always use normalised states, so confusion might not be there.

12.2 Invariant Measure

Define the *projector* onto the single-particle state,

$$\mathbb{1} := \int \frac{d^3 \mathbf{p}}{(2\pi)^3} |\mathbf{p}\rangle \langle \mathbf{p}|$$

Here we have used the states with proper normalisation, obeying Lorentz invariance. The LHS is Lorentz invariant, since the operator essentially checks whether we have single particle or not, which is a frame independent fact. Thus, the only thing to change is the *integral measure*!

Consider the following integral,

$$\int d^4 p \delta((p^0)^2 - \mathbf{p}^2 - m^2) \Theta(p^0)$$

Note that, $d^4 p$ is Lorentz invariant. Also, from the dispersion relation, we have:

$$p^\mu p_\mu = (p^0)^2 - \mathbf{p}^2 = m^2$$

The dispersion relation is also Lorentz invariant, since it contains scalars $p^\mu p_\mu$ and m^2 . Solving for p^0 , we have $p^0 = \pm \sqrt{\mathbf{p}^2 + m^2}$. Now, the choice of which branch to choose (positive or negative) is also Lorentz

invariant, since here we are considering only *orthochronous transformations*, which does not change the time component of the momentum. Thus, overall the integral is Lorentz-invariant.

$$\begin{aligned}
 \int d^4p \, \delta((p^0)^2 - \mathbf{p}^2 - m^2) \Theta(p^0) &= \int d^4p \, \delta((p^0)^2 - E_{\mathbf{p}}^2) \Theta(p^0) \\
 &= \int d^3\mathbf{p} \int_{-\infty}^{\infty} dp^0 \, \delta((p^0)^2 - E_{\mathbf{p}}^2) \Theta(p^0) \\
 &= \int d^3\mathbf{p} \int_{-\infty}^{\infty} dp^0 \, \frac{1}{2E_{\mathbf{p}}} [\delta(p^0 + E_{\mathbf{p}}) + \delta(p^0 - E_{\mathbf{p}})] \Theta(p^0) \\
 &= \int \frac{d^3\mathbf{p}}{2E_{\mathbf{p}}}
 \end{aligned}$$

Hence, whatever measure we were taking earlier, that is, $\frac{d^3\mathbf{p}}{(2\pi)^3}$ was not Lorentz-invariant, however, as seen above,

$$\frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2E_{\mathbf{p}}} \text{ is manifestly Lorentz invariant}$$

Nice! So, henceforth we will always try to write the integrals in manifestly Lorentz invariant form, for example, the field expression becomes:

$$\hat{\phi}(x) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} \left(\sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} + \sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}}^{\dagger} e^{-i\mathbf{p}\cdot\mathbf{x}} \right)$$

We could technically absorb the $\sqrt{2\omega_{\mathbf{p}}}$ in the definition of the creation and annihilation operators which would make the new field expressions similar to the old ones, but it is not really needed!

Finally, we can rewrite the identity operator on one-particle states as,

$$\mathbb{1} := \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} |\mathbf{p}\rangle \langle \mathbf{p}|$$

12.3 Position states?

Consider the field operator acting on the vacuum. Since annihilation operators kill the vacuum, we only have:

$$\hat{\phi}(x) |\Omega\rangle = \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} e^{-i\mathbf{p}\cdot\mathbf{x}} |p\rangle$$

Apart from the factor $\frac{1}{2\omega_{\mathbf{p}}}$, this looks more or less like some position state since it is linear superposition of all well-defined momentum states. Also, note that:

$$\begin{aligned}
 \langle \Omega | \hat{\phi}(\mathbf{x}) | \mathbf{p}' \rangle &= \langle \Omega | \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{2\omega_{\mathbf{p}}} \left(\sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}} + \sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}}^{\dagger} e^{-i\mathbf{p}\cdot\mathbf{x}} \right) \sqrt{2\omega_{\mathbf{p}}} \hat{a}_{\mathbf{p}'}^{\dagger} | \Omega \rangle \\
 &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} (e^{i\mathbf{p}\cdot\mathbf{x}}) \langle \Omega | \hat{a}_{\mathbf{p}} \hat{a}_{\mathbf{p}'}^{\dagger} | \Omega \rangle \\
 &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} (e^{i\mathbf{p}\cdot\mathbf{x}}) \langle \Omega | (2\pi)^3 \delta^{(3)}(\mathbf{p} - \mathbf{p}') + \hat{a}_{\mathbf{p}'}^{\dagger} \hat{a}_{\mathbf{p}} | \Omega \rangle \\
 &= e^{i\mathbf{p}'\cdot\mathbf{x}}
 \end{aligned}$$

We can interpret this as the position-space representation of the single-particle state $|\mathbf{p}\rangle$, exactly as in non-relativistic quantum mechanics. Thus, we have:

$$\langle \Omega | \hat{\phi}(\mathbf{x}) | \mathbf{p} \rangle = \langle x | p \rangle = e^{i\mathbf{p}\cdot\mathbf{x}}$$

Lecture 13: Heisenberg Picture

Till now, we saw that the operator $\hat{\phi}(\mathbf{x})$ has no time dependence (we defined it at $t = 0$). However, space and time should be on equal footing. Hence, it is better to define these in the Heisenberg picture, where the operators evolve in time while the states don't. Recall that a Heisenberg operator is defined from the Schrödinger operator via,

$$\mathcal{O}_H = e^{+iHt} \mathcal{O}_s e^{-iHt} \implies \frac{d\mathcal{O}_H}{dt} = i[H, \mathcal{O}_H] + \frac{\partial \mathcal{O}_H}{\partial t}$$

We consider here only free fields, that is, there is no interactions. However, in general, the Hamiltonian can be decomposed as

$$\hat{H} = \hat{H}_0 + \hat{H}_{\text{int.}}$$

where $\hat{H}_0$ is the *free field*. In that case, it would be better to work in the interaction picture. However, for now we assume $\hat{H}_{\text{int.}} = 0$. Note that from Eq. 1 and Eq. 2, we can obtain the time evolution of the ladder operators as:

$$\begin{aligned} \frac{d\hat{a}_{\mathbf{p}}^\dagger}{dt} &= i\omega_{\mathbf{p}} \hat{a}_{\mathbf{p}}^\dagger \implies \hat{a}_{\mathbf{p}}^\dagger(t) = \hat{a}_{\mathbf{p}}^\dagger(0) e^{i\omega_{\mathbf{p}} t} \implies e^{iH_0 t} \hat{a}_{\mathbf{p}}^\dagger e^{-iH_0 t} = \hat{a}_{\mathbf{p}}^\dagger e^{i\omega_{\mathbf{p}} t} \\ \frac{d\hat{a}_{\mathbf{p}}}{dt} &= -i\omega_{\mathbf{p}} \hat{a}_{\mathbf{p}} \implies \hat{a}_{\mathbf{p}}(t) = \hat{a}_{\mathbf{p}}(0) e^{-i\omega_{\mathbf{p}} t} \implies e^{iH_0 t} \hat{a}_{\mathbf{p}} e^{-iH_0 t} = \hat{a}_{\mathbf{p}} e^{-i\omega_{\mathbf{p}} t} \end{aligned}$$

Now, we calculate the field operator in the Heisenberg picture which gives:

$$\begin{aligned} \hat{\phi}(\mathbf{x}, t) &= e^{iH_0 t} \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}}^\dagger e^{-i\mathbf{p}\cdot\mathbf{x}} + \hat{a}_{\mathbf{p}} e^{i\mathbf{p}\cdot\mathbf{x}}) e^{-iH_0 t} \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (e^{iH_0 t} \hat{a}_{\mathbf{p}}^\dagger e^{-iH_0 t} e^{-i\mathbf{p}\cdot\mathbf{x}} + e^{iH_0 t} \hat{a}_{\mathbf{p}} e^{-iH_0 t} e^{i\mathbf{p}\cdot\mathbf{x}}) \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}}^\dagger e^{i\omega_{\mathbf{p}} t} e^{-i\mathbf{p}\cdot\mathbf{x}} + \hat{a}_{\mathbf{p}} e^{-i\omega_{\mathbf{p}} t} e^{i\mathbf{p}\cdot\mathbf{x}}) \\ &= \int \frac{d^3\mathbf{p}}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\mathbf{p}}}} (\hat{a}_{\mathbf{p}}^\dagger e^{ip\cdot x} + \hat{a}_{\mathbf{p}} e^{-ip\cdot x}) \end{aligned}$$

where $p \cdot x = p^\mu x_\mu = \omega_{\mathbf{p}} t - \mathbf{p} \cdot \mathbf{x}$ is the inner product of four momentum and four position in Minkowski space. Similarly, the conjugate momentum operator will be obtained as:

$$\hat{\pi}(\mathbf{x}, t) = \int \frac{d^3\mathbf{p}}{(2\pi)^3} (-i) \sqrt{\frac{\omega_{\mathbf{p}}}{2}} (\hat{a}_{\mathbf{p}} e^{-ip\cdot x} - \hat{a}_{\mathbf{p}}^\dagger e^{ip\cdot x})$$

Lecture 14 & 15: Propagators and Causality