

vv256: Week 2-3.  
Intervals of Existence. Direction fields.  
Autonomous equations. Singular solutions. Linear  
spaces.

Dr.Olga Danilkina

UM-SJTU Joint Institute

September 20, 2019

# Outline

- 1 Lecture 4: Intervals of Existence of a solution. Direction fields. Autonomous equations.
  - Direction fields
  - Intervals of Existence of a solution.
  - Autonomous Equations
- 2 Lecture 5: Intervals of Existence. Gronwall's and Bihari's inequalities.
- 3 Lecture 6: Implicit first-order ODEs. Singular solutions.
  - Implicit first-order ODEs
  - Singular Solutions

# Outline

- 1 Lecture 4: Intervals of Existence of a solution. Direction fields. Autonomous equations.
  - Direction fields
  - Intervals of Existence of a solution.
  - Autonomous Equations
- 2 Lecture 5: Intervals of Existence. Gronwall's and Bihari's inequalities.
- 3 Lecture 6: Implicit first-order ODEs. Singular solutions.
  - Implicit first-order ODEs
  - Singular Solutions

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No.

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE?

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

$$\bullet y' = f(t, y) \Rightarrow$$



# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

•  $y' = f(t, y) \Rightarrow$  What is  $f(t, y)$ ?

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

- $y' = f(t, y) \Rightarrow$  What is  $f(t, y)$ ?
- $f(x, y)$  is the slope of the tangent to the solution curve  $y = y(t)$  at a generic point  $(t, y)$ .

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

- $y' = f(t, y) \Rightarrow$  What is  $f(t, y)$ ?
- $f(x, y)$  is the slope of the tangent to the solution curve  $y = y(t)$  at a generic point  $(t, y)$ .
- How will we use it?

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!

- $y' = f(t, y) \Rightarrow$  What is  $f(t, y)$ ?
- $f(x, y)$  is the slope of the tangent to the solution curve  $y = y(t)$  at a generic point  $(t, y)$ .
- How will we use it?
- Choose a lattice/grid in the  $(t, y)$ -plane and draw short segments of the line with the slope  $f(t, y)$  at each node.

# Direction fields

Can we solve any nonlinear first-order ODE by means of integrals?

No. Is it possible to obtain information about the behavior of a solution/solutions without solving the ODE? Yes!


- $y' = f(t, y) \Rightarrow$  What is  $f(t, y)$ ?
- $f(x, y)$  is the slope of the tangent to the solution curve  $y = y(t)$  at a generic point  $(t, y)$ .
- How will we use it?
- Choose a lattice/grid in the  $(t, y)$ -plane and draw short segments of the line with the slope  $f(t, y)$  at each node.
- Thus obtained picture is called a **direction/slope field**.

## Direction fields

For example, for the equation  $y' = -5y$  the direction field is

## Direction fields

For example, for the equation  $y' = -5y$  the direction field is




## Direction fields

The direction field for  $y' = -5y$  with the solution satisfying  $y(0) = 1$ :



## Direction fields

The direction field for  $y' = -5y$  with the solution satisfying  $y(0) = 1$ :




## Direction fields

The direction field for  $y' = -5y$  with the solutions satisfying other ICs:

# Direction fields

The direction field for  $y' = -5y$  with the solutions satisfying other ICs:




# Direction fields

For the equation  $y' = y(4 - y)$  the direction field is

# Direction fields

For the equation  $y' = y(4 - y)$  the direction field is




## Direction fields

The direction field for  $y' = y(4 - y)$  with the solution satisfying  $y(0) = 1$ :

## Direction fields

The direction field for  $y' = y(4 - y)$  with the solution satisfying  $y(0) = 1$ :




# Direction fields

The direction field for  $y' = y(4 - y)$  with the solutions satisfying other ICs:




## Direction fields

The direction field for  $y' = y(4 - y)$  with the solutions satisfying other ICs:




## Direction fields

For the equation  $y' = 1 - 2t + (4t - 1)y - 2ty^2$  the direction field is




# Direction fields

The direction field for  $y' = 1 - 2t + (4t - 1)y - 2ty^2$  with the solution satisfying  $y(0) = 1$ :



## Direction fields

The direction field for  $y' = 1 - 2t + (4t - 1)y - 2ty^2$  with the solutions satisfying other ICs:



# Intervals of existence

**Theorem:** Let  $J$  be an open interval of the form  $a < t < b$  and  $t_0$  be a point in  $J$ . Consider the IVP

$$y' + p(t)y = q(t), y(t_0) = y_0,$$

where  $y_0$  is a given initial value.

If  $p$  and  $q$  are continuous on  $J$ , then the IVP has a unique solution on  $J$  for any  $y_0$ .

# Intervals of existence

**Theorem:** Let  $J$  be an open interval of the form  $a < t < b$  and  $t_0$  be a point in  $J$ . Consider the IVP

$$y' + p(t)y = q(t), y(t_0) = y_0,$$

where  $y_0$  is a given initial value.

If  $p$  and  $q$  are continuous on  $J$ , then the IVP has a unique solution on  $J$  for any  $y_0$ .

**Definition:** The largest open interval  $J$  on which an IVP has a unique solution is called the **maximal interval of existence** for that solution.

## Intervals of existence: Examples

Consider the IVP

$$y' - (t^2 + 1)y = \sin t, \quad y(1) = 2.$$

# Intervals of existence: Examples

Consider the IVP

$$y' - (t^2 + 1)y = \sin t, \quad y(1) = 2.$$

The functions  $p(t) = -(t^2 + 1)$ ,  $q(t) = \sin t$  are continuous on the interval  $(-\infty, +\infty) \Rightarrow$



# Intervals of existence: Examples

Consider the IVP

$$y' - (t^2 + 1)y = \sin t, \quad y(1) = 2.$$

The functions  $p(t) = -(t^2 + 1)$ ,  $q(t) = \sin t$  are continuous on the interval  $(-\infty, +\infty) \Rightarrow$  the IVP has a unique solution on the entire real line.

# Intervals of existence: Examples

Consider the IVP

$$y' - (t^2 + 1)y = \sin t, \quad y(1) = 2.$$

The functions  $p(t) = -(t^2 + 1)$ ,  $q(t) = \sin t$  are continuous on the interval  $(-\infty, +\infty) \Rightarrow$  the IVP has a unique solution on the entire real line.

For the IVP

$$(t + 1)y' + y = e^{2t}, \quad y(2) = 3,$$

$$p(t) = \frac{1}{t + 1}, \quad q(t) = \frac{e^{2t}}{t + 1} \text{ and hence,}$$

the functions  $p$ ,  $q$  are continuous on each of the open intervals  $-\infty < t < -1$  and  $-1 < t < +\infty$ .

# Intervals of existence: Examples

Consider the IVP

$$y' - (t^2 + 1)y = \sin t, \quad y(1) = 2.$$

The functions  $p(t) = -(t^2 + 1)$ ,  $q(t) = \sin t$  are continuous on the interval  $(-\infty, +\infty) \Rightarrow$  the IVP has a unique solution on the entire real line.

For the IVP

$$(t + 1)y' + y = e^{2t}, \quad y(2) = 3,$$

$$p(t) = \frac{1}{t + 1}, \quad q(t) = \frac{e^{2t}}{t + 1} \text{ and hence,}$$

the functions  $p$ ,  $q$  are continuous on each of the open intervals  $-\infty < t < -1$  and  $-1 < t < +\infty$ .

The IC is given at  $t_0 = 2 > -1$ , therefore the IVP has a unique solution in the interval  $-1 < t < +\infty$ .

## Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

## Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

- If  $t_0 \neq 0$ , then

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on



# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0,$$

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0, \quad a < t < b < 0, \quad t_0 < 0.$$

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const}$ .

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0, \quad a < t < b < 0, \quad t_0 < 0.$$

Direct verification shows that the maximal interval of existence is the entire real line.

# Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const}$ .

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0, \quad a < t < b < 0, \quad t_0 < 0.$$

Direct verification shows that the maximal interval of existence is the entire real line.

- If  $t_0 = 0$  but  $y_0 \neq 0$  then the IVP has no solution since

## Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const}$ .

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0, \quad a < t < b < 0, \quad t_0 < 0.$$

Direct verification shows that the maximal interval of existence is the entire real line.

- If  $t_0 = 0$  but  $y_0 \neq 0$  then the IVP has no solution since the equality  $y_0 = Ct_0 = 0$  is impossible for any value of  $C$ .

## Intervals of existence: Examples

Consider the IVP

$$ty' - y = 0, \quad y(t_0) = y_0$$

$p(t) = -1/t$  is not defined at the point  $t = 0$ , and  $q(t) = 0$  is continuous everywhere.

The general solution of the DE is  $y(t) = Ct$ ,  $C = \text{const.}$

- If  $t_0 \neq 0$ , then there exists a unique solution  $y(t) = \frac{y_0}{t_0}t$  on

$$0 < a < t < b, \quad t_0 > 0, \quad a < t < b < 0, \quad t_0 < 0.$$

Direct verification shows that the maximal interval of existence is the entire real line.

- If  $t_0 = 0$  but  $y_0 \neq 0$  then the IVP has no solution since the equality  $y_0 = Ct_0 = 0$  is impossible for any value of  $C$ .
- If  $t_0 = y_0 = 0$  then the IVP has infinitely many solutions, given by  $y(t) = Ct$  with any constant  $C$  on the entire real line.

# Intervals of existence

**Theorem.** Consider the IVP

$$y' = f(t, y), y(t_0) = y_0,$$

where  $f, f_y$  are continuous in an open rectangle  $R$  functions,  
 $R = \{(t, y) : a < t < b, c < y < d\}$ . If  $(t_0, y_0) \in R$  then the IVP  
has a unique solution in some open interval  $J$  of the form  
 $t_0 - h < t < t_0 + h$  contained in the interval  $a < t < b$ .

The conditions of the theorems are sufficient.

# Intervals of existence

**Theorem.** Consider the IVP

$$y' = f(t, y), y(t_0) = y_0,$$

where  $f, f_y$  are continuous in an open rectangle  $R$  functions,  $R = \{(t, y) : a < t < b, c < y < d\}$ . If  $(t_0, y_0) \in R$  then the IVP has a unique solution in some open interval  $J$  of the form  $t_0 - h < t < t_0 + h$  contained in the interval  $a < t < b$ .

The conditions of the theorems are sufficient. Are they necessary?

**Definition:** We say that a mathematical model is well-posed (correctly formulated) if there exists its unique solution.



## Intervals of existence: Examples

Consider the IVP

$$2(y - 1)y' = 2t + 1, y(2) = -1.$$

## Intervals of existence: Examples

Consider the IVP

$$2(y - 1)y' = 2t + 1, y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ .

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ . We can apply the theorem with any rectangle  $R$  that does not intersect the line  $y = 1$

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ . We can apply the theorem with any rectangle  $R$  that does not intersect the line  $y = 1 \Rightarrow 2-h < t < 2+h$ .

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ . We can apply the theorem with any rectangle  $R$  that does not intersect the line  $y = 1 \Rightarrow 2-h < t < 2+h$ .

Solve the equation:  $y(t) = 1 - (t^2 + t - 2)^{1/2}$ . **Why? Explain!**

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ . We can apply the theorem with any rectangle  $R$  that does not intersect the line  $y = 1 \Rightarrow 2-h < t < 2+h$ .

Solve the equation:  $y(t) = 1 - (t^2 + t - 2)^{1/2}$ . **Why? Explain!**

$$t^2 + t - 2 = (t-1)(t+2) \geq 0 \Rightarrow t \leq -2 \text{ or } t \geq 1.$$

# Intervals of existence: Examples

Consider the IVP

$$2(y-1)y' = 2t+1, \quad y(2) = -1.$$

What are the functions  $f, f_y$ ? Are they continuous?

$$f(t, y) = \frac{2t+1}{2(y-1)}, \quad f_y(t, y) = -\frac{2t+1}{2(y-1)^2}$$

Both continuous everywhere in the  $(t, y)$ -plane except on the line  $y = 1$ . We can apply the theorem with any rectangle  $R$  that does not intersect the line  $y = 1 \Rightarrow 2-h < t < 2+h$ .

Solve the equation:  $y(t) = 1 - (t^2 + t - 2)^{1/2}$ . Why? Explain!

$$t^2 + t - 2 = (t-1)(t+2) \geq 0 \Rightarrow t \leq -2 \text{ or } t \geq 1.$$

$t_0 = 2 \Rightarrow$  the maximal interval of existence  $1 < t < +\infty$ .



## Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

## Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

The point  $(0, 1)$  lies on the line  $y = 1$  where  $f, f_y$  are undefined.

# Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

The point  $(0, 1)$  lies on the line  $y = 1$  where  $f, f_y$  are undefined. So, the corresponding theorem does not work in this case!!!

# Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

The point  $(0, 1)$  lies on the line  $y = 1$  where  $f$ ,  $f_y$  are undefined. So, the corresponding theorem does not work in this case!!!

- Solving the IVP we obtain  $y(t) = 1 \pm (t^2 + t)^{1/2}$ .

# Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

The point  $(0, 1)$  lies on the line  $y = 1$  where  $f, f_y$  are undefined. So, the corresponding theorem does not work in this case!!!

- Solving the IVP we obtain  $y(t) = 1 \pm (t^2 + t)^{1/2}$ .
- This function is well-defined on  $-\infty < t < -1$  and  $0 < t < +\infty$ .

# Intervals of existence: Examples

What happens if the IC is replaced by  $y(0) = 1$ ?

The point  $(0, 1)$  lies on the line  $y = 1$  where  $f, f_y$  are undefined. So, the corresponding theorem does not work in this case!!!

- Solving the IVP we obtain  $y(t) = 1 \pm (t^2 + t)^{1/2}$ .
- This function is well-defined on  $-\infty < t < -1$  and  $0 < t < +\infty$ .
- Thus, there are two different solutions with the maximal interval of existence  $0 < t < +\infty$ . **Explain!**

# Intervals of existence: Exercises

Find the largest open interval on which the conditions of existence and uniqueness are satisfied, without solving the IVP itself.

- $(2t + 1)y' - 2y = \sin t, y(0) = -2$
- $(t^2 - 3t + 2)y' + ty = e^t, y(3/2) = -1$
- $(t^2 - t - 2)^{1/2}y' + 3y = (t - 3)^{1/2}, y(4) = 1$

Solve the DE with each of the given ICs and find the maximum interval of existence of the solution

- $y' = 4ty^2, y(0) = 2, y(-1) = -2, y(3) = -1$
- $(y - 2)y' = t, y(2) = 3, y(-2) = 1, y(0) = 1, y(1) = 2$
- $y' = 4(y - 1)^{1/2}, y(0) = 5, y(-1) = 2$

# Autonomous Equations: Definition

An autonomous equation is represented in the form

$$y' = f(y),$$

where the function  $f$  does not depend explicitly on  $t$ .

Is it a separable equation?



# Autonomous Equations: Definition

An autonomous equation is represented in the form

$$y' = f(y),$$

where the function  $f$  does not depend explicitly on  $t$ .

Is it a separable equation? Yes.

# Autonomous Equations: Definition

An autonomous equation is represented in the form

$$y' = f(y),$$

where the function  $f$  does not depend explicitly on  $t$ .

Is it a separable equation? Yes.

Did we consider autonomous equations before?

# Autonomous Equations: Definition

An autonomous equation is represented in the form

$$y' = f(y),$$

where the function  $f$  does not depend explicitly on  $t$ .

Is it a separable equation? Yes.

Did we consider autonomous equations before? Yes, the simplest population model is

$$y' - ry = 0, y(0) = y_0.$$

What are limitations of this model?

# Autonomous Equations: Definition

An autonomous equation is represented in the form

$$y' = f(y),$$

where the function  $f$  does not depend explicitly on  $t$ .

Is it a separable equation? Yes.

Did we consider autonomous equations before? Yes, the simplest population model is

$$y' - ry = 0, y(0) = y_0.$$

What are limitations of this model?

We need to make the model realistic!

# Population with logistic growth

Replace the constant rate  $r$  by a variable rate  $u(y)$ . Will it remain a physically plausible model?

# Population with logistic growth

Replace the constant rate  $r$  by a variable rate  $u(y)$ . Will it remain a physically plausible model? Yes, if  $u(y)$  satisfies the following conditions:

# Population with logistic growth

Replace the constant rate  $r$  by a variable rate  $u(y)$ . Will it remain a physically plausible model? Yes, if  $u(y)$  satisfies the following conditions:

- $u(y) \approx r$  for small  $y$ ,
- $u(y)$  decreases as  $y$  increases,
- $u(y) < 0$  for large  $y$ .

Derive the simplest function  $u(y)$ .

# Population with logistic growth

Replace the constant rate  $r$  by a variable rate  $u(y)$ . Will it remain a physically plausible model? Yes, if  $u(y)$  satisfies the following conditions:

- $u(y) \approx r$  for small  $y$ ,
- $u(y)$  decreases as  $y$  increases,
- $u(y) < 0$  for large  $y$ .

Derive the simplest function  $u(y)$ .

$$u(y) = r - ay, \quad a = \text{const} > 0.$$



# Population with logistic growth

$$u(y) = r - ay, \quad a = \text{const} > 0.$$

We obtain the logistic equation of the form

$$y' = r(1 - \frac{y}{B})y, \quad B = r/a.$$

# Population with logistic growth

$$u(y) = r - ay, \quad a = \text{const} > 0.$$

We obtain the logistic equation of the form

$$y' = r(1 - \frac{y}{B})y, \quad B = r/a.$$

What are the roots of the algebraic equation  $f(y) = 0$ ?

# Population with logistic growth

$$u(y) = r - ay, \quad a = \text{const} > 0.$$

We obtain the logistic equation of the form

$$y' = r(1 - \frac{y}{B})y, \quad B = r/a.$$

What are the roots of the algebraic equation  $f(y) = 0$ ?

$$f(y) = r(1 - y/B)y \Rightarrow y = 0, y = B \leftarrow \text{critical points/equilibrium sol.}$$

# Population with logistic growth

$$u(y) = r - ay, \quad a = \text{const} > 0.$$

We obtain the logistic equation of the form

$$y' = r(1 - \frac{y}{B})y, \quad B = r/a.$$

What are the roots of the algebraic equation  $f(y) = 0$ ?

$$f(y) = r(1 - y/B)y \Rightarrow y = 0, y = B \leftarrow \text{critical points/equilibrium sol.}$$

The solution of the logistic equation is

$$\frac{y}{B - y} = Ce^{rt}, \quad C = \text{const} \neq 0. \quad \text{Obtain it!}$$

# Population with logistic growth

If  $y(0) = y_0 > 0$  then the solution is

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{-rt}}.$$

Is the function  $y(t)$  well-defined?

# Population with logistic growth

If  $y(0) = y_0 > 0$  then the solution is

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{-rt}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $t > 0$ . Check what happens if  $y_0 > B$ !

# Population with logistic growth

If  $y(0) = y_0 > 0$  then the solution is

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{-rt}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $t > 0$ . Check what happens if  $y_0 > B$ ! Letting  $t \rightarrow \infty$ , we obtain

$$\lim_{t \rightarrow \infty} y(t) = B.$$

What does it mean?

# Population with logistic growth

If  $y(0) = y_0 > 0$  then the solution is

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{-rt}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $t > 0$ . Check what happens if  $y_0 > B$ ! Letting  $t \rightarrow \infty$ , we obtain

$$\lim_{t \rightarrow \infty} y(t) = B.$$

What does it mean?

$B$  is the size of the largest population that the environment can sustain long term  $\Rightarrow B$  is the environmental carrying capacity.



# Population with logistic growth

An equilibrium solution  $y_0$  is called

- **stable** if any other solution starting close to  $y_0$  remains close to  $y_0$  for all time.
- **asymptotically stable** if it is stable and any solution starting close to  $y_0$  becomes arbitrarily close to  $y_0$  as  $t$  increases.

An equilibrium solution that is not stable is called **unstable**.

Are equilibrium solutions  $y_0 = 0$  and  $y_0 = B$  stable or unstable?  
 $y_0 = B$  is asymptotically stable and  $y_0 = 0$  is unstable.

# Population with logistic growth: Example

Consider the IVP

$$y' = 400y - 4y^2, \quad y(0) = y_0.$$

## Population with logistic growth: Example

Consider the IVP

$$y' = 400y - 4y^2, \quad y(0) = y_0.$$

The critical points/equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 100$  and hence,

# Population with logistic growth: Example

Consider the IVP

$$y' = 400y - 4y^2, \quad y(0) = y_0.$$

The critical points/equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 100$  and hence,

$$y(t) = \frac{100y_0}{y_0 + (100 - y_0)e^{-400t}}.$$

# Population with logistic growth: Example

Consider the IVP

$$y' = 400y - 4y^2, \quad y(0) = y_0.$$

The critical points/equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 100$  and hence,

$$y(t) = \frac{100y_0}{y_0 + (100 - y_0)e^{-400t}}.$$


Sketch the graph of the obtained solution:

- $y' > 0$  for  $0 < y < 100$ , and  $y' < 0$  when  $y > 100$ ,
- $y'' = 32y(50 - y)(100 - y)$  and hence,  
 $y'' > 0$ ,  $0 < y < 50$ ,  $y > 100$  and  $y'' < 0$ ,  $50 < y < 100$ ,

and let  $y_0$  be 25, 75, 125 to obtain the particular solutions

$$y_1(t) = \frac{100}{1 + 3e^{-400t}}, \quad y_2(t) = \frac{300}{3 + e^{-400t}}, \quad y_3(t) = \frac{500}{5 - e^{-400t}}$$

## Population with logistic growth: Example



## Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

# Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population,



# Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population, and a sample problem

$$y' = 28 \left( 1 - \frac{y}{7} \right) y - 40 = 4(y - 2)(5 - y).$$

# Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population, and a sample problem

$$y' = 28 \left( 1 - \frac{y}{7} \right) y - 40 = 4(y - 2)(5 - y).$$

The equilibrium solutions are

# Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population, and a sample problem

$$y' = 28 \left( 1 - \frac{y}{7} \right) y - 40 = 4(y - 2)(5 - y).$$

The equilibrium solutions are  $y(t) = 2$ ,  $y(t) = 5$  and the general solution has the form

# Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population, and a sample problem

$$y' = 28 \left( 1 - \frac{y}{7} \right) y - 40 = 4(y - 2)(5 - y).$$

The equilibrium solutions are  $y(t) = 2$ ,  $y(t) = 5$  and the general solution has the form

$$\frac{y - 2}{y - 5} = C e^{12t}, \quad C = \text{const} \neq 0.$$

## Logistic population with harvesting

Consider a mathematical model of logistic population with harvesting governed by the DE of the form

$$y' = r \left( 1 - \frac{y}{B} \right) y - \alpha,$$

where  $\alpha = \text{const} > 0$  is the rate of harvesting  $\Rightarrow \alpha$  contributes to the decline of the population, and a sample problem

$$y' = 28 \left( 1 - \frac{y}{7} \right) y - 40 = 4(y - 2)(5 - y).$$

The equilibrium solutions are  $y(t) = 2$ ,  $y(t) = 5$  and the general solution has the form

$$\frac{y - 2}{y - 5} = C e^{12t}, \quad C = \text{const} \neq 0.$$

Any IC of the form  $y(0) = y_0$  yields  $C = (y_0 - 2)/(y_0 - 5)$  and

## Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined?

# Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $y_0 > 2$ . Check what happens if  $0 < y_0 < 2$ ,  $2 < y_0 < 5$ ,  $y_0 > 5$ !

# Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $y_0 > 2$ . Check what happens if  $0 < y_0 < 2$ ,  $2 < y_0 < 5$ ,  $y_0 > 5$ !

$$\lim_{t \rightarrow \infty} y(t) = 5.$$



# Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $y_0 > 2$ . Check what happens if  $0 < y_0 < 2$ ,  $2 < y_0 < 5$ ,  $y_0 > 5$ !

$$\lim_{t \rightarrow \infty} y(t) = 5.$$

For  $0 < y_0 < 2$ , the solution  $y(t) = 0$  at the point

$$t^* = \frac{1}{12} \ln \frac{2(5 - y_0)}{5(2 - y_0)} > 0.$$

# Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $y_0 > 2$ . Check what happens if  $0 < y_0 < 2$ ,  $2 < y_0 < 5$ ,  $y_0 > 5$ !

$$\lim_{t \rightarrow \infty} y(t) = 5.$$

For  $0 < y_0 < 2$ , the solution  $y(t) = 0$  at the point

$$t^* = \frac{1}{12} \ln \frac{2(5 - y_0)}{5(2 - y_0)} > 0.$$

Populations starting with a size less than 2 are not large enough to survive and die out in finite time.

# Logistic population with harvesting

$$y(t) = \frac{5(y_0 - 2) - 2(y_0 - 5)e^{-12t}}{y_0 - 2 - (y_0 - 5)e^{-12t}}.$$

Is the function  $y(t)$  well-defined? Yes, for all  $y_0 > 2$ . Check what happens if  $0 < y_0 < 2$ ,  $2 < y_0 < 5$ ,  $y_0 > 5$ !


$$\lim_{t \rightarrow \infty} y(t) = 5.$$

For  $0 < y_0 < 2$ , the solution  $y(t) = 0$  at the point

$$t^* = \frac{1}{12} \ln \frac{2(5 - y_0)}{5(2 - y_0)} > 0.$$

Populations starting with a size less than 2 are not large enough to survive and die out in finite time. Check what happens with the solution at other points  $t!!!$  We plot the particular solutions for  $y_0 = 1, 3, 4, 6$ .

## Logistic population with harvesting: Example



## Population with a critical threshold

The model of a population with a critical threshold is of the form

$$y' = -r \left( 1 - \frac{y}{B} \right) y, \quad r, B = \text{const} > 0$$

# Population with a critical threshold

The model of a population with a critical threshold is of the form

$$y' = -r \left( 1 - \frac{y}{B} \right) y, \quad r, B = \text{const} > 0$$

with the solution

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0) e^{rt}} \quad \text{for } y(0) = y_0.$$

## Population with a critical threshold

The model of a population with a critical threshold is of the form

$$y' = -r \left( 1 - \frac{y}{B} \right) y, \quad r, B = \text{const} > 0$$

with the solution

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{rt}} \quad \text{for } y(0) = y_0.$$

What happens when  $y_0 < B$  and  $y_0 > B$ ?

- If  $y_0 < B$ , then the solution is well-defined for all  $t > 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ .

## Population with a critical threshold

The model of a population with a critical threshold is of the form

$$y' = -r \left( 1 - \frac{y}{B} \right) y, \quad r, B = \text{const} > 0$$

with the solution

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0) e^{rt}} \quad \text{for } y(0) = y_0.$$

What happens when  $y_0 < B$  and  $y_0 > B$ ?

- If  $y_0 < B$ , then the solution is well-defined for all  $t > 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ .
- For  $y_0 > B$ , the solution is not defined at the point

$$t^* = \frac{1}{r} \ln \frac{y_0}{y_0 - B} > 0.$$

What does it mean?



## Population with a critical threshold

The model of a population with a critical threshold is of the form

$$y' = -r \left(1 - \frac{y}{B}\right) y, \quad r, B = \text{const} > 0$$

with the solution

$$y(t) = \frac{y_0 B}{y_0 + (B - y_0)e^{rt}} \quad \text{for } y(0) = y_0.$$

What happens when  $y_0 < B$  and  $y_0 > B$ ?

- If  $y_0 < B$ , then the solution is well-defined for all  $t > 0$  and  $\lim_{t \rightarrow \infty} y(t) = 0$ .
- For  $y_0 > B$ , the solution is not defined at the point

$$t^* = \frac{1}{r} \ln \frac{y_0}{y_0 - B} > 0.$$

What does it mean? A population that starts with a size above the value of  $B$  grows without bound in finite time.

## Population with a critical threshold

A population that starts with a size above the value of  $B$  grows without bound in finite time. The number  $B$  in this model is called a critical threshold.

## Population with a critical threshold

A population that starts with a size above the value of  $B$  grows without bound in finite time. The number  $B$  in this model is called a critical threshold.

Consider an example.

$$y' = -40 \left(1 - \frac{y}{20}\right) y = 2y(y - 20).$$

The equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 20$ .

# Population with a critical threshold

A population that starts with a size above the value of  $B$  grows without bound in finite time. The number  $B$  in this model is called a critical threshold.

Consider an example.

$$y' = -40 \left(1 - \frac{y}{20}\right) y = 2y(y - 20).$$

The equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 20$ . A solution that corresponds to the IC  $y(0) = y_0$  is

$$y(t) = \frac{20y_0}{y_0 - (y_0 - 20)e^{40t}}.$$

# Population with a critical threshold

A population that starts with a size above the value of  $B$  grows without bound in finite time. The number  $B$  in this model is called a critical threshold.

Consider an example.

$$y' = -40 \left(1 - \frac{y}{20}\right) y = 2y(y - 20).$$


The equilibrium solutions are  $y(t) = 0$ ,  $y(t) = 20$ . A solution that corresponds to the IC  $y(0) = y_0$  is

$$y(t) = \frac{20y_0}{y_0 - (y_0 - 20)e^{40t}}.$$

Sketch the particular solutions for the cases  $y_0 = 5, 15, 25$ :

$$y_1(t) = \frac{20}{1 + 3e^{40t}}, \quad y_1(t) = \frac{60}{3 + e^{40t}}, \quad y_1(t) = \frac{100}{5 - 3e^{40t}}.$$

## Population with a critical threshold: Example



# Exercises

Find the critical points and equilibrium solutions of the given DE and solve the DE with each of the prescribed ICs. Sketch the graphs of the solutions obtained and comment on the stability/instability of the equilibrium solutions. Identify the model governed by the IVP, if any, and describe its main elements.

- $y' = 300y - 2y^2, y(0) = 50, y(0) = 100, y(0) = 200$
- $y' = 240y - 3y^2, y(0) = 20, y(0) = 60, y(0) = 100$
- $y' = 15y - y^2/2, y(0) = 10, y(0) = 20, y(0) = 40$
- $y' = 8y - 2y^2 - 6, y(0) = 1/2, y(0) = 3/2, y(0) = 5/2, y(0) = 7/2$
- $y' = y^2 + y - 6, y(0) = -4, y(0) = -2, y(0) = 1, y(0) = 3$

# Outline

- 1 Lecture 4: Intervals of Existence of a solution. Direction fields. Autonomous equations.
  - Direction fields
  - Intervals of Existence of a solution.
  - Autonomous Equations
- 2 Lecture 5: Intervals of Existence. Gronwall's and Bihari's inequalities.
- 3 Lecture 6: Implicit first-order ODEs. Singular solutions.
  - Implicit first-order ODEs
  - Singular Solutions



## Lemma 1: Gronwall-Bellman

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$\forall t \geq t_0 \quad u(t) \leq C + \int_{t_0}^t f(t_1) u(t_1) dt_1,$$

where  $C$  is a positive constant.

Then

$$\forall t \geq t_0 \quad u(t) \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right). \quad (1)$$

**Proof.**

## Lemma 1: Gronwall-Bellman

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$\forall t \geq t_0 \quad u(t) \leq C + \int_{t_0}^t f(t_1)u(t_1) dt_1,$$

where  $C$  is a positive constant.

Then

$$\forall t \geq t_0 \quad u(t) \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right). \quad (1)$$

**Proof.**

$$\frac{u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq 1 \Rightarrow$$

## Lemma 1: Gronwall-Bellman

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$\forall t \geq t_0 \quad u(t) \leq C + \int_{t_0}^t f(t_1)u(t_1) dt_1,$$

where  $C$  is a positive constant.

Then

$$\forall t \geq t_0 \quad u(t) \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right). \quad (1)$$

**Proof.**

$$\frac{u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq 1 \Rightarrow \frac{f(t)u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq f(t)$$

## Lemma 1: Gronwall-Bellman

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$\forall t \geq t_0 \quad u(t) \leq C + \int_{t_0}^t f(t_1)u(t_1) dt_1,$$

where  $C$  is a positive constant.

Then

$$\forall t \geq t_0 \quad u(t) \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right). \quad (1)$$

**Proof.**

$$\frac{u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq 1 \Rightarrow \frac{f(t)u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq f(t)$$

Integrate the last inequality with respect to  $t$  from  $t_0$  to  $t$ .

## Lemma 1: Gronwall-Bellman

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$\forall t \geq t_0 \quad u(t) \leq C + \int_{t_0}^t f(t_1)u(t_1) dt_1,$$

where  $C$  is a positive constant.

Then

$$\forall t \geq t_0 \quad u(t) \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right). \quad (1)$$

**Proof.**

$$\frac{u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq 1 \Rightarrow \frac{f(t)u(t)}{C + \int_{t_0}^t f(t_1)u(t_1) dt_1} \leq f(t)$$

Integrate the last inequality with respect to  $t$  from  $t_0$  to  $t$ . Since

$$\frac{d}{dt} \left[ C + \int_{t_0}^t f(t_1)u(t_1) dt_1 \right] = f(t)u(t),$$

## Lemma 1: Gronwall-Bellman

we obtain

$$\ln \left| C + \int_{t_0}^t f(t_1) u(t_1) dt_1 \right| - \ln C \leq \int_{t_0}^t f(t) dt.$$

Therefore,

$$u(t) \leq C + \int_{t_0}^t f(t_1) u(t_1) dt_1 \leq C \exp \left( \int_{t_0}^t f(t_1) dt_1 \right).$$

## Lemma 1: Gronwall-Bellman

**Corollary:** Let  $u(t)$  be a positive continuous function satisfying

$$u(t) \leq u(\tau) + \int_{\tau}^t f(t_1)u(t_1) dt_1 \quad \forall t, \tau \in (a, b),$$

where  $f(t) \in C(a, b)$  and  $f(t) \geq 0 \forall t \in (a, b)$ .

Then for all  $a < t_0 \leq t < b$

$$u(t_0) \exp \left[ - \int_{t_0}^t f(t_1) dt_1 \right] \leq u(t) \leq u(t_0) \exp \left[ \int_{t_0}^t f(t_1) dt_1 \right].$$

**Proof:** Leave as an exercise.

## Lemma 2: Bihari- LaSalle

Let  $u(t) \geq 0$ ,  $f(t) \geq 0$  for all  $t \geq t_0$ ,  $u(t), f(t) \in C[t_0, +\infty)$ , and

$$u(t) \leq C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1,$$

where  $C$  is a positive constant,  $\Phi(u)$  is a positive non-decreasing continuous function for all  $0 < u < \bar{u}$  ( $\bar{u} \leq \infty$ ). Define

$$\Psi(u) = \int_C^u \frac{du_1}{\Phi(u_1)}, \quad 0 < u < \bar{u}.$$

If  $\int_{t_0}^t f(t_1) dt_1 < \Psi(\bar{u} - 0)$ ,  $t_0 \leq t < \infty$ , then

$$u(t) \leq \Psi^{-1} \left[ \int_{t_0}^t f(t_1) dt_1 \right] \quad \forall t_0 \leq t < \infty \quad (2)$$



## Bihari-LaSalle: Proof

**Proof.** Since  $\Phi$  is non-decreasing, so

## Bihari-LaSalle: Proof

**Proof.** Since  $\Phi$  is non-decreasing, so

$$\Phi(u(t)) \leq \Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)$$

and hence,

## Bihari-LaSalle: Proof

**Proof.** Since  $\Phi$  is non-decreasing, so

$$\Phi(u(t)) \leq \Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)$$

and hence,

$$\frac{f(t) \Phi(u(t))}{\Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)} \leq f(t).$$

## Bihari-LaSalle: Proof

**Proof.** Since  $\Phi$  is non-decreasing, so

$$\Phi(u(t)) \leq \Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)$$

and hence,

$$\frac{f(t) \Phi(u(t))}{\Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)} \leq f(t).$$

Integrating with respect to  $t$  from  $t_0$  to  $t$ , we have

$$\int_{t_0}^t \frac{w'(t) dt}{\Phi(w(t))} = \int_{w(t_0)}^{w(t)} \frac{dw}{\Phi(w(t))} \leq \int_{t_0}^t f(t_1) dt_1,$$

where  $w(t) = C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1$ .

## Bihari-LaSalle: Proof

**Proof.** Since  $\Phi$  is non-decreasing, so

$$\Phi(u(t)) \leq \Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)$$

and hence,

$$\frac{f(t) \Phi(u(t))}{\Phi \left( C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 \right)} \leq f(t).$$

Integrating with respect to  $t$  from  $t_0$  to  $t$ , we have

$$\int_{t_0}^t \frac{w'(t) dt}{\Phi(w(t))} = \int_{w(t_0)}^{w(t)} \frac{dw}{\Phi(w(t))} \leq \int_{t_0}^t f(t_1) dt_1,$$

where  $w(t) = C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1$ . Since  $w(t_0) = C > 0$

and  $w(t) \geq C > 0$  then  $\Psi(w(t)) \leq \int_{t_0}^t f(t_1) dt_1$ .

## Bihari-LaSalle: Proof

Recall that  $\Psi'(u) = \frac{1}{\Phi(u)} > 0$  for all  $0 < u < \bar{u}$ .

## Bihari-LaSalle: Proof

Recall that  $\Psi'(u) = \frac{1}{\Phi(u)} > 0$  for all  $0 < u < \bar{u}$ . Therefore, the function  $v = \Psi(u)$  has a unique inverse function  $u = \Psi^{-1}(v)$  (continuous and monotonically increasing) defined on  $\Psi(+0) < v < \Psi(\bar{u} - 0)$ , where  $\Psi(+0) < 0$ .

## Bihari-LaSalle: Proof

Recall that  $\Psi'(u) = \frac{1}{\Phi(u)} > 0$  for all  $0 < u < \bar{u}$ . Therefore, the function  $v = \Psi(u)$  has a unique inverse function  $u = \Psi^{-1}(v)$  (continuous and monotonically increasing) defined on  $\Psi(+0) < v < \Psi(\bar{u} - 0)$ , where  $\Psi(+0) < 0$ . We can express  $w(t)$  from the last inequality provided  $\int_{t_0}^t f(t_1) dt_1 < \Psi(\bar{u} - 0)$  as follows

$$w(t) \leq \Psi^{-1} \left[ \int_{t_0}^t f(t_1) dt_1 \right]$$

It gives

$$u(t) \leq C + \int_{t_0}^t f(t_1) \Phi(u(t_1)) dt_1 = w(t) \leq \Psi^{-1} \left[ \int_{t_0}^t f(t_1) dt_1 \right].$$



## Bihari-LaSalle: intervals of existence

Find an interval of existence of the solution for the following IVP

$$y' = t + y^3, \quad y(0) = 0.$$

**Solution.**

We have  $f(t, y) = t + y^3$ ,  $\frac{\partial f}{\partial y}(t, y) = 3y^2$ ,  $(t_0, y_0) = (0, 0)$ . **What are the conditions for existence of the unique solution?**  $f, f_y$  are continuous in any rectangular  $R$  that contains  $(t_0, y_0) = (0, 0)$ , for example  $R = \{(t, y) \in \mathbb{R}^2 : |t| \leq a, |y| \leq b\}$ .  $\Rightarrow$  there exists a unique solution of the problem defined on  $0 - h < t < 0 + h$ . **How can we find the interval of existence without solving the equation?** Use Bihari's inequality.

The equation implies that

$$y(t) = \int_0^t (t_1 + y^3(t_1)) dt_1 = \frac{t^2}{2} + \int_0^t y^3(t_1) dt_1$$

## Bihari-LaSalle: intervals of existence

$$|y(t)| \leq \frac{a^2}{2} + \int_0^t |y(t_1)|^3 dt_1.$$

Denote  $u(t) = |y(t)|$ ,  $C = a^2/2$ ,  $f(t) = 1$ ,  $\Phi(u) = u^3$ . Then

$$v = \Psi(u) = \int_C^u \frac{du_1}{u_1^3} = \frac{1}{2} \left( \frac{1}{C^2} - \frac{1}{u^2} \right)$$

$$u = \Psi^{-1}(v) = \frac{C}{\sqrt{1 - 2C^2v}}$$

Then the Bihari inequality implies that

$$u(t) \leq \Psi^{-1} \left[ \int_0^t 1 dt_1 \right] = \frac{C}{\sqrt{1 - 2C^2t}}.$$

That is,

$$|y(t)| \leq \frac{C}{\sqrt{1 - 2C^2t}}, \forall t > 0.$$

## Bihari-LaSalle: intervals of existence

What is the restriction for  $t$ ?

$$0 < t < \frac{1}{2C^2}$$

From the equation  $a = \frac{1}{2C^2}$  find that  $\max a = \sqrt[5]{2}$ . Then the solution exists on the interval  $[0, \sqrt[5]{2})$ .

Can we also extend the interval of existence along the negative part of the  $t$ -axis? Yes, change  $t$  to  $-t$  ( $t \geq 0$ ) in the equation and proceed with the same procedure.

We obtain  $u(t) \leq \frac{C}{\sqrt{1+2C^2t}}$ ,  $t \leq 0$ .  $\Rightarrow$  the solution exists for  $-\sqrt[5]{2} < t < 0$  as well.

**Answer:** We can guarantee the existence of the unique solution in the interval  $-\sqrt[5]{2} < t < \sqrt[5]{2}$ .

## Bihari-LaSalle: intervals of existence

Using Bihari's lemma, find an interval of existence of the solution for the following IVPs:

$$1. y' = 2y^2 - t, y(1) = 1.$$

$$2. \begin{cases} y_1' = y_2^2, & y_1(0) = 1, \\ y_2' = y_1^2, & y_2(0) = 2. \end{cases}$$

# Outline

- 1 Lecture 4: Intervals of Existence of a solution. Direction fields. Autonomous equations.
  - Direction fields
  - Intervals of Existence of a solution.
  - Autonomous Equations
- 2 Lecture 5: Intervals of Existence. Gronwall's and Bihari's inequalities.
- 3 Lecture 6: Implicit first-order ODEs. Singular solutions.
  - Implicit first-order ODEs
  - Singular Solutions

# Implicit first-order ODEs

Consider the ODE

$$F(t, y, y') = 0, \quad (3)$$

where  $F$  is a known smooth function. Consider the following cases

# Implicit first-order ODEs

Consider the ODE

$$F(t, y, y') = 0, \quad (3)$$

where  $F$  is a known smooth function. Consider the following cases

- The derivative  $y'$  can be expressed explicitly in, say,  $k$  ways

$$y' = f_1(t, y), y' = f_2(t, y), \dots, y' = f_k(t, y) \quad (4)$$

# Implicit first-order ODEs

Consider the ODE

$$F(t, y, y') = 0, \quad (3)$$

where  $F$  is a known smooth function. Consider the following cases

- The derivative  $y'$  can be expressed explicitly in, say,  $k$  ways

$$y' = f_1(t, y), y' = f_2(t, y), \dots, y' = f_k(t, y) \quad (4)$$

**What does it mean?** It means that there are  $k$  integral curves passing through each point of the domain  $G$  where right-hand sides of the equations are defined.



# Implicit first-order ODEs

Consider the ODE

$$F(t, y, y') = 0, \quad (3)$$

where  $F$  is a known smooth function. Consider the following cases

- The derivative  $y'$  can be expressed explicitly in, say,  $k$  ways

$$y' = f_1(t, y), y' = f_2(t, y), \dots, y' = f_k(t, y) \quad (4)$$

**What does it mean?** It means that there are  $k$  integral curves passing through each point of the domain  $G$  where right-hand sides of the equations are defined. Let

$$\Phi_1(t, y, C) = 0, \Phi_2(t, y, C) = 0, \dots, \Phi_k(t, y, C) = 0$$

be general solutions of (4).

# Implicit first-order ODEs

Consider the ODE

$$F(t, y, y') = 0, \quad (3)$$

where  $F$  is a known smooth function. Consider the following cases

- The derivative  $y'$  can be expressed explicitly in, say,  $k$  ways

$$y' = f_1(t, y), y' = f_2(t, y), \dots, y' = f_k(t, y) \quad (4)$$

**What does it mean?** It means that there are  $k$  integral curves passing through each point of the domain  $G$  where right-hand sides of the equations are defined. Let

$$\Phi_1(t, y, C) = 0, \Phi_2(t, y, C) = 0, \dots, \Phi_k(t, y, C) = 0$$

be general solutions of (4). **What is the general solution of the equation (3)?**

$$\Phi_1(t, y, C) \cdot \Phi_2(t, y, C) \cdot \dots \cdot \Phi_k(t, y, C) = 0$$

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $t$ :  $t = \varphi(y, y')$ .

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $t$ :  $t = \varphi(y, y')$ . Introduce a parameter  $p = y'$ , that is  $t = \varphi(y, p)$ .

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $t$ :  $t = \varphi(y, y')$ . Introduce a parameter  $p = y'$ , that is  $t = \varphi(y, p)$ . Differentiate it w.r.t  $y$  to obtain

$$\frac{1}{p} = \frac{\partial \varphi}{\partial y} + \frac{\partial \varphi}{\partial p} \cdot \frac{dp}{dy} \Rightarrow \frac{dy}{dp} = \frac{p \frac{\partial \varphi}{\partial p}}{1 - p \frac{\partial \varphi}{\partial y}}$$

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $t$ :  $t = \varphi(y, y')$ . Introduce a parameter  $p = y'$ , that is  $t = \varphi(y, p)$ . Differentiate it w.r.t  $y$  to obtain

$$\frac{1}{p} = \frac{\partial \varphi}{\partial y} + \frac{\partial \varphi}{\partial p} \cdot \frac{dp}{dy} \Rightarrow \frac{dy}{dp} = \frac{p \frac{\partial \varphi}{\partial p}}{1 - p \frac{\partial \varphi}{\partial y}}$$

If a general solution of this equation has the representation  $y = \Theta(p, C)$ , where  $\Theta$  is known and  $C$  is a constant, then

$$\begin{cases} t = \varphi(\Theta(p, C), p) \\ y = \Theta(p, C) \end{cases}$$

is the general solution of the equation  $t = \varphi(y, y')$  in the parametric form. **Non-parametric form?**  $\Rightarrow$  **Eliminate  $p$ .**

**Exercise:** Find the general solution if  $t = \varphi(y')$ .

# Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $y$  :  $y = \psi(t, y')$ .

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $y$  :  $y = \psi(t, y')$ .  
Letting  $p = y'$  we obtain  $y = \psi(t, p)$ .



## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $y$  :  $y = \psi(t, y')$ .  
Letting  $p = y'$  we obtain  $y = \psi(t, p)$ . Again, we differentiate the obtained equation w.r.t.  $t$

$$p = \frac{\partial \psi}{\partial t} + \frac{\partial \psi}{\partial p} \cdot \frac{dp}{dt} \Rightarrow$$

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $y$  :  $y = \psi(t, y')$ .  
Letting  $p = y'$  we obtain  $y = \psi(t, p)$ . Again, we differentiate the obtained equation w.r.t.  $t$

$$p = \frac{\partial \psi}{\partial t} + \frac{\partial \psi}{\partial p} \cdot \frac{dp}{dt} \Rightarrow \frac{dt}{dp} = \frac{\frac{\partial \psi}{\partial p}}{p - \frac{\partial \psi}{\partial t}}$$

## Implicit first-order ODEs

- The eq. (3) can be solved with respect to  $y$  :  $y = \psi(t, y')$ .  
Letting  $p = y'$  we obtain  $y = \psi(t, p)$ . Again, we differentiate the obtained equation w.r.t.  $t$

$$p = \frac{\partial \psi}{\partial t} + \frac{\partial \psi}{\partial p} \cdot \frac{dp}{dt} \Rightarrow \frac{dt}{dp} = \frac{\frac{\partial \psi}{\partial p}}{p - \frac{\partial \psi}{\partial t}}$$

If a general solution  $t = \Theta(p, C)$  of this equation exists then

$$\begin{cases} t = \Theta(p, C) \\ y = \psi(\Theta(p, C), p) \end{cases}$$

is the general solution of the equation  $y = \psi(t, y')$  in the parametric form.

What happens if  $y = \psi(y')$ ?

# Implicit first-order ODEs

What is common in cases 2 and 3?

- In both cases equations are explicit with respect to either  $t$  or  $y$ , and
- we differentiate w.r.t. another variable.
- New equations are explicit w.r.t. corresponding derivatives

However, new explicit equations may not have analytical representation of the solution!!!

We are to consider two types of equations for which the approach described above works and explicit equations are solvable.

# Lagrange Equation

The equation

$$y = t\varphi(y') + \psi(y')$$

is called the Lagrange equation.

# Lagrange Equation

The equation

$$y = t\varphi(y') + \psi(y')$$

is called the Lagrange equation.

We apply the procedure from the Case 3 ( $y' = p$ ) to obtain

# Lagrange Equation

The equation

$$y = t\varphi(y') + \psi(y')$$

is called the Lagrange equation.

We apply the procedure from the Case 3 ( $y' = p$ ) to obtain

$$(\varphi(p) - p)\frac{dt}{dp} + \varphi'(p)t + \psi'(p) = 0.$$

# Lagrange Equation

The equation

$$y = t\varphi(y') + \psi(y')$$

is called the Lagrange equation.

We apply the procedure from the Case 3 ( $y' = p$ ) to obtain

$$(\varphi(p) - p)\frac{dt}{dp} + \varphi'(p)t + \psi'(p) = 0.$$

**What is the type of this equation?** Linear  $\Rightarrow$  Find its solution  $t = \Phi(p, C)$  and obtain the general solution of the Lagrange equation in the form

$$\begin{cases} t = \Phi(p, C) \\ y = \Phi(p, C)\varphi(p) + \psi(p) \end{cases}$$

Attention! The Lagrange equation may also have special solutions of the form  $y = \varphi(c)t + \psi(c)$ , where  $c$  is the root of the equation  $\varphi(c) = 0$ . We will consider the question of special solutions later.



## Clairaut equation

If  $\varphi(y') = y'$  in the Lagrange equation then the equation

$$y = ty' + \psi(y')$$

is called the Clairaut equation.

## Clairaut equation

If  $\varphi(y') = y'$  in the Lagrange equation then the equation

$$y = ty' + \psi(y')$$

is called the Clairaut equation.

Letting  $y' = p$  and differentiating, we obtain  $p = C$  and  $t = -\psi'(p)$ . **Verify it.**

## Clairaut equation

If  $\varphi(y') = y'$  in the Lagrange equation then the equation

$$y = ty' + \psi(y')$$

is called the Clairaut equation.

Letting  $y' = p$  and differentiating, we obtain  $p = C$  and  $t = -\psi'(p)$ . **Verify it.**

What is the general solution?  $y = Ct + \psi(C)$ .

## Clairaut equation

If  $\varphi(y') = y'$  in the Lagrange equation then the equation

$$y = ty' + \psi(y')$$

is called the Clairaut equation.

Letting  $y' = p$  and differentiating, we obtain  $p = C$  and  $t = -\psi'(p)$ . **Verify it.**

What is the general solution?  $y = Ct + \psi(C)$ .

Eliminating the parameter  $p$  from the system

$$\begin{cases} t = -\psi'(p) \\ y = pt + \psi(p) \end{cases}$$

we obtain the singular solution as well.

## Clairaut equation

If  $\varphi(y') = y'$  in the Lagrange equation then the equation

$$y = ty' + \psi(y')$$

is called the Clairaut equation.

Letting  $y' = p$  and differentiating, we obtain  $p = C$  and  $t = -\psi'(p)$ . **Verify it.**

What is the general solution?  $y = Ct + \psi(C)$ .

Eliminating the parameter  $p$  from the system

$$\begin{cases} t = -\psi'(p) \\ y = pt + \psi(p) \end{cases}$$

we obtain the singular solution as well.

Therefore, plugging  $C$  instead of  $y'$  in Clairaut's equation we immediately obtain the general solution. **How we can get a singular solution from the general one? Differentiate w.r.t  $C$ .**

## Examples

1. Find the solution of the equation  $F(y') = 0$ .

Let  $y' = p$  and  $F(p_0) = 0$ . Then  $F(\frac{y - C}{t}) = 0$ .

2. Solve the equation  $(y')^2 - (t + y)y' + ty = 0$ .

We have  $y' = t$  and  $y' = y$ . Then  $y = \frac{t^2}{2} + C$ ,  $y = Ce^t$  and the general solution is  $(y - \frac{t^2}{2} - C)(y - Ce^t) = 0$ .

3. Consider the equation  $y = y' + (y')^2 e^{y'}$ .

Case 2  $\Rightarrow y' = p$  and  $y = p + p^2 e^p$ . Therefore,  $dt = \frac{1+(p^2+2p)e^p}{p} dp$  and  $t = \ln |p| + (p + 1)e^p + C$ . The general solution has the form

$$\begin{cases} t = \ln |p| + (p + 1)e^p + C \\ y = p + p^2 e^p \end{cases}$$

Moreover, we need to complement it with the obvious solution  $y = 0$ .

## Exercises

Solve the following ODEs:

$$1. (y')^2 - 2ty' - 8t^2 = 0.$$

$$2. t^2(y')^2 + 3tyy' + 2y^2 = 0.$$

$$3. (y')^3 - y(y')^2 - t^2y' + t^2y = 0.$$

$$4. t = \ln y' + \sin y'.$$

$$5. y = \sin^{-1} y' + \ln(1 + (y')^2).$$

$$6. y = ty' + y' + \sqrt{y'}.$$

$$7. y = y' \ln y'.$$

$$y = 3/2ty' + e^{y'}.$$

# Singular Solutions

We have already considered ODEs with singular solutions (check examples above). Intuitively, a singular solution is a special solution that is not contained in the general solution for any values of the constant  $C$  including  $C = \pm\infty$ . **What do singular solution mean geometrically? How can we plot them?**

**Definition.** A solution  $y = \varphi(t)$  of the differential equation

$$F(t, y, y') = 0 \quad (5)$$

is called singular if the uniqueness property does not hold at any of its points, that is,

- there is another solution of the same ODEs passing through each point  $(t_0, y_0)$  of the singular solution, and
- both solutions have the same tangent at the point  $(t_0, y_0)$  but
- another non-singular solution is different from the singular one in any arbitrary small neighborhood of the point  $(t_0, y_0)$ .



# Singular Solutions

Does a singular solution satisfies the equation (5)? Yes. Moreover, if  $F(t, y, y')$ ,  $\frac{\partial F}{\partial y}$ ,  $\frac{\partial F}{\partial y'}$  are continuous with respect to all arguments  $t, y, y'$  then any singular solution satisfies the equation

$$\frac{\partial F(t, y, y')}{\partial y'} = 0. \quad (6)$$

How can we find a singular solutions from (5) and (6)?  $\Rightarrow$   
Eliminate  $y'$ . Elimination gives us an equation

$$\psi_p(t, y) = 0$$

which is called  $p$ -discriminant of the equation (5), and the integral curve corresponding  $p$ -discriminant is called the  $p$ -discriminant integral curve.

Is a  $p$ -discriminant curve unique? Does it define a singular solution? In general, no.  $\Rightarrow$  Double-check.


# Singular Solutions

Consider the equation

$$\Phi(t, y, C) = 0$$

with a parameter  $C$  and continuous  $\Phi_t, \Phi_y, \Phi_C$ . It defines a family of curves depending on one parameter.

An **envelope** of the family of curves with a parameter is a smooth curve  $\Gamma$  that touches one curve of the family at any of its points and any its segment is touched by an infinite number of curves from the family. **What does it mean if curves touch? A common tangent.**



# Singular Solutions

Does the envelope satisfies the definition of a singular integral curve? Yes.  $\Rightarrow$  the envelope defines a singular solution.

An envelope is a part of a  $C$ -discriminant curve defined a by

$$\begin{cases} \Phi(t, y, C) = 0 \\ \frac{\partial \Phi(t, y, C)}{\partial C} = 0 \end{cases}$$

To make sure that a branch of a  $C$ -discriminant curve is an envelope, we check the following conditions.

- there exist bounded partial derivatives
$$\left| \frac{\partial \Phi}{\partial t} \right| \leq M, \left| \frac{\partial \Phi}{\partial y} \right| \leq N, M, N = \text{const},$$
- $\frac{\partial \Phi}{\partial t} \neq 0$ , or,  $\frac{\partial \Phi}{\partial y} \neq 0$

Are these condition are necessary or sufficient? Sufficient.  $\Rightarrow$  if they are not satisfied on a branch of the  $C$ -discriminant curve, it can still be an envelope.




# Singular Solutions

The equations of  $p$ -discriminant and  $C$ -discriminant have a certain structure

$$\psi_p(t, y) = E \cdot C \cdot T^2 = 0,$$

$$\psi_C(t, y) = E \cdot N^2 \cdot C^3 = 0,$$

where  $E = 0$  is the equation of the envelope,  $C = 0$  is the equation of the cusp locus,  $N = 0$  is the equation of nodal locus,  $T = 0$  is the equation of the tac locus. Attention! Over all locus points only the envelope is a singular solution.



# Singular Solutions: Example 1

Consider the equation

$$ty' + (y')^2 - y = 0.$$

1. Find its  $p$ -discriminant curve

$$y = -\frac{t^2}{4}.$$

2. Check if it is a solution of the DE.
3. Check if it is a singular solution. Find the general solution of the equation  $y = Ct + C^2$ . **Why?** Check the type of the equation. If two curves  $y = y_1(t)$  and  $y = y_2(t)$  touch at the point  $t = t_0$  then


$$y_1(t_0) = y_2(t_0), y_1'(t_0) = y_2'(t_0).$$

## Singular Solutions: Example 1

It gives us

$$-\frac{t_0^2}{4} = Ct_0 + C^2, \quad -\frac{t_0}{2} = C.$$

and hence,  $-\frac{t_0^2}{4} = -\frac{t_0^2}{4} \Rightarrow$  at each point of the curve  $y = -\frac{t^2}{4}$ ,  
another curve of the form  $y = Ct + C^2$  touches it, with  $C = -\frac{t_0}{2}$ .  
Therefore,  $y = -\frac{t^2}{4}$  is a singular solution.




## Singular Solutions: Example 2

Find singular solutions of the equation

$$t(y')^2 - 2yy' + 4t = 0, \quad t > 0$$

with the general solution  $t^2 = C(y - C)$ .

1. Find the  $C$ -discriminant curve  $y = \pm 2t$
2. Verify that both functions are solutions of the equation.
3. Prove that each function is a singular solution. Use the sufficient condition.




## Singular Solutions: Example 3

Find a singular solution of the equation

$$(y')^2 = 4t^2.$$

1.  $p$ -discriminant is  $2y' = 0$ ,  $(y')^2 = 4t^2$  and hence,  $t^2 = 0$ .
2. Is  $t = 0$  an integral curve of the equation? No. But it may be tac points locus. Why?
3. The general solution of the equation is  $y = \pm t^2 + C$ . Therefore, the line  $t = 0$  is indeed the tac points locus.






## Singular Solutions: Example 4

Find singular solution of the equation

$$(y')^2(2 - 3y)^2 = 4(1 - y).$$

1. Find the  $p$ -discriminant  $(2 - 3y)^2(1 - y) = 0$ . What are the conclusions?
2. Find the general solution of the equation  $y^2(1 - y) = (t - C)^2$ .
3. Find the  $C$ -discriminant curve  $y^2(1 - y) = 0$ .



## Singular Solutions: Exercises

For the following equations, find singular solutions if they exist.

1.  $(1 + (y')^2)y^2 - 4yy' - 4t = 0,$

2.  $(y')^2 - 4y = 0,$

3.  $(y')^3 - 4tyy' + 8y^2 = 0,$

4.  $(y')^2 - y^2 = 0,$

5.  $(ty' + y)^2 + 3t^5(ty' - 2y) = 0.$

Use  $C$ -discriminant to find singular solutions for the following equations 1.  $y = (y')^2 - ty' + t^2/2, y = Ct + C^2 + t^2/2,$

2.  $(ty' + y)^2 = y^2y', y(C - t) = C^2,$

3.  $y^2(y')^2 + y^2 = 1, (x - C)^2 + y^2 = 1,$

4.  $(y')^2 - yy' + e^t = 0, y = Ce^t + 1/C.$