

vv256: Laplace Transform

Dr.Olga Danilkina

UM-SJTU Joint Institute



November 29, 2019

General Integral Transforms

The Laplace Transform is the method for solving IVPs for ODEs and PDEs by converting the differential equation into an algebraic equation.

General Integral Transforms

The **Laplace Transform** is the method for **solving** IVPs for ODEs and PDEs by **converting the differential equation into an algebraic equation**.

A **general integral transform** of a function $f(t)$ into another function $\bar{f}(s)$ is defined by

$$\bar{f}(s) = \int_{\alpha}^{\beta} K(s, t) f(t) dt.$$

General Integral Transforms

The **Laplace Transform** is the method for **solving** IVPs for ODEs and PDEs by **converting the differential equation into an algebraic equation**.

A **general integral transform** of a function $f(t)$ into another function $\bar{f}(s)$ is defined by

$$\bar{f}(s) = \int_{\alpha}^{\beta} K(s, t) f(t) dt.$$

The function $K(s, t)$ is called the **kernel** of the transform.

General Integral Transforms

The **Laplace Transform** is the method for **solving** IVPs for ODEs and PDEs by **converting the differential equation into an algebraic equation**.

A **general integral transform** of a function $f(t)$ into another function $\bar{f}(s)$ is defined by

$$\bar{f}(s) = \int_{\alpha}^{\beta} K(s, t) f(t) dt.$$

The function $K(s, t)$ is called the **kernel** of the transform.

The Laplace transform is a special case with $\alpha = 0$, $\beta = \infty$, $K(s, t) = e^{-st} \Rightarrow$

General Integral Transforms

The **Laplace Transform** is the method for **solving** IVPs for ODEs and PDEs by **converting the differential equation into an algebraic equation**.

A **general integral transform** of a function $f(t)$ into another function $\bar{f}(s)$ is defined by

$$\bar{f}(s) = \int_{\alpha}^{\beta} K(s, t) f(t) dt.$$

The function $K(s, t)$ is called the **kernel** of the transform.

The Laplace transform is a special case with $\alpha = 0$, $\beta = \infty$, $K(s, t) = e^{-st} \Rightarrow$ improper integral

We assume that $f(t)$ as a real-valued function defined for $t > 0$.

Definition of the Laplace Transform

The Laplace transform of $f(t)$ is

$$\bar{f}(s) = L[f(t): t \rightarrow s] = \int_0^{\infty} e^{-st} f(t) dt.$$

The function $K(s, t)$ is called the kernel of the transform.

Definition of the Laplace Transform

The Laplace transform of $f(t)$ is

$$\bar{f}(s) = L[f(t): t \rightarrow s] = \int_0^{\infty} e^{-st} f(t) dt.$$

The function $K(s, t)$ is called the kernel of the transform.

Reverse the procedure and obtain the inverse Laplace transform: if $L[f(t): t \rightarrow s]$ then

$$L^{-1} [\bar{f}(s): s \rightarrow t] = f(t).$$

Definition of the Laplace Transform

The Laplace transform of $f(t)$ is

$$\bar{f}(s) = L[f(t): t \rightarrow s] = \int_0^{\infty} e^{-st} f(t) dt.$$

The function $K(s, t)$ is called the kernel of the transform.

Reverse the procedure and obtain the inverse Laplace transform: if $L[f(t): t \rightarrow s]$ then

$$L^{-1} [\bar{f}(s): s \rightarrow t] = f(t).$$

1. What functions have a Laplace transform?

Definition of the Laplace Transform

The Laplace transform of $f(t)$ is

$$\bar{f}(s) = L[f(t): t \rightarrow s] = \int_0^{\infty} e^{-st} f(t) dt.$$

The function $K(s, t)$ is called the kernel of the transform.

Reverse the procedure and obtain the inverse Laplace transform: if $L[f(t): t \rightarrow s]$ then

$$L^{-1} [\bar{f}(s): s \rightarrow t] = f(t).$$

1. What functions have a Laplace transform?
2. Can two functions $f(t)$, $g(t)$ have the same Laplace transform?

Definition of the Laplace Transform

The **Laplace transform** of $f(t)$ is

$$\bar{f}(s) = L[f(t): t \rightarrow s] = \int_0^{\infty} e^{-st} f(t) dt.$$

The function $K(s, t)$ is called the **kernel** of the transform.

Reverse the procedure and obtain the **inverse Laplace transform**: if $L[f(t): t \rightarrow s]$ then

$$L^{-1} [\bar{f}(s): s \rightarrow t] = f(t).$$

1. What functions have a Laplace transform?
2. Can two functions $f(t)$, $g(t)$ have the same Laplace transform?

The **existence and uniqueness of the Laplace transform** of a function $f(t)$ is guaranteed if there exist real K , M and a such that

1. $f(t)$ is piecewise continuous for $t > 0$,
2. $|f(t)| \leq Ke^{at}$ for $t \geq M$.

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$L[t: t \rightarrow s] = \int_0^{\infty} te^{-st} =$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$L[t: t \rightarrow s] = \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt =$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$L[t: t \rightarrow s] = \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \frac{1}{s} \left. \frac{e^{-st}}{-s} \right|_0^{\infty}$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$\begin{aligned} L[t: t \rightarrow s] &= \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \left. \frac{1}{s} \frac{e^{-st}}{-s} \right|_0^{\infty} \\ &= \frac{1}{s^2}, \quad s > 0 \end{aligned}$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$\begin{aligned} L[t: t \rightarrow s] &= \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \frac{1}{s} \left. \frac{e^{-st}}{-s} \right|_0^{\infty} \\ &= \frac{1}{s^2}, \quad s > 0 \end{aligned}$$

Example 3. Find the Laplace transform of the function $f(t) = e^{\alpha t}$.

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$\begin{aligned} L[t: t \rightarrow s] &= \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \left. \frac{1}{s} \frac{e^{-st}}{-s} \right|_0^{\infty} \\ &= \frac{1}{s^2}, \quad s > 0 \end{aligned}$$

Example 3. Find the Laplace transform of the function $f(t) = e^{\alpha t}$.

$$L[e^{\alpha t}: t \rightarrow s] = \int_0^{\infty} e^{\alpha t} e^{-st} =$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$\begin{aligned} L[t: t \rightarrow s] &= \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \left. \frac{1}{s} \frac{e^{-st}}{-s} \right|_0^{\infty} \\ &= \frac{1}{s^2}, \quad s > 0 \end{aligned}$$

Example 3. Find the Laplace transform of the function $f(t) = e^{\alpha t}$.

$$L[e^{\alpha t}: t \rightarrow s] = \int_0^{\infty} e^{\alpha t} e^{-st} = \left. \frac{e^{(\alpha-s)t}}{\alpha-s} \right|_0^{\infty} =$$

Examples

Example 1. Find the Laplace transform of the function $f(t) = 1$.

$$L[1: t \rightarrow s] = \int_0^{\infty} e^{-st} = \left. \frac{e^{-st}}{-s} \right|_0^{\infty} = \frac{1}{s}$$

Remark: The result is limited to $s > 0$ only.

Example 2. Find the Laplace transform of the function $f(t) = t$.

$$\begin{aligned} L[t: t \rightarrow s] &= \int_0^{\infty} te^{-st} = \left. \frac{te^{-st}}{-s} \right|_0^{\infty} - \int_0^{\infty} \frac{e^{-st}}{(-s)} dt = \left. \frac{1}{s} \frac{e^{-st}}{-s} \right|_0^{\infty} \\ &= \frac{1}{s^2}, \quad s > 0 \end{aligned}$$

Example 3. Find the Laplace transform of the function $f(t) = e^{\alpha t}$.

$$L[e^{\alpha t}: t \rightarrow s] = \int_0^{\infty} e^{\alpha t} e^{-st} = \left. \frac{e^{(\alpha-s)t}}{\alpha-s} \right|_0^{\infty} = \frac{1}{s-\alpha}, \quad s > \alpha$$

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

$$L [\cos at + i \sin at : t \rightarrow s] = L [e^{iat} : t \rightarrow s] =$$

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

$$L[\cos at + i \sin at: t \rightarrow s] = L[e^{iat}: t \rightarrow s] = \frac{1}{s - ia} =$$

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

$$L[\cos at + i \sin at: t \rightarrow s] = L[e^{iat}: t \rightarrow s] = \frac{1}{s - ia} =$$

$$= \frac{s + ia}{s^2 + a^2}, \quad s > 0$$

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

$$L[\cos at + i \sin at: t \rightarrow s] = L[e^{iat}: t \rightarrow s] = \frac{1}{s - ia} =$$

$$= \frac{s + ia}{s^2 + a^2}, \quad s > 0$$

Separate real and imaginary parts:

Examples

Example 4. Find the Laplace transform of the functions $f(t) = \sin at$ and $f(t) = \cos at$.

$$L[\cos at + i \sin at: t \rightarrow s] = L[e^{iat}: t \rightarrow s] = \frac{1}{s - ia} =$$

$$= \frac{s + ia}{s^2 + a^2}, \quad s > 0$$

Separate real and imaginary parts:

$$L[\cos at: t \rightarrow s] = \frac{s}{s^2 + a^2}, \quad L[\sin at: t \rightarrow s] = \frac{a}{s^2 + a^2}$$

Some General Properties

► **Linearity.**

If $\bar{f}(s) = L[f(t): t \rightarrow s]$ and $\bar{g}(s) = L[g(t): t \rightarrow s]$ exist then $L[af(t) + bg(t): t \rightarrow s]$ exists for all constants a and b and

$$L[af(t) + bg(t): t \rightarrow s] = a\bar{f}(s) + b\bar{g}(s)$$

Some General Properties

► **Linearity.**

If $\bar{f}(s) = L[f(t): t \rightarrow s]$ and $\bar{g}(s) = L[g(t): t \rightarrow s]$ exist then $L[af(t) + bg(t): t \rightarrow s]$ exists for all constants a and b and

$$L[af(t) + bg(t): t \rightarrow s] = a\bar{f}(s) + b\bar{g}(s)$$

► **First shifting property.**

Suppose $\bar{f}(s) = L[f(t): t \rightarrow s]$ exists and that a is a constant then $L[e^{at}f(t): t \rightarrow s]$ exists and

$$L[e^{at}f(t): t \rightarrow s] = \bar{f}(s - a)$$

or

$$L^{-1}[\bar{f}(s - a): s \rightarrow t] = e^{at}f(t)$$

Laplace Transforms of Selected Functions

$$f(t) = L^{-1} [\bar{f}(s): s \rightarrow t] \quad \bar{f}(s) = L[f(t): t \rightarrow s]$$

$$1 \qquad \frac{1}{s} \qquad \operatorname{Re}(s) > 0$$

$$e^{at} \qquad \frac{1}{s-a} \qquad \operatorname{Re}(s) > 0$$

$$t^n, n \in \mathbb{Z}_+ \qquad \frac{n!}{s^{n+1}} \qquad \operatorname{Re}(s) > 0$$

$$t^p, p > -1 \qquad \frac{\Gamma(p+1)}{s^{p+1}} \qquad \operatorname{Re}(s) > 0$$

$$\cos at \qquad \frac{s}{s^2 + a^2} \qquad \operatorname{Re}(s) > 0$$

$$\sin at \qquad \frac{a}{s^2 + a^2} \qquad \operatorname{Re}(s) > 0$$

Laplace Transforms of Selected Functions

$$f(t) = L^{-1} [\bar{f}(s): s \rightarrow t] \quad \bar{f}(s) = L [f(t): t \rightarrow s]$$
$$\cosh at \qquad \frac{s}{s^2 - a^2} \qquad \operatorname{Re}(s) > |a|$$

$$\sinh at \qquad \frac{a}{s^2 - a^2} \qquad \operatorname{Re}(s) > |a|$$

$$e^{at} \cos bt \qquad \frac{s - a}{(s - a)^2 + b^2} \qquad \operatorname{Re}(s) > a$$

$$e^{at} \sin bt \qquad \frac{b}{(s - a)^2 + b^2} \qquad \operatorname{Re}(s) > a$$

$$t^n e^{at}, \quad n \in \mathbb{Z}_+ \qquad \frac{n!}{(s - a)^{n+1}} \qquad \operatorname{Re}(s) > a$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] =$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- ▶ The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- ▶ But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow f(t) = 1, g(t) = t$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- ▶ The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- ▶ But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow f(t) = 1, g(t) = t$$

Therefore, $f(t)g(t) = t$ while

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- ▶ The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- ▶ But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow f(t) = 1, g(t) = t$$

Therefore, $f(t)g(t) = t$ while

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] =$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow f(t) = 1, g(t) = t$$

Therefore, $f(t)g(t) = t$ while

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] = L^{-1} [1/s^3: s \rightarrow t] =$$

Convolution

Let $\bar{f}(s) = L[f(t): t \rightarrow s]$, $\bar{g}(s) = L[g(t): t \rightarrow s]$

be Laplace transforms of $f(t)$ and $g(t)$.

- The linearity of the Laplace transform operator guarantees:

$$L^{-1} [\bar{f}(s) + \bar{g}(s): s \rightarrow t] =$$

$$= L^{-1} [\bar{f}(s): s \rightarrow t] + L^{-1} [\bar{g}(s): s \rightarrow t] = f(t) + g(t)$$

- But the inverse Laplace transform of a product **does not equal** the product of the inverse Laplace transforms of the factors:

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] \neq f(t)g(t)$$

For example,

$$\bar{f}(s) = \frac{1}{s}, \bar{g}(s) = \frac{1}{s^2} \Rightarrow f(t) = 1, g(t) = t$$

Therefore, $f(t)g(t) = t$ while

$$L^{-1} [\bar{f}(s)\bar{g}(s): s \rightarrow t] = L^{-1} [1/s^3: s \rightarrow t] = t^2/2 \neq f(t)g(t) = t$$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

$$L[(f \star g)(t): t \rightarrow s] =$$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

$$L[(f \star g)(t): t \rightarrow s] = \int_0^\infty \left(\int_0^t f(t-u)g(u) du \right) e^{-st} dt$$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

$$\begin{aligned} L[(f \star g)(t): t \rightarrow s] &= \int_0^\infty \left(\int_0^t f(t-u)g(u) du \right) e^{-st} dt \\ &= \int_0^\infty g(u) \left(\int_u^\infty f(t-u)e^{-st} dt \right) du \end{aligned}$$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

$$\begin{aligned} L[(f \star g)(t): t \rightarrow s] &= \int_0^\infty \left(\int_0^t f(t-u)g(u) du \right) e^{-st} dt \\ &= \int_0^\infty g(u) \left(\int_u^\infty f(t-u)e^{-st} dt \right) du \\ &= \int_0^\infty g(u) \left(\int_0^\infty f(w)e^{-s(w+u)} dw \right) du = \end{aligned}$$

Convolution

Given two functions $f(t)$ and $g(t)$, the **convolution** $f \star g$ of f and g is defined by the integral

$$(f \star g)(t) = \int_0^t f(t-u)g(u) du$$

whenever this integral exists.

- ▶ $(f \star g)(t) = (g \star f)(t)$ **verify by simple change of variable**
- ▶ $L[(f \star g)(t): t \rightarrow s] = \bar{f}(s)\bar{g}(s)$

$$\begin{aligned} L[(f \star g)(t): t \rightarrow s] &= \int_0^\infty \left(\int_0^t f(t-u)g(u) du \right) e^{-st} dt \\ &= \int_0^\infty g(u) \left(\int_u^\infty f(t-u)e^{-st} dt \right) du \\ &= \int_0^\infty g(u) \left(\int_0^\infty f(w)e^{-s(w+u)} dw \right) du = \bar{f}(s)\bar{g}(s) \end{aligned}$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f(\bar{s})\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f(\bar{s})\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] =$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f(\bar{s})\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] = L^{-1} \left[\frac{s}{s^2 + 1} \frac{1}{s^2 + 1} : s \rightarrow t \right]$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f(\bar{s})\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] = L^{-1} \left[\frac{s}{s^2 + 1} \frac{1}{s^2 + 1} : s \rightarrow t \right] = (\sin \star \cos)(t)$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f(\bar{s})\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] = L^{-1} \left[\frac{s}{s^2 + 1} \frac{1}{s^2 + 1} : s \rightarrow t \right] = (\sin \star \cos)(t)$$

$$(\sin \star \cos)(t) = \int_0^t \sin(t - u) \cos u \, du$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f\bar{(s)}\bar{g}(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] = L^{-1} \left[\frac{s}{s^2 + 1} \frac{1}{s^2 + 1} : s \rightarrow t \right] = (\sin \star \cos)(t)$$

$$(\sin \star \cos)(t) = \int_0^t \sin(t - u) \cos u \, du$$

$$= \sin t \int_0^t \cos^2 u \, du - \cos t \int_0^t \sin u \cos u \, du =$$

Examples

Example. Compute $L^{-1} [s/(s^2 + 1)^2: s \rightarrow t]$.

The convolution property

$$L^{-1} [(f\bar{g})(s): s \rightarrow t] = (f \star g)(t)$$

asserts that

$$L^{-1} \left[\frac{s}{(s^2 + 1)^2} : s \rightarrow t \right] = L^{-1} \left[\frac{s}{s^2 + 1} \frac{1}{s^2 + 1} : s \rightarrow t \right] = (\sin \star \cos)(t)$$

$$(\sin \star \cos)(t) = \int_0^t \sin(t - u) \cos u \, du$$

$$= \sin t \int_0^t \cos^2 u \, du - \cos t \int_0^t \sin u \cos u \, du = \frac{t \sin t}{2}$$

You can check the result $L[(t/2) \sin t: t \rightarrow s]$ by direct calculation as well

Examples

Example. Show that the solution of the integral equation

$$f(t) + \int_0^t e^u f(t-u) du = g(t)$$

can be written in the form

$$f(t) = g(t) - \int_0^t g(u) du.$$

Examples

Example. Show that the solution of the integral equation

$$f(t) + \int_0^t e^u f(t-u) du = g(t)$$

can be written in the form

$$f(t) = g(t) - \int_0^t g(u) du.$$

- ▶ Let $\bar{f}(s)$ be the Laplace transform of $f(t)$. Take Laplace transforms of the integral equation

$$\bar{f} + L[(f \star e)(t): t \rightarrow s] = \bar{g}$$

Examples

Example. Show that the solution of the integral equation

$$f(t) + \int_0^t e^u f(t-u) du = g(t)$$

can be written in the form

$$f(t) = g(t) - \int_0^t g(u) du.$$

- ▶ Let $\bar{f}(s)$ be the Laplace transform of $f(t)$. Take Laplace transforms of the integral equation

$$\bar{f} + L[(f \star e)(t): t \rightarrow s] = \bar{g}$$

- ▶ and use the property of the Laplace transform of a convolution

$$\bar{f} + \bar{f}L[e^t: t \rightarrow s] = \bar{g} \Rightarrow$$

Examples

Example. Show that the solution of the integral equation

$$f(t) + \int_0^t e^u f(t-u) du = g(t)$$

can be written in the form

$$f(t) = g(t) - \int_0^t g(u) du.$$

- ▶ Let $\bar{f}(s)$ be the Laplace transform of $f(t)$. Take Laplace transforms of the integral equation

$$\bar{f} + L[(f \star e)(t): t \rightarrow s] = \bar{g}$$

- ▶ and use the property of the Laplace transform of a convolution

$$\bar{f} + \bar{f} L[e^t: t \rightarrow s] = \bar{g} \Rightarrow \bar{f} = \bar{g} - \frac{\bar{g}}{s}$$

Examples

Example. Show that the solution of the integral equation

$$f(t) + \int_0^t e^u f(t-u) du = g(t)$$

can be written in the form

$$f(t) = g(t) - \int_0^t g(u) du.$$

- ▶ Let $\bar{f}(s)$ be the Laplace transform of $f(t)$. Take Laplace transforms of the integral equation

$$\bar{f} + L[(f \star e)(t): t \rightarrow s] = \bar{g}$$

- ▶ and use the property of the Laplace transform of a convolution

$$\bar{f} + \bar{f}L[e^t: t \rightarrow s] = \bar{g} \Rightarrow \bar{f} = \bar{g} - \frac{\bar{g}}{s}$$

$$f(t) = g(t) - L^{-1}\left[\frac{\bar{g}(s)}{s}: s \rightarrow t\right] = g(t) - \int_0^t g(x) dx$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$L \left[\dot{f}(t) : t \rightarrow s \right] =$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$L \left[\dot{f}(t) : t \rightarrow s \right] = \int_0^{\infty} \dot{f}(t) e^{-st} dt =$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = \end{aligned}$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = s \bar{f}(s) - f(0) \end{aligned}$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = s \bar{f}(s) - f(0) \end{aligned}$$

► Similarly,

$$L \left[\ddot{f}(t) : t \rightarrow s \right] = \int_0^{\infty} \ddot{f}(t) e^{-st} dt =$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = s \bar{f}(s) - f(0) \end{aligned}$$

► Similarly,

$$\begin{aligned} L \left[\ddot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \ddot{f}(t) e^{-st} dt = \left[\dot{f}(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} \dot{f}(t) e^{-st} dt = \end{aligned}$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = s \bar{f}(s) - f(0) \end{aligned}$$

► Similarly,

$$\begin{aligned} L \left[\ddot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \ddot{f}(t) e^{-st} dt = \left[\dot{f}(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} \dot{f}(t) e^{-st} dt = -\dot{f}(0) + s L \left[\dot{f}(t) : t \rightarrow s \right] \end{aligned}$$

Application to Initial-Value Problems

We assume that a function f has a Laplace transform and its derivative \dot{f} is piecewise continuous in the interval $(0, \infty)$. Then



$$\begin{aligned} L \left[\dot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \dot{f}(t) e^{-st} dt = \left[f(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} f(t) e^{-st} dt = s \bar{f}(s) - f(0) \end{aligned}$$

► Similarly,

$$\begin{aligned} L \left[\ddot{f}(t) : t \rightarrow s \right] &= \int_0^{\infty} \ddot{f}(t) e^{-st} dt = \left[\dot{f}(t) e^{-st} \right]_0^{\infty} + \\ &+ s \int_0^{\infty} \dot{f}(t) e^{-st} dt = -\dot{f}(0) + s L \left[\dot{f}(t) : t \rightarrow s \right] \\ &= s^2 \bar{f}(s) - s f(0) - \dot{f}(0) \end{aligned}$$

► Both results are valid for $s > 0$.

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Denote $\bar{y}(s) = L[y(t): t \rightarrow s]$ and

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Denote $\bar{y}(s) = L[y(t): t \rightarrow s]$ and apply the Laplace transform operation to the differential equation

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) - 4\bar{y}(s) = 0.$$

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Denote $\bar{y}(s) = L[y(t): t \rightarrow s]$ and apply the Laplace transform operation to the differential equation

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) - 4\bar{y}(s) = 0.$$

Solve the algebraic equation for $\bar{y}(s)$ to obtain

$$\bar{y}(s) = \frac{sy(0) + \dot{y}(0)}{s^2 - 4} =$$

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Denote $\bar{y}(s) = L[y(t): t \rightarrow s]$ and apply the Laplace transform operation to the differential equation

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) - 4\bar{y}(s) = 0.$$

Solve the algebraic equation for $\bar{y}(s)$ to obtain

$$\bar{y}(s) = \frac{sy(0) + \dot{y}(0)}{s^2 - 4} = \frac{s + 2}{s^2 - 4} = \frac{1}{s - 2}.$$

Examples

Example 1. Find the solution of the initial value problem

$$\ddot{y} - 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Denote $\bar{y}(s) = L[y(t): t \rightarrow s]$ and apply the Laplace transform operation to the differential equation

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) - 4\bar{y}(s) = 0.$$

Solve the algebraic equation for $\bar{y}(s)$ to obtain

$$\bar{y}(s) = \frac{sy(0) + \dot{y}(0)}{s^2 - 4} = \frac{s + 2}{s^2 - 4} = \frac{1}{s - 2}.$$

It immediately follows that

$$y(t) = L^{-1} \left[\frac{1}{s - 2} : s \rightarrow t \right] = e^{2t}.$$

Examples

Example 2. Find the solution of the initial value problem

$$\ddot{y} + 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Examples

Example 2. Find the solution of the initial value problem

$$\ddot{y} + 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$,

Examples

Example 2. Find the solution of the initial value problem

$$\ddot{y} + 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$, then the usual procedure shows that $\bar{y}(s)$ satisfies

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) + 4\bar{y}(s) = 0$$

and

Examples

Example 2. Find the solution of the initial value problem

$$\ddot{y} + 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$, then the usual procedure shows that $\bar{y}(s)$ satisfies

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) + 4\bar{y}(s) = 0$$

and therefore,

$$\bar{y}(s) = \frac{s+2}{s^2+4} = \frac{s}{s^2+4} + \frac{2}{s^2+4}.$$

Examples

Example 2. Find the solution of the initial value problem

$$\ddot{y} + 4y = 0, \quad y(0) = 1, \quad \dot{y}(0) = 2.$$

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$, then the usual procedure shows that $\bar{y}(s)$ satisfies

$$s^2 \bar{y}(s) - sy(0) - \dot{y}(0) + 4\bar{y}(s) = 0$$

and therefore,

$$\bar{y}(s) = \frac{s+2}{s^2+4} = \frac{s}{s^2+4} + \frac{2}{s^2+4}.$$

Thus

$$y(t) = \cos 2t + \sin 2t$$

As you can see, the method as useful as far we can take inverse Laplace transforms easily.

Example: electrical circuit

Example 3. Consider an electrical circuit with

- ▶ an inductor of inductance $L = 10^3$ Henry,
- ▶ a resistor of resistance $R = 6000\text{ Ohm}$,
- ▶ an uncharged capacitor of capacitance $C = (1/9)10^3$ Farad and
- ▶ a battery of electromotive force $U(t) = 250 \cos t$ Volt

which are connected in a serial arrangement. The circuit is activated at $t = 0$. Find the charge on the capacitor and the electrical current flowing in the circuit for $t \geq 0$.

Example: electrical circuit

Example 3. Consider an electrical circuit with

- ▶ an inductor of inductance $L = 10^3$ Henry,
- ▶ a resistor of resistance $R = 6000\text{ Ohm}$,
- ▶ an uncharged capacitor of capacitance $C = (1/9)10^3$ Farad and
- ▶ a battery of electromotive force $U(t) = 250 \cos t$ Volt

which are connected in a serial arrangement. The circuit is activated at $t = 0$. Find the charge on the capacitor and the electrical current flowing in the circuit for $t \geq 0$.

Apply Kirchhoff's Second Law (the sum of all voltage drops around a closed circuit is zero) to obtain the differential equation w.r.t the electrical current I and the rate of change of charge Q

$$L\dot{I}(t) + RI(t) + \frac{Q}{C} = U(t)$$

Example: electrical circuit

With $\dot{Q}(t) = I(t)$, initial conditions $Q(0) = 0$ and $I(0) = 0$ and the given values for L , R , C , the IVP becomes

$$\ddot{Q}(t) + 6\dot{Q}(t) + 9Q(t) = \frac{1}{4} \cos t, \quad Q(0) = 0, \quad \dot{Q}(0) = 0.$$

Example: electrical circuit

With $\dot{Q}(t) = I(t)$, initial conditions $Q(0) = 0$ and $I(0) = 0$ and the given values for L , R , C , the IVP becomes

$$\ddot{Q}(t) + 6\dot{Q}(t) + 9Q(t) = \frac{1}{4} \cos t, \quad Q(0) = 0, \quad \dot{Q}(0) = 0.$$

Denote $\bar{Q}(s) = L[Q(t): t \rightarrow s]$ and apply the Laplace transform operation to the equation:

Example: electrical circuit

With $\dot{Q}(t) = I(t)$, initial conditions $Q(0) = 0$ and $I(0) = 0$ and the given values for L , R , C , the IVP becomes

$$\ddot{Q}(t) + 6\dot{Q}(t) + 9Q(t) = \frac{1}{4} \cos t, \quad Q(0) = 0, \quad \dot{Q}(0) = 0.$$

Denote $\bar{Q}(s) = L[Q(t): t \rightarrow s]$ and apply the Laplace transform operation to the equation:

$$s^2 \bar{Q}(s) - sQ(0) - \dot{Q}(0) + 6(s\bar{Q}(s) - Q(0)) + 9\bar{Q}(s) = \frac{1}{4} \frac{s}{s^2 + 1}$$

Example: electrical circuit

With $\dot{Q}(t) = I(t)$, initial conditions $Q(0) = 0$ and $I(0) = 0$ and the given values for L, R, C , the IVP becomes

$$\ddot{Q}(t) + 6\dot{Q}(t) + 9Q(t) = \frac{1}{4} \cos t, \quad Q(0) = 0, \quad \dot{Q}(0) = 0.$$

Denote $\bar{Q}(s) = L[Q(t): t \rightarrow s]$ and apply the Laplace transform operation to the equation:

$$s^2 \bar{Q}(s) - sQ(0) - \dot{Q}(0) + 6(s\bar{Q}(s) - Q(0)) + 9\bar{Q}(s) = \frac{1}{4} \frac{s}{s^2 + 1}$$

Express $\bar{Q}(s)$ and apply partial fraction decomposition:

Example: electrical circuit

With $\dot{Q}(t) = I(t)$, initial conditions $Q(0) = 0$ and $I(0) = 0$ and the given values for L , R , C , the IVP becomes

$$\ddot{Q}(t) + 6\dot{Q}(t) + 9Q(t) = \frac{1}{4} \cos t, \quad Q(0) = 0, \quad \dot{Q}(0) = 0.$$

Denote $\bar{Q}(s) = L[Q(t): t \rightarrow s]$ and apply the Laplace transform operation to the equation:

$$s^2 \bar{Q}(s) - sQ(0) - \dot{Q}(0) + 6(s\bar{Q}(s) - Q(0)) + 9\bar{Q}(s) = \frac{1}{4} \frac{s}{s^2 + 1}$$

Express $\bar{Q}(s)$ and apply partial fraction decomposition:

$$\begin{aligned} \bar{Q}(s) &= \frac{s}{4(s^2 + 1)(s^2 + 6s + 9)} = \\ &= \frac{1}{200} \left(\frac{4s}{s^2 + 1} + \frac{3}{s^2 + 1} - \frac{4}{s + 3} - \underbrace{\frac{15}{(s + 3)^2}}_{\text{apply the shift formula}} \right) \end{aligned}$$

Example: electrical circuit

Extract the solution

$$Q(t) = \frac{1}{200}(4 \cos t + 3 \sin t - 4e^{-3t} - 15te^{-3t})$$

Example: electrical circuit

Extract the solution

$$Q(t) = \frac{1}{200}(4 \cos t + 3 \sin t - 4e^{-3t} - 15te^{-3t})$$

The current $I(t)$ follows immediately from $I(t) = \dot{Q}(t)$ and is given by

$$I(t) = \frac{1}{200}(-4 \sin t + 3 \cos t - 3e^{-3t} + 45te^{-3t}).$$

Example: electrical circuit

Extract the solution

$$Q(t) = \frac{1}{200}(4 \cos t + 3 \sin t - 4e^{-3t} - 15te^{-3t})$$

The current $I(t)$ follows immediately from $I(t) = \dot{Q}(t)$ and is given by

$$I(t) = \frac{1}{200}(-4 \sin t + 3 \cos t - 3e^{-3t} + 45te^{-3t}).$$

The long-term behavior of the solution is described by the result

$$\lim_{t \rightarrow \infty} \left[Q(t) - \frac{1}{200}(4 \cos t + 3 \sin t) \right] = 0.$$

Example: electrical circuit

Extract the solution

$$Q(t) = \frac{1}{200}(4 \cos t + 3 \sin t - 4e^{-3t} - 15te^{-3t})$$

The current $I(t)$ follows immediately from $I(t) = \dot{Q}(t)$ and is given by

$$I(t) = \frac{1}{200}(-4 \sin t + 3 \cos t - 3e^{-3t} + 45te^{-3t}).$$

The long-term behavior of the solution is described by the result

$$\lim_{t \rightarrow \infty} \left[Q(t) - \frac{1}{200}(4 \cos t + 3 \sin t) \right] = 0.$$

The solution for large time is dominated by the inhomogeneous term in the differential equation, while the contributions from the complementary function are damped out.

Example: electrical circuit

Extract the solution

$$Q(t) = \frac{1}{200}(4 \cos t + 3 \sin t - 4e^{-3t} - 15te^{-3t})$$

The current $I(t)$ follows immediately from $I(t) = \dot{Q}(t)$ and is given by

$$I(t) = \frac{1}{200}(-4 \sin t + 3 \cos t - 3e^{-3t} + 45te^{-3t}).$$

The long-term behavior of the solution is described by the result

$$\lim_{t \rightarrow \infty} \left[Q(t) - \frac{1}{200}(4 \cos t + 3 \sin t) \right] = 0.$$

The solution for large time is dominated by the inhomogeneous term in the differential equation, while the contributions from the complementary function are damped out.

In this example, the forcing function $U(t) = 250 \cos t$ is a continuous function of time. In many applications, such a function is a **step function** or an **impulse function**.

The Unit Step Function

The unit step function (or Heaviside function) is defined by

$$H(t - a) = \begin{cases} 0 & t < a \\ 1 & t \geq a. \end{cases}$$

The Unit Step Function

The **unit step function** (or **Heaviside function**) is defined by

$$H(t - a) = \begin{cases} 0 & t < a \\ 1 & t \geq a. \end{cases}$$



The unit step function can be used to construct more complicated step functions, for example $f(t) = H(t - a) - H(t - b)$.

The Unit Step Function

The **unit step function** (or **Heaviside function**) is defined by

$$H(t - a) = \begin{cases} 0 & t < a \\ 1 & t \geq a. \end{cases}$$

The unit step function can be used to construct more complicated step functions, for example $f(t) = H(t - a) - H(t - b)$.



The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt =$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt =$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

Example. Compute $L^{-1} \left[\frac{(1-e^{-\pi s/2})}{1+s^2} : s \rightarrow t \right]$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

Example. Compute $L^{-1}\left[\frac{(1-e^{-\pi s/2})}{1+s^2}: s \rightarrow t\right]$

$$L^{-1}\left[\frac{1-e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right] = L^{-1}\left[\frac{1}{1+s^2}: s \rightarrow t\right] - L^{-1}\left[\frac{e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right]$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

Example. Compute $L^{-1}\left[\frac{(1-e^{-\pi s/2})}{1+s^2}: s \rightarrow t\right]$

$$\begin{aligned} L^{-1}\left[\frac{1-e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right] &= L^{-1}\left[\frac{1}{1+s^2}: s \rightarrow t\right] - L^{-1}\left[\frac{e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right] \\ &= \sin t - \end{aligned}$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

Example. Compute $L^{-1}\left[\frac{(1-e^{-\pi s/2})}{1+s^2}: s \rightarrow t\right]$

$$L^{-1}\left[\frac{1-e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right] = L^{-1}\left[\frac{1}{1+s^2}: s \rightarrow t\right] - L^{-1}\left[\frac{e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right]$$

$$= \sin t - H(t - \pi/2) \sin(t - \pi/2) =$$

The Unit Step Function

The Laplace transform of the unit step function is

$$L[H(t-a): t \rightarrow s] = \int_0^{\infty} e^{-st} H(t-a) dt = \int_a^{\infty} e^{-st} dt = \frac{e^{-as}}{s}, s > 0$$

The second shifting property

$$L[f(t-a)H(t-a): t \rightarrow s] = e^{-as}\bar{f}(s)$$

Example. Compute $L^{-1}\left[\frac{(1-e^{-\pi s/2})}{1+s^2}: s \rightarrow t\right]$

$$L^{-1}\left[\frac{1-e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right] = L^{-1}\left[\frac{1}{1+s^2}: s \rightarrow t\right] - L^{-1}\left[\frac{e^{-\pi s/2}}{1+s^2}: s \rightarrow t\right]$$

$$= \sin t - H(t - \pi/2) \sin(t - \pi/2) = \sin t + H(t - \pi/2) \cos t$$

The Unit Step Function

Example. Solve the initial value problem

$$\ddot{y} + y = \begin{cases} 1 & 0 \leq t < 1, \\ 0 & t \geq 1, \end{cases}, \quad y(0) = 0, \dot{y}(0) = 0.$$

Such a problem may appear when an undamped oscillator is disturbed from rest in its equilibrium state by a constant external driving force which is active for a finite period only.

The Unit Step Function

Example. Solve the initial value problem

$$\ddot{y} + y = \begin{cases} 1 & 0 \leq t < 1, \\ 0 & t \geq 1, \end{cases}, \quad y(0) = 0, \dot{y}(0) = 0.$$

Such a problem may appear when an undamped oscillator is disturbed from rest in its equilibrium state by a constant external driving force which is active for a finite period only.

Represent the driving force as

$$f(t) = H(t) - H(t - 1)$$

and let $\bar{y}(s) = L[y(t): t \rightarrow s]$.

The Unit Step Function

Example. Solve the initial value problem

$$\ddot{y} + y = \begin{cases} 1 & 0 \leq t < 1, \\ 0 & t \geq 1, \end{cases}, \quad y(0) = 0, \dot{y}(0) = 0.$$

Such a problem may appear when an undamped oscillator is disturbed from rest in its equilibrium state by a constant external driving force which is active for a finite period only.

Represent the driving force as

$$f(t) = H(t) - H(t - 1)$$

and let $\bar{y}(s) = L[y(t): t \rightarrow s]$.

Take the Laplace transform of the equation to obtain

$$s^2 \bar{y}(s) - s\dot{y}(0) - y(0) + \bar{y}(s) = \frac{1}{s} - \frac{e^{-s}}{s}$$

The Unit Step Function

Then

$$\bar{y}(s) = \frac{1 - e^{-s}}{s(s^2 + 1)} =$$

The Unit Step Function

Then

$$\bar{y}(s) = \frac{1 - e^{-s}}{s(s^2 + 1)} = \frac{1}{s} - \frac{s}{s^2 + 1} - \underbrace{\frac{e^{-s}}{s} + \frac{se^{-s}}{s^2 + 1}}_{\text{apply the 2nd shifting pr.}}$$

The Unit Step Function

Then

$$\bar{y}(s) = \frac{1 - e^{-s}}{s(s^2 + 1)} = \frac{1}{s} - \frac{s}{s^2 + 1} - \underbrace{\frac{e^{-s}}{s} + \frac{se^{-s}}{s^2 + 1}}_{\text{apply the 2nd shifting pr.}}$$

The solution of the IVP is

$$y(t) = 1 - \cos t - H(t - 1) + H(t - 1) \cos(t - 1)$$

The Unit Step Function

Then

$$\bar{y}(s) = \frac{1 - e^{-s}}{s(s^2 + 1)} = \frac{1}{s} - \frac{s}{s^2 + 1} - \underbrace{\frac{e^{-s}}{s} + \frac{se^{-s}}{s^2 + 1}}_{\text{apply the 2nd shifting pr.}}$$

The solution of the IVP is

$$y(t) = 1 - \cos t - H(t - 1) + H(t - 1) \cos(t - 1)$$

$$y(t) = \begin{cases} 1 - \cos t & 0 \leq t < 1, \\ \cos(t - 1) - \cos t & t \geq 1, \end{cases}$$

The Unit Step Function

Then

$$\bar{y}(s) = \frac{1 - e^{-s}}{s(s^2 + 1)} = \frac{1}{s} - \frac{s}{s^2 + 1} - \underbrace{\frac{e^{-s}}{s} + \frac{se^{-s}}{s^2 + 1}}_{\text{apply the 2nd shifting pr.}}$$

The solution of the IVP is

$$y(t) = 1 - \cos t - H(t - 1) + H(t - 1) \cos(t - 1)$$

$$y(t) = \begin{cases} 1 - \cos t & 0 \leq t < 1, \\ \cos(t - 1) - \cos t & t \geq 1, \end{cases}$$

The solution is continuous at $t = 1$ even though the driving force $f(t)$ is not.

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The **unit impulse function** $\delta(t)$ is defined through the integral relation

$$\int_{-\infty}^{+\infty} \delta(t - a) f(t) dt = f(a)$$

for any integrable function $f(t)$.

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The **unit impulse function** $\delta(t)$ is defined through the integral relation

$$\int_{-\infty}^{+\infty} \delta(t - a) f(t) dt = f(a)$$

for any integrable function $f(t)$.

The delta function possesses good analytical properties because the delta function may be thought of as a limit of a sequence of classical functions.

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The **unit impulse function** $\delta(t)$ is defined through the integral relation

$$\int_{-\infty}^{+\infty} \delta(t - a) f(t) dt = f(a)$$

for any integrable function $f(t)$.

The delta function possesses good analytical properties because the delta function may be thought of as a limit of a sequence of classical functions.

Sequence 1: Let $\delta_\varepsilon(t - a) = \begin{cases} \frac{1}{2\varepsilon} & |t - a| < \varepsilon, \\ 0 & |t - a| \geq \varepsilon. \end{cases}$

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The **unit impulse function** $\delta(t)$ is defined through the integral relation

$$\int_{-\infty}^{+\infty} \delta(t-a)f(t) dt = f(a)$$

for any integrable function $f(t)$.

The delta function possesses good analytical properties because the delta function may be thought of as a limit of a sequence of classical functions.

Sequence 1: Let $\delta_{\varepsilon}(t-a) = \begin{cases} \frac{1}{2\varepsilon} & |t-a| < \varepsilon, \\ 0 & |t-a| \geq \varepsilon. \end{cases}$

$$\int_{-\infty}^{+\infty} \delta_{\varepsilon}(t-a)f(t) dt = \frac{1}{2\varepsilon} \int_{a-\varepsilon}^{a+\varepsilon} f(t) dt =$$

The Unit Impulse Function

Consider the **unit impulse function** $\delta(t)$ (or delta or Dirac delta function) which is not an ordinary function but belongs to the class of so-called **generalized functions** or **distributions**.

The **unit impulse function** $\delta(t)$ is defined through the integral relation

$$\int_{-\infty}^{+\infty} \delta(t-a)f(t) dt = f(a)$$

for any integrable function $f(t)$.

The delta function possesses good analytical properties because the delta function may be thought of as a limit of a sequence of classical functions.

Sequence 1: Let $\delta_\varepsilon(t-a) = \begin{cases} \frac{1}{2\varepsilon} & |t-a| < \varepsilon, \\ 0 & |t-a| \geq \varepsilon. \end{cases}$

$$\int_{-\infty}^{+\infty} \delta_\varepsilon(t-a)f(t) dt = \frac{1}{2\varepsilon} \int_{a-\varepsilon}^{a+\varepsilon} f(t) dt = f(a+\theta\varepsilon), |\theta| < 1$$

The Unit Impulse Function

We used the fact that f is integrable and applied the integral mean value theorem above.

The Unit Impulse Function

We used the fact that f is integrable and applied the integral mean value theorem above. Therefore,

$$\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{+\infty} \delta_{\varepsilon}(t - a) f(t) dt = f(a)$$

for any real integrable function f . Therefore,

$$\delta(t - a) = \lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon}(t - a)$$

The Unit Impulse Function

We used the fact that f is integrable and applied the integral mean value theorem above. Therefore,

$$\lim_{\varepsilon \rightarrow 0} \int_{-\infty}^{+\infty} \delta_{\varepsilon}(t - a) f(t) dt = f(a)$$


for any real integrable function f . Therefore,

$$\delta(t - a) = \lim_{\varepsilon \rightarrow 0} \delta_{\varepsilon}(t - a)$$

Sequence 2: Another popular description of $\delta(t)$ ($a = 0$) is the limit of the sequence

$$\delta_n(t) = \lim_{n \rightarrow \infty} \frac{n}{\sqrt{\pi}} e^{-n^2 t^2}$$

The Unit Impulse Function



The graph of the function $(n/\sqrt{\pi})e^{-n^2 t^2}$ for $n = 1, 5, 10$.

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

- ▶ The unit step function is related to the unit impulse function

$$H(t - a) = \int_{-\infty}^t \delta(u - a) du \Rightarrow \frac{d}{dt} H(t - a) = \delta(t - a)$$

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

- ▶ The unit step function is related to the unit impulse function

$$H(t - a) = \int_{-\infty}^t \delta(u - a) du \Rightarrow \frac{d}{dt} H(t - a) = \delta(t - a)$$

- ▶ The Laplace transform of $\delta(t - a)$ is

$$L[\delta(t - a): t \rightarrow s] =$$

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

- ▶ The unit step function is related to the unit impulse function

$$H(t - a) = \int_{-\infty}^t \delta(u - a) du \Rightarrow \frac{d}{dt} H(t - a) = \delta(t - a)$$

- ▶ The Laplace transform of $\delta(t - a)$ is

$$L[\delta(t - a): t \rightarrow s] = \int_0^{\infty} \delta(t - a) e^{-st} dt$$

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

- ▶ The unit step function is related to the unit impulse function

$$H(t - a) = \int_{-\infty}^t \delta(u - a) du \Rightarrow \frac{d}{dt} H(t - a) = \delta(t - a)$$

- ▶ The Laplace transform of $\delta(t - a)$ is

$$\begin{aligned} L[\delta(t - a): t \rightarrow s] &= \int_0^{\infty} \delta(t - a) e^{-st} dt \\ &= \int_{-\infty}^{\infty} \delta(t - a) e^{-st} dt = \end{aligned}$$

The Unit Impulse Function

In the strict mathematical sense neither of these limits exists, but they provide useful conceptual ways towards an understanding of the delta function.

- ▶ Letting $f(t) = 1$ in the definition of the delta function,

$$\int_{-\infty}^{+\infty} \delta(t - a) dt = 1$$

For this reason $\delta(t - a)$ is called the **unit impulse function**.

- ▶ The unit step function is related to the unit impulse function

$$H(t - a) = \int_{-\infty}^t \delta(u - a) du \Rightarrow \frac{d}{dt} H(t - a) = \delta(t - a)$$

- ▶ The Laplace transform of $\delta(t - a)$ is


$$\begin{aligned} L[\delta(t - a): t \rightarrow s] &= \int_0^{\infty} \delta(t - a) e^{-st} dt \\ &= \int_{-\infty}^{\infty} \delta(t - a) e^{-st} dt = e^{-as}, \quad a > 0 \end{aligned}$$

The Unit Impulse Function

Consider an elastic ball of mass m moving at velocity v_0 toward a rigid wall.


The Unit Impulse Function

Consider an elastic ball of mass m moving at velocity v_0 toward a rigid wall.



The Unit Impulse Function


Consider an elastic ball of mass m moving at velocity v_0 toward a rigid wall.



- At time t_0 , the ball collides with the wall, the wall exerts a force $f(t)$ on the ball over a short period of time ε .

The Unit Impulse Function

Consider an elastic ball of mass m moving at velocity v_0 toward a rigid wall.



- ▶ At time t_0 , the ball collides with the wall, the wall exerts a force $f(t)$ on the ball over a short period of time ε .
- ▶ During this time, the ball is in contact with the wall and the velocity of the ball reduces from v_0 to 0 and then changes its direction, finally leaving the wall with velocity $-v_1$.

The Unit Impulse Function


- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall.

The Unit Impulse Function

- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall. It is opposite to the direction of the initial velocity $v_0 \Rightarrow f(t)$ is negative.


The Unit Impulse Function

- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall. It is opposite to the direction of the initial velocity $v_0 \Rightarrow f(t)$ is negative.



The Unit Impulse Function


- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall. It is opposite to the direction of the initial velocity $v_0 \Rightarrow f(t)$ is negative.



- ▶ The area under the force curve is called the **impulse** /

The Unit Impulse Function

- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall. It is opposite to the direction of the initial velocity $v_0 \Rightarrow f(t)$ is negative.




- ▶ The area under the force curve is called the **impulse** I

$$I = \int_{t_0}^{t_0 + \epsilon} f(t) dt$$

The Unit Impulse Function

- ▶ The force $f(t)$ depends on the contact between the elastic ball and the rigid wall. It is opposite to the direction of the initial velocity $v_0 \Rightarrow f(t)$ is negative.



- ▶ The area under the force curve is called the **impulse** I

$$I = \int_{t_0}^{t_0 + \epsilon} f(t) dt$$

- ▶ The **Impulse-Momentum Principle**: the change in momentum of mass m is equal to the total impulse on m :
 $m(-v_1) - mv_0 = I$.

The Unit Impulse Function

- ▶ A mathematical idealization:

The Unit Impulse Function

- ▶ A mathematical idealization: consider an impulse function $f(t)$ over a time interval $t_0 < t < t_0 + \varepsilon$ with constant amplitude I/ε such that the area under the function $f(t)$ is I

The Unit Impulse Function

- ▶ A mathematical idealization: consider an impulse function $f(t)$ over a time interval $t_0 < t < t_0 + \varepsilon$ with constant amplitude I/ε such that the area under the function $f(t)$ is I

$$f(t) = \frac{I}{\varepsilon}(H(t - t_0) - H(t - (t_0 + \varepsilon)))$$

The Unit Impulse Function

- ▶ A mathematical idealization: consider an impulse function $f(t)$ over a time interval $t_0 < t < t_0 + \varepsilon$ with constant amplitude I/ε such that the area under the function $f(t)$ is I


$$f(t) = \frac{I}{\varepsilon}(H(t - t_0) - H(t - (t_0 + \varepsilon)))$$

- ▶ For $I = 1$, the limiting function

$$\lim_{\varepsilon \rightarrow 0} f(t) = \lim_{\varepsilon \rightarrow 0} \frac{H(t - t_0) - H(t - (t_0 + \varepsilon))}{\varepsilon}$$

is called the unit impulse function

The Unit Impulse Function



The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an external driving force of infinite magnitude acting for an infinitely short time such that unit impulse is imparted to the system at time $t = 1$.

The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an external driving force of infinite magnitude acting for an infinitely short time such that unit impulse is imparted to the system at time $t = 1$.

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$ and take the Laplace transform of the differential equation:

The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an external driving force of infinite magnitude acting for an infinitely short time such that unit impulse is imparted to the system at time $t = 1$.

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$ and take the Laplace transform of the differential equation:

$$s^2\bar{y}(s) - 2s - 3 + 2(s\bar{y}(s) - 2) + \bar{y}(s) = e^{-s}.$$

The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an external driving force of infinite magnitude acting for an infinitely short time such that unit impulse is imparted to the system at time $t = 1$.

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$ and take the Laplace transform of the differential equation:

$$s^2\bar{y}(s) - 2s - 3 + 2(s\bar{y}(s) - 2) + \bar{y}(s) = e^{-s}.$$

$$\Rightarrow \bar{y}(s) = \frac{2s + 7 + e^{-s}}{s^2 + 2s + 1} =$$

The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an **external driving force of infinite magnitude acting for an infinitely short time** such that unit impulse is imparted to the system at time $t = 1$.

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$ and take the Laplace transform of the differential equation:

$$s^2\bar{y}(s) - 2s - 3 + 2(s\bar{y}(s) - 2) + \bar{y}(s) = e^{-s}.$$

$$\Rightarrow \bar{y}(s) = \frac{2s + 7 + e^{-s}}{s^2 + 2s + 1} = \frac{2}{s + 1} + \frac{5}{(s + 1)^2} + \frac{e^{-s}}{(s + 1)^2}$$

after taking partial fractions.

The Unit Impulse Function: Examples

Solve the IVP

$$\ddot{y} + 2\dot{y} + y = \delta(t - 1), y(0) = 2, \dot{y}(0) = 3$$

that represents a damped oscillator with an **external driving force of infinite magnitude acting for an infinitely short time** such that unit impulse is imparted to the system at time $t = 1$.

Let $\bar{y}(s) = L[y(t): t \rightarrow s]$ and take the Laplace transform of the differential equation:

$$s^2\bar{y}(s) - 2s - 3 + 2(s\bar{y}(s) - 2) + \bar{y}(s) = e^{-s}.$$

$$\Rightarrow \bar{y}(s) = \frac{2s + 7 + e^{-s}}{s^2 + 2s + 1} = \frac{2}{s + 1} + \frac{5}{(s + 1)^2} + \frac{e^{-s}}{(s + 1)^2}$$

after taking partial fractions. Apply the first and the second shifting properties to obtain

$$y(t) = 2e^{-t} + 5te^{-t} + (t - 1)e^{-(t-1)}H(t - 1)$$

Laplace Transforms of Selected Functions

$$f(t) = L^{-1} [\bar{f}(s): s \rightarrow t] \quad \bar{f}(s) = L[f(t): t \rightarrow s]$$

$$H(t-a) \quad \frac{e^{-as}}{s} \quad \text{Re}(s) > 0$$

$$\frac{H(t-a)f(t-a)}{e^{at}f(t)} \quad \frac{e^{-as}\bar{f}(s)}{\bar{f}(s-a)}$$

$$\frac{f(at)}{\int_0^t f(t-w)g(w)dw} \quad \frac{\frac{1}{a}\bar{f}\left(\frac{s}{a}\right)}{\bar{f}(s)\bar{g}(s)} \quad a > 0$$

$$\delta(t-a) \quad e^{-as}$$

$$\frac{(-t)^n f(t)}{f^{(n)}(t)} \quad \frac{\bar{f}^{(n)}(s)}{s^n \bar{f}(s) - \sum_{j=0}^{n-1} s^{n-j-1} f^{(j)}(0)}$$

Periodic Functions

Let $f(t)$ be a real-valued function defined for $0 \leq t < \infty$ with periodicity T :

$$f(t) = f(t + kT), \quad k = 1, 2, \dots$$

Periodic Functions

Let $f(t)$ be a real-valued function defined for $0 \leq t < \infty$ with periodicity T :

$$f(t) = f(t + kT), \quad k = 1, 2, \dots$$

The Laplace transform of the periodic function f is

$$L[f(t): t \rightarrow s] = \bar{f}(s) = \int_0^{\infty} f(t)e^{-st} dt = \sum_{k=0}^{\infty} \int_{kT}^{(k+1)T} f(t)e^{-st} dt$$

Use the substitution $t = u + kT$

Periodic Functions

Let $f(t)$ be a real-valued function defined for $0 \leq t < \infty$ with periodicity T :

$$f(t) = f(t + kT), \quad k = 1, 2, \dots$$

The Laplace transform of the periodic function f is

$$L[f(t): t \rightarrow s] = \bar{f}(s) = \int_0^{\infty} f(t)e^{-st} dt = \sum_{k=0}^{\infty} \int_{kT}^{(k+1)T} f(t)e^{-st} dt$$

Use the substitution $t = u + kT$ and exploit the periodicity of f :

$$\bar{f}(s) = \sum_{k=0}^{\infty} \int_0^T f(u+kT)e^{-s(u+kT)} du =$$

Periodic Functions

Let $f(t)$ be a real-valued function defined for $0 \leq t < \infty$ with periodicity T :

$$f(t) = f(t + kT), \quad k = 1, 2, \dots$$

The Laplace transform of the periodic function f is

$$L[f(t): t \rightarrow s] = \bar{f}(s) = \int_0^{\infty} f(t)e^{-st} dt = \sum_{k=0}^{\infty} \int_{kT}^{(k+1)T} f(t)e^{-st} dt$$

Use the substitution $t = u + kT$ and exploit the periodicity of f :

$$\bar{f}(s) = \sum_{k=0}^{\infty} \int_0^T f(u+kT)e^{-s(u+kT)} du = \sum_{k=0}^{\infty} e^{-kTu} \int_0^T f(u)e^{-su} du$$

Periodic Functions

Let $f(t)$ be a real-valued function defined for $0 \leq t < \infty$ with periodicity T :

$$f(t) = f(t + kT), \quad k = 1, 2, \dots$$

The Laplace transform of the periodic function f is

$$L[f(t): t \rightarrow s] = \bar{f}(s) = \int_0^{\infty} f(t)e^{-st} dt = \sum_{k=0}^{\infty} \int_{kT}^{(k+1)T} f(t)e^{-st} dt$$

Use the substitution $t = u + kT$ and exploit the periodicity of f :

$$\bar{f}(s) = \sum_{k=0}^{\infty} \int_0^T f(u+kT)e^{-s(u+kT)} du = \sum_{k=0}^{\infty} e^{-kTs} \int_0^T f(u)e^{-su} du$$

$$\bar{f}(s) = \frac{1}{1 - e^{-sT}} \int_0^T f(t)e^{-st} dt$$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$.

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated

$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated

$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Since $\sin at$ has periodicity $T = 2\pi/a \Rightarrow \sin aT = 0, \cos aT = 1,$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated

$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Since $\sin at$ has periodicity $T = 2\pi/a \Rightarrow \sin aT = 0, \cos aT = 1$,

$$\int_0^T e^{-st} \sin at dt =$$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated

$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Since $\sin at$ has periodicity $T = 2\pi/a \Rightarrow \sin aT = 0, \cos aT = 1$,

$$\int_0^T e^{-st} \sin at dt = \left. \frac{-e^{-st}}{s} \sin at \right|_0^T + \frac{a}{s} \int_0^T e^{-st} \cos at dt$$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated

$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Since $\sin at$ has periodicity $T = 2\pi/a \Rightarrow \sin aT = 0, \cos aT = 1$,

$$\begin{aligned} \int_0^T e^{-st} \sin at dt &= \left. \frac{-e^{-st}}{s} \sin at \right|_0^T + \frac{a}{s} \int_0^T e^{-st} \cos at dt \\ &= \frac{a}{s} \left. \frac{-e^{-st}}{s} \cos at \right|_0^T - \frac{a^2}{s^2} \int_0^T e^{-st} \sin at dt \end{aligned}$$

Periodic Functions: Examples

To demonstrate the validity of the obtained formula, we shall use it to calculate the Laplace transform of $\sin at$. We have already calculated



$$L[\sin at: t \rightarrow s] = \frac{a}{a^2 + s^2}$$

Since $\sin at$ has periodicity $T = 2\pi/a \Rightarrow \sin aT = 0, \cos aT = 1$,

$$\begin{aligned}\int_0^T e^{-st} \sin at dt &= \left. \frac{-e^{-st}}{s} \sin at \right|_0^T + \frac{a}{s} \int_0^T e^{-st} \cos at dt \\&= \frac{a}{s} \left. \frac{-e^{-st}}{s} \cos at \right|_0^T - \frac{a^2}{s^2} \int_0^T e^{-st} \sin at dt \\&= \frac{a}{s^2} (1 - e^{-sT}) - \frac{a^2}{s^2} \int_0^T e^{-st} \sin at dt\end{aligned}$$



Common periodic waveforms

The periodic waveforms **saw tooth** and **square wave** of amplitude V and periods T and $2T$ are shown below



Common periodic waveforms

The periodic waveforms **saw tooth** and **square wave** of amplitude V and periods T and $2T$ are shown below




Since

$$\int_0^T f(t) e^{-st} dt =$$

Common periodic waveforms

The periodic waveforms **saw tooth** and **square wave** of amplitude V and periods T and $2T$ are shown below




Since

$$\int_0^T f(t)e^{-st}dt = \frac{V}{T} \int_0^T te^{-st}dt =$$

Common periodic waveforms

The periodic waveforms **saw tooth** and **square wave** of amplitude V and periods T and $2T$ are shown below



Since

$$\int_0^T f(t)e^{-st}dt = \frac{V}{T} \int_0^T te^{-st}dt = \frac{V}{Ts^2}(1 - e^{-sT} - sTe^{-sT}),$$

Common periodic waveforms

so the Laplace transform of the saw tooth waveform of amplitude V and period T is

$$\bar{f}(s) = \frac{V}{Ts^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

Common periodic waveforms

so the Laplace transform of the saw tooth waveform of amplitude V and period T is

$$\bar{f}(s) = \frac{V}{Ts^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

The Laplace transform of the square wave of amplitude V and period $2T$ is

$$\bar{f}(s) = \frac{V}{1 - e^{-2sT}} \int_0^T te^{-st} dt = \frac{V}{s} \frac{1}{1 + e^{-sT}}$$

Common periodic waveforms

so the Laplace transform of the saw tooth waveform of amplitude V and period T is

$$\bar{f}(s) = \frac{V}{Ts^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

The Laplace transform of the square wave of amplitude V and period $2T$ is

$$\bar{f}(s) = \frac{V}{1 - e^{-2sT}} \int_0^T te^{-st} dt = \frac{V}{s} \frac{1}{1 + e^{-sT}}$$

Other periodic pulses.

Meander function

$$f(t) = \begin{cases} V & t \in (0, T) \\ -V & t \in (T, 2T) \end{cases}$$

Full wave rectification

$$f(t) = \begin{cases} V \sin(\frac{\pi t}{T}) & t \in (0, T) \end{cases}$$

Triangular wave

$$f(t) = \begin{cases} tV/T & t \in (0, T) \\ \frac{(2T-t)V}{T} & t \in (T, 2T) \end{cases}$$

Half wave rectification

$$f(t) = \begin{cases} V \sin(\pi t/T) & t \in (0, T) \\ 0 & t \in (T, 2T) \end{cases}$$

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

The current $I(t)$ in the circuit is governed by the differential equation

$$L\dot{I} + RI = U_{\text{saw}}(T), \quad I(0) = 0.$$

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

The current $I(t)$ in the circuit is governed by the differential equation

$$L\dot{I} + RI = U_{\text{saw}}(T), \quad I(0) = 0.$$

Therefore, the Laplace transform $\bar{I}(s) = L[I(t): t \rightarrow s]$ satisfies the algebraic equation

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

The current $I(t)$ in the circuit is governed by the differential equation

$$L\dot{I} + RI = U_{\text{saw}}(T), \quad I(0) = 0.$$

Therefore, the Laplace transform $\bar{I}(s) = L[I(t): t \rightarrow s]$ satisfies the algebraic equation

$$(Ls + R)\bar{I} = \frac{V}{T} \frac{1}{s^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right).$$

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

The current $I(t)$ in the circuit is governed by the differential equation

$$L\dot{I} + RI = U_{\text{saw}}(T), \quad I(0) = 0.$$

Therefore, the Laplace transform $\bar{I}(s) = L[I(t): t \rightarrow s]$ satisfies the algebraic equation

$$(Ls + R)\bar{I} = \frac{V}{T} \frac{1}{s^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right).$$

Denote $a = R/L$. Then

Example

Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a saw tooth voltage of period T and amplitude V . Assume that no current is flowing initially.

The current $I(t)$ in the circuit is governed by the differential equation

$$L\dot{I} + RI = U_{\text{saw}}(T), \quad I(0) = 0.$$

Therefore, the Laplace transform $\bar{I}(s) = L[I(t): t \rightarrow s]$ satisfies the algebraic equation

$$(Ls + R)\bar{I} = \frac{V}{T} \frac{1}{s^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right).$$

Denote $a = R/L$. Then

$$\bar{I} = \frac{V}{RT} \frac{a}{s^2(s + a)} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right) = \frac{V}{RT} \frac{1}{s^2} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right) -$$

Example

$$-\frac{V}{RT} \frac{1}{s(s+a)} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

Example

$$-\frac{V}{RT} \frac{1}{s(s+a)} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

Note, that the first term in the last expression is a multiple of $L[U_{saw}(t) : t \rightarrow s]$ and hence,

Example

$$-\frac{V}{RT} \frac{1}{s(s+a)} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

Note, that the first term in the last expression is a multiple of $L[U_{saw}(t) : t \rightarrow s]$ and hence,

$$\bar{I} - \frac{1}{R} \bar{U}_{saw} = -\frac{VL}{R^2 T} \frac{a}{s(s+a)} + \frac{V}{R} \frac{1}{s+a} \frac{e^{-sT}}{1 - e^{-sT}}$$

Example

$$-\frac{V}{RT} \frac{1}{s(s+a)} \left(1 - \frac{sTe^{-sT}}{1 - e^{-sT}} \right)$$

Note, that the first term in the last expression is a multiple of $L[U_{saw}(t) : t \rightarrow s]$ and hence,

$$\begin{aligned} \bar{I} - \frac{1}{R} \bar{U}_{saw} &= -\frac{VL}{R^2 T} \frac{a}{s(s+a)} + \frac{V}{R} \frac{1}{s+a} \frac{e^{-sT}}{1 - e^{-sT}} \\ &= -\frac{VL}{R^2 T} \frac{a}{s(s+a)} + \frac{V}{R} \sum_{k=1}^{\infty} \frac{e^{-skT}}{s+a} \end{aligned}$$

Example

Using the inverse Laplace transforms

$$L^{-1} \left[\frac{a}{s(s+a)} : s \rightarrow t \right] = 1 - e^{-at}$$

Example

Using the inverse Laplace transforms

$$L^{-1} \left[\frac{a}{s(s+a)} : s \rightarrow t \right] = 1 - e^{-at}$$

$$L^{-1} \left[\frac{e^{-skT}}{(s+a)} : s \rightarrow t \right] = e^{-a(t-kT)} H(t - kT)$$

Example

Using the inverse Laplace transforms

$$L^{-1} \left[\frac{a}{s(s+a)} : s \rightarrow t \right] = 1 - e^{-at}$$

$$L^{-1} \left[\frac{e^{-skT}}{(s+a)} : s \rightarrow t \right] = e^{-a(t-kT)} H(t - kT)$$

we obtain that

$$I(t) = \frac{U_{saw}(t)}{R} - \frac{VL}{R^2 T} (1 - e^{-at}) + \frac{V}{R} \sum_{k=1}^{\infty} e^{-a(t-kT)} H(t - kT)$$

Exercises

1. Find the Laplace transform of $f(t) = e^{-t-1/2}$ and $f(t) = \cos(at + b)$.
2. Find the inverse Laplace transform of $\bar{f}(s) = (s - 2)/(s^2 - 2)$, $\bar{f}(s) = 3/(s^2 + 4s + 9)$.
3. Solve the convolution integral equations

$$A. \quad f(t) = 1 + \int_0^t f(u) \cos(t - u) du,$$

$$B. \quad \sin t - t = \int_0^t (t - u)^2 f(u) du.$$

4. Use Laplace transforms to solve the initial value problems

$$A. \quad \ddot{y}(t) - 5\dot{y}(t) + 6y(t) = 0, \quad y(0) = 2, \quad \dot{y}(0) = 1,$$

$$B. \quad \ddot{y}(t) - y(t) = te^{2t}, \quad y(0) = 0, \quad \dot{y}(0) = 1.$$

5. Using the definition of the Laplace transform, prove the second shifting property.

Exercises

6. Write $f(t)$ as a step function and calculate its Laplace transform

$$f(t) = \begin{cases} 1 & t < 0 \\ 3 & 1 \leq t < 7 \\ 5 & t \geq 7. \end{cases}$$

7. Use Laplace transforms to solve the initial value problem

$$4\ddot{y}(t) + 4\dot{y}(t) + 5y(t) = g(t) \quad y(0) = \dot{y}(0) = 0,$$

$$g(t) = \begin{cases} 4 & 0 \leq t < \pi \\ 0 & t \geq \pi \end{cases} \quad \text{and sketch the solution } y(t).$$

8. Solve the initial value problem

$$\ddot{y}(t) + 2\dot{y}(t) + 2y(t) = \delta(t - \pi) \quad y(0) = \dot{y}(0) = 0,$$

by using Laplace transforms and sketch the solution $y(t)$.

9. Determine the current in a series circuit containing an inductor of L Henry and a resistor of R Ohm when the circuit is driven by a square wave voltage of period $2T$ and amplitude V . No current is flowing initially.

Matlab Codes

How to solve an IVP using Laplace transforms in Matlab?

Matlab Codes

How to solve an IVP using Laplace transforms in Matlab?

Here we are:

$$y''(t) + 6y'(t) + 5y(t) = f(t), \quad y(0) = 1, y'(0) = -1$$

Let $f(t) = 3e^{-2t}$

Matlab Codes

How to solve an IVP using Laplace transforms in Matlab?

Here we are:

$$y''(t) + 6y'(t) + 5y(t) = f(t), \quad y(0) = 1, y'(0) = -1$$

Let $f(t) = 3e^{-2t}$

```
>> syms s t Y
f = exp(-2*t)*3
F=laplace(f) % define the Laplace transform of the RHS part f(t)
y2=Y*s.^2-1*s-(-1) % define the Laplace transform of y''
y1=Y*s-1 % define the Laplace transform of y'
Yimage = solve(y2 + 6*y1 + 5*Y-F,Y) % express the image of y
ilaplace(Yimage,s,t) % find the original, i.e. the solution of the IVP
```



Yimage =

$(s + 3/(s + 2) + 5)/(s^2 + 6*s + 5)$

ans =

$(7*\exp(-t))/4 - \exp(-2*t) + \exp(-5*t)/4$

Matlab Codes



Matlab Codes

$$f(t) = \begin{cases} \sin t & 0 < t < \pi \\ -1 & \pi < t < 2\pi \\ 0 & t > 2\pi \end{cases}$$

Matlab Codes

$$f(t) = \begin{cases} \sin t & 0 < t < \pi \\ -1 & \pi < t < 2\pi \\ 0 & t > 2\pi \end{cases}$$

$$f(t) = \sin t + H(t - \pi)(-1 - \sin t) + H(t - 2\pi)(0 - (-1))$$

Matlab Codes

$$f(t) = \begin{cases} \sin t & 0 < t < \pi \\ -1 & \pi < t < 2\pi \\ 0 & t > 2\pi \end{cases}$$

$$f(t) = \sin t + H(t - \pi)(-1 - \sin t) + H(t - 2\pi)(0 - (-1))$$

```
>> f=sin(t)+ (-1-sin(t))*heaviside(t-pi) + heaviside(t-2*pi)
```

Matlab Codes

$$f(t) = \begin{cases} \sin t & 0 < t < \pi \\ -1 & \pi < t < 2\pi \\ 0 & t > 2\pi \end{cases}$$

$$f(t) = \sin t + H(t - \pi)(-1 - \sin t) + H(t - 2\pi)(0 - (-1))$$

```
>> f=sin(t)+ (-1-sin(t))*heaviside(t-pi) + heaviside(t-2*pi)
```

$$f(t) = 1 + 2\delta(t - 1)$$

Matlab Codes

$$f(t) = \begin{cases} \sin t & 0 < t < \pi \\ -1 & \pi < t < 2\pi \\ 0 & t > 2\pi \end{cases}$$

$$f(t) = \sin t + H(t - \pi)(-1 - \sin t) + H(t - 2\pi)(0 - (-1))$$

```
>> f=sin(t)+ (-1-sin(t))*heaviside(t-pi) + heaviside(t-2*pi)
```

$$f(t) = 1 + 2\delta(t - 1)$$

```
>> f = 1 + 2*dirac(t-1)
```

```
f =
```

```
2*dirac(t - 1) + 1
```

```
>> F=laplace(f)
```

```
F =
```

```
2*exp(-s) + 1/s
```