

vv256: Week 1-2. Introduction to solutions of differential equations. First-order ODEs.

Dr.Olga Danilkina

UM-SJTU Joint Institute

September 20, 2019

Outline

- 1 Lecture 1: Introduction
 - Course Info
 - Differential Equations and Their Solutions
 - Classification of DE
- 2 Lecture 2: Separable and linear ordinary differential equations
 - Separable Equations
 - Linear equations
- 3 Lecture 3: Other first-order ordinary differential equations
 - Homogeneous Polar Equations
 - Bernoulli Equations
 - Riccati Equations
 - Exact Equations

Outline

- 1 Lecture 1: Introduction
 - Course Info
 - Differential Equations and Their Solutions
 - Classification of DE
- 2 Lecture 2: Separable and linear ordinary differential equations
 - Separable Equations
 - Linear equations
- 3 Lecture 3: Other first-order ordinary differential equations
 - Homogeneous Polar Equations
 - Bernoulli Equations
 - Riccati Equations
 - Exact Equations

Recommended Literature

- George F. Simmons. Differential equations with applications and historical notes, third edition, CRC Press, 2017.
- William E. Boyce, Richard C. DiPrima, Douglas B. Meade. Elementary Differential Equations and Boundary Value Problems, 11th edition, Wiley, 2017.
- Kam Tim Chau. Theory of differential equations in engineering and mechanics, CRC Press, 2017.
- Yuefan Deng. Lectures, problems and solutions for ordinary differential equations, second addition, World Scientific, 2018.
- Marcelo Epstein. Partial differential equations. Mathematical techniques for engineers, Springer, 2017.

Course Assessment

Final Grade (100%)

**Assignments (20%) + Midterm Exam I (20%) +
Midterm Exam 2 (20%) + Final Exam (40%)**

Differential Equations and Their Solutions

Physical (chemical, biological etc) process \rightarrow mathematical model
 \rightarrow differential equation

Definition. A differential equation (DE) is a relation that contains an unknown function, say y , and one or more of its derivatives. If $y = y(x)$ then the general form of an ordinary differential equation (ODE) is


$$F(y, y', y'', \dots, y^{(n)}) = 0.$$

If $y = y(x_1, x_2)$ then the general representation of a partial differential equation (PDE) is

$$T\left(y, \frac{\partial y}{\partial x_1}, \frac{\partial y}{\partial x_2}, \frac{\partial^2 y}{\partial x_1 \partial x_2}, \frac{\partial^2 y}{\partial x_1^2}, \frac{\partial^2 y}{\partial x_2^2}, \dots, \frac{\partial^n y}{\partial x_1^n}, \frac{\partial^n y}{\partial x_2^n}, \frac{\partial^n y}{\partial x_1^{n-1} \partial x_2}\right) = 0.$$

Example 1: RC Circuit

Consider the circuit with resistor and capacitor as shown below. Kirchhoff's voltage law \rightarrow the sum of the voltage drops around a closed circuit is zero.



Example 1: RC Circuit

Ohm's law \rightarrow

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Kirchhoff's current law \rightarrow

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Kirchhoff's current law \rightarrow the current through the resistor is equal the rate of change of the charge on the capacitor $\rightarrow I = \frac{dq}{dt}$.

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Kirchhoff's current law \rightarrow the current through the resistor is equal
the rate of change of the charge on the capacitor $\rightarrow I = \frac{dq}{dt}$.

The charge of the capacitor is $q = CV$.

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Kirchhoff's current law \rightarrow the current through the resistor is equal the rate of change of the charge on the capacitor $\rightarrow I = \frac{dq}{dt}$.

The charge of the capacitor is $q = CV$.

$$\frac{dV}{dt} = -\frac{1}{RC}V, \quad V(0) = V_0.$$

Example 1: RC Circuit

Ohm's law \rightarrow the voltage drop over the resistor is IR .

$$IR + V = 0$$

Kirchhoff's current law \rightarrow the current through the resistor is equal the rate of change of the charge on the capacitor $\rightarrow I = \frac{dq}{dt}$.
The charge of the capacitor is $q = CV$.


$$\frac{dV}{dt} = -\frac{1}{RC}V, \quad V(0) = V_0.$$

It is the initial-value problem (IVP) with the solution

$$V(t) = V_0 e^{-\frac{1}{RC}t}.$$

Example 1: RC Circuit

Graph of $V(t) = V_0 \exp(-\frac{t}{RC})$ for $RC = 0.5, 1$, and 2 .



Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law \rightarrow

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object).

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object). **What are forces that act on the particle?**

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object). **What are forces that act on the particle?**

The gravity exerts a force equal mg , where $g \approx 9.8$ is the acceleration due to gravity, it acts in the downward (positive) direction.

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object). **What are forces that act on the particle?**

The gravity exerts a force equal mg , where $g \approx 9.8$ is the acceleration due to gravity, it acts in the downward (positive) direction.

What else?

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object). **What are forces that act on the particle?**

The gravity exerts a force equal mg , where $g \approx 9.8$ is the acceleration due to gravity, it acts in the downward (positive) direction.

What else? There is also a force due to the resistance of the ambient medium to motion, its proportional to the velocity and acts in upward (negative direction), it equals $-\gamma v$.

Example 2: Free fall in gravity

If v is the velocity at time $t > 0$ of a material particle falling in a gravitational field then $v = v(t)$.

The second Newton's law $\rightarrow ma = F$ (the mass of the object times its acceleration is equal to the net force on the object). **What are forces that act on the particle?**

The gravity exerts a force equal mg , where $g \approx 9.8$ is the acceleration due to gravity, it acts in the downward (positive) direction.

What else? There is also a force due to the resistance of the ambient medium to motion, its proportional to the velocity and acts in upward (negative direction), it equals $-\gamma v$.

$$m \frac{dv}{dt} = mg - \gamma v$$

Example 2: Free fall in gravity

Exercise: Let an object of mass $m = 10$ dropped into a liquid-filled reservoir reach the bottom with velocity $v_b = 24.5$. If $g = 9.8$ and the motion resistance coefficient of the liquid is $\gamma = 2$, compute the depth of the reservoir.

Example 2: Free fall in gravity

Exercise: Let an object of mass $m = 10$ dropped into a liquid-filled reservoir reach the bottom with velocity $v_b = 24.5$. If $g = 9.8$ and the motion resistance coefficient of the liquid is $\gamma = 2$, compute the depth of the reservoir.

Hint:

$$v(t) = \frac{mg}{\gamma} + \left(v_0 - \frac{mg}{\gamma}\right)e^{-\frac{\gamma}{m}t}$$

Example 2: Free fall in gravity

Exercise: Let an object of mass $m = 10$ dropped into a liquid-filled reservoir reach the bottom with velocity $v_b = 24.5$. If $g = 9.8$ and the motion resistance coefficient of the liquid is $\gamma = 2$, compute the depth of the reservoir.

Hint:

$$v(t) = \frac{mg}{\gamma} + \left(v_0 - \frac{mg}{\gamma}\right)e^{-\frac{\gamma}{m}t}$$

The position $y(t) = v'(t)$ and

Example 2: Free fall in gravity

Exercise: Let an object of mass $m = 10$ dropped into a liquid-filled reservoir reach the bottom with velocity $v_b = 24.5$. If $g = 9.8$ and the motion resistance coefficient of the liquid is $\gamma = 2$, compute the depth of the reservoir.

Hint:

$$v(t) = \frac{mg}{\gamma} + \left(v_0 - \frac{mg}{\gamma}\right)e^{-\frac{\gamma}{m}t}$$

The position $y(t) = \int v(t) dt$ and $v(0) = v_0 = 0$.

Example 3: Population growth

Let $P(t)$ be the size of a population at time t , and let $\beta(t)$ and $\delta(t)$ be the birth and death rates (measured per unit of population per unit time) within the population.

Example 3: Population growth

Let $P(t)$ be the size of a population at time t , and let $\beta(t)$ and $\delta(t)$ be the birth and death rates (measured per unit of population per unit time) within the population. For a very short interval of time Δt , we may assume that β and δ are constant, so we approximate the change ΔP in the population during that interval by

$$\Delta P \approx (\beta - \delta)P\Delta t$$

or

Example 3: Population growth

Let $P(t)$ be the size of a population at time t , and let $\beta(t)$ and $\delta(t)$ be the birth and death rates (measured per unit of population per unit time) within the population. For a very short interval of time Δt , we may assume that β and δ are constant, so we approximate the change ΔP in the population during that interval by

$$\Delta P \approx (\beta - \delta)P\Delta t$$

or

$$P' = (\beta - \delta)P$$

Example 3: Population growth

Let $P(t)$ be the size of a population at time t , and let $\beta(t)$ and $\delta(t)$ be the birth and death rates (measured per unit of population per unit time) within the population. For a very short interval of time Δt , we may assume that β and δ are constant, so we approximate the change ΔP in the population during that interval by

$$\Delta P \approx (\beta - \delta)P\Delta t$$

or

$$P' = (\beta - \delta)P$$

If the size of the population at $t = 0$ is $P(0) = P_0$ then $P(t) = P_0 e^{(\beta - \delta)t}$.

Example 3: Population growth

What happens when $\delta > \beta$ and $\beta < \delta$?

Example 3: Population growth

What happens when $\delta > \beta$ and $\beta < \delta$?

The population decreases asymptotically to zero if $\delta > \beta$ and increases exponentially without bound if $\beta < \delta$.

Example 3: Population growth

What happens when $\delta > \beta$ and $\beta < \delta$?

The population decreases asymptotically to zero if $\delta > \beta$ and increases exponentially without bound if $\beta < \delta$.

\Rightarrow the model governed by equation $P' = (\beta - \delta)P$ is valid only for relatively small populations and for a finite time interval.

Example 3: Population growth

What happens when $\delta > \beta$ and $\beta < \delta$?

The population decreases asymptotically to zero if $\delta > \beta$ and increases exponentially without bound if $\beta < \delta$.

\Rightarrow the model governed by equation $P' = (\beta - \delta)P$ is valid only for relatively small populations and for a finite time interval.

If $\beta = \beta(t)$, $\delta = \delta(t)$ then we obtain the same model with the solution

$$P(t) = P_0 e^{\int_0^t (\beta(\tau) - \delta(\tau)) d\tau}, \quad t > 0.$$

Example 3: Population growth

What happens when $\delta > \beta$ and $\beta < \delta$?

The population decreases asymptotically to zero if $\delta > \beta$ and increases exponentially without bound if $\beta < \delta$.

\Rightarrow the model governed by equation $P' = (\beta - \delta)P$ is valid only for relatively small populations and for a finite time interval.

If $\beta = \beta(t)$, $\delta = \delta(t)$ then we obtain the same model with the solution

$$P(t) = P_0 e^{\int_0^t (\beta(\tau) - \delta(\tau)) d\tau}, \quad t > 0.$$

Exercise: Suppose that the annual birth and death rates in a population of the initial size $P_0 = 100$ are $\beta(t) = 2t + 1$ and $\delta(t) = 4t + 4$. Determine how long it will take for the size of the population to decrease to 40.

Example 4: Radioactive decay

If $k = \text{const} > 0$ is the rate of decay of a radioactive isotope (the number of decaying atoms per unit of atom 'population' per unit time), then the approximate change ΔN in the number $N(t)$ of atoms during a very short time interval Δt is

$$\Delta N \approx -kN\Delta t.$$

Example 4: Radioactive decay

If $k = \text{const} > 0$ is the rate of decay of a radioactive isotope (the number of decaying atoms per unit of atom 'population' per unit time), then the approximate change ΔN in the number $N(t)$ of atoms during a very short time interval Δt is

$$\Delta N \approx -kN\Delta t.$$

We obtain the DE

$$N' = -kN.$$

Example 4: Radioactive decay

If $k = \text{const} > 0$ is the rate of decay of a radioactive isotope (the number of decaying atoms per unit of atom 'population' per unit time), then the approximate change ΔN in the number $N(t)$ of atoms during a very short time interval Δt is

$$\Delta N \approx -kN\Delta t.$$

We obtain the DE

$$N' = -kN.$$

After separating the variables, integrating, and using an IC of the form $N(0) = N_0$, we find the solution

$$N(t) = N_0 e^{-kt}, \quad t > 0.$$

Example 4: Radioactive decay

An important characteristic of a radioactive substance is its so-called half-life, which is the length of time t^* it takes the substance to decay by half.

Example 4: Radioactive decay

An important characteristic of a radioactive substance is its so-called half-life, which is the length of time t^* it takes the substance to decay by half. As $N(t^*) = N_0/2$, so from $N(t) = N_0 e^{-kt}$ it follows that

$$\frac{1}{2}N_0 = N_0 e^{-kt^*}.$$

Example 4: Radioactive decay

An important characteristic of a radioactive substance is its so-called half-life, which is the length of time t^* it takes the substance to decay by half. As $N(t^*) = N_0/2$, so from $N(t) = N_0e^{-kt}$ it follows that

$$\frac{1}{2}N_0 = N_0e^{-kt^*}.$$

Divide both sides by N_0 and take logarithms to find that

$$t^* = \frac{\ln 2}{k}.$$

Example 5: Newton's law of cooling

The idea: to describe convection cooling in which the heat transfer coefficient is independent of the temperature difference between the object and its thermo environment (Newton, 1701).

Example 5: Newton's law of cooling

The idea: to describe convection cooling in which the heat transfer coefficient is independent of the temperature difference between the object and its thermo environment (Newton, 1701).

Newton's law of cooling \rightarrow the rate of change of the temperature of an object is proportional to the difference between its own temperature and the ambient temperature.

Example 5: Newton's law of cooling

The idea: to describe convection cooling in which the heat transfer coefficient is independent of the temperature difference between the object and its thermo environment (Newton, 1701).

Newton's law of cooling \rightarrow the rate of change of the temperature of an object is proportional to the difference between its own temperature and the ambient temperature.

Let $T(t)$ be the temperature at time $t > 0$ of an object immersed in an outside medium of temperature θ and k is the heat transfer coefficient of the object material.

Example 5: Newton's law of cooling

The idea: to describe convection cooling in which the heat transfer coefficient is independent of the temperature difference between the object and its thermo environment (Newton, 1701).

Newton's law of cooling \rightarrow the rate of change of the temperature of an object is proportional to the difference between its own temperature and the ambient temperature.

Let $T(t)$ be the temperature at time $t > 0$ of an object immersed in an outside medium of temperature θ and k is the heat transfer coefficient of the object material.

$$T' = -k(T - \theta), \quad k > 0$$

Why do we need negative sign on the right-hand of the equation?

Exercise: Solve the equation.

Exercise: Solve the equation.

Answer:

Exercise: Solve the equation.

Answer:

$$T = \theta + (T_0 - \theta)e^{-kt}, \quad t > 0.$$

Exercise: Solve the equation.

Answer:

$$T = \theta + (T_0 - \theta)e^{-kt}, \quad t > 0.$$

Problem: Take a pot with boiling water and put it in a sink with running water with the constant temperature 5°C . The temperature of hot water goes to 60°C in 10 minutes. How long will it take to cool hot water to 20°C ?

Classification of Differential Equations

- ODEs and PDEs (number of independent variables)
- Order of a DE
- Linear and nonlinear DEs
- Homogeneous and non-homogeneous DEs

Classification of Differential Equations

- ODEs and PDEs (number of independent variables)
- Order of a DE
- Linear and nonlinear DEs
- Homogeneous and non-homogeneous DEs
- Initial-value problems (propagation problems)
- Boundary-value problems (equilibrium problems)

Outline

- 1 Lecture 1: Introduction
 - Course Info
 - Differential Equations and Their Solutions
 - Classification of DE
- 2 Lecture 2: Separable and linear ordinary differential equations
 - Separable Equations
 - Linear equations
- 3 Lecture 3: Other first-order ordinary differential equations
 - Homogeneous Polar Equations
 - Bernoulli Equations
 - Riccati Equations
 - Exact Equations

Existence and Uniqueness Theorem for the IVP

Consider the Initial Value Problem (IVP)

$$y' = f(x, y), \quad y(x_0) = y_0.$$

Existence and Uniqueness Theorem for the IVP

Consider the Initial Value Problem (IVP)

$$y' = f(x, y), \quad y(x_0) = y_0.$$

Suppose $f(x, y)$ and $\frac{\partial f}{\partial y}(x, y)$ are continuous functions in some open rectangle $R = \{(x, y) : |x - x_0| < a, |y - y_0| < b\}$, $a, b > 0$, and hence, there exist $K, L > 0$ such that

$$(a) |f(x, y)| \leq K, \quad (b) \left| \frac{\partial f}{\partial y} \right| \leq L \quad \forall (x, y).$$

Then the IVP has a unique solution in the interval $|x - x_0| \leq \alpha$, where $\alpha = \min\{a, \frac{b}{K}\}$.

Separable equations

Definition. A separable ODE is an equation of the form

$$\frac{dy}{dx} = f(x)g(y),$$

where f and g are given functions.

Separable equations

Definition. A separable ODE is an equation of the form

$$\frac{dy}{dx} = f(x)g(y),$$

where f and g are given functions.

Remark. If there is any value y_0 such that $g(y_0) = 0$, then $y = y_0$ is a solution (equilibrium solution). To find other (non-constant) solutions, we assume that $g(y) \neq 0$.

Separable equations

- Apply the definition of the differential

Separable equations

- Apply the definition of the differential

$$dy =$$

Separable equations

- Apply the definition of the differential

$$dy = y'(x)dx =$$

Separable equations

- Apply the definition of the differential

$$dy = y'(x)dx = \frac{dy}{dx}dx =$$

Separable equations

- Apply the definition of the differential

$$dy = y'(x)dx = \frac{dy}{dx}dx = f(x)g(y)dx. \quad (1)$$

Separable equations

- Apply the definition of the differential

$$dy = y'(x)dx = \frac{dy}{dx}dx = f(x)g(y)dx. \quad (1)$$

- Separate variables

$$\frac{1}{g(y)}dy = f(x)dx, \quad \text{and}$$

Separable equations

- Apply the definition of the differential

$$dy = y'(x)dx = \frac{dy}{dx}dx = f(x)g(y)dx. \quad (1)$$

- Separate variables

$$\frac{1}{g(y)}dy = f(x)dx, \quad \text{and}$$

- Integrate each side with respect to its variable to obtain

$$G(y) = F(x) + C, \quad (2)$$

where F and G are anti-derivatives of f and $1/g$, and C is an arbitrary constant.

Separable equations

- For each value of C , (2) is a connection between x and $y \Rightarrow$ A function $y = y(x)$ is defined implicitly.

Separable equations

- For each value of C , (2) is a connection between x and $y \Rightarrow$ A function $y = y(x)$ is defined implicitly.
- The solution of (2) also satisfies (1):

$$\frac{d}{dx} G(y(x)) = \frac{d}{dy} G(y) \frac{dy}{dx} = \frac{1}{g(y)} \frac{dy}{dx} = f(x).$$

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$.

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$. Then for $y \neq 0$

$$\int \frac{dy}{y} = \int (-8x)dx,$$

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$. Then for $y \neq 0$

$$\int \frac{dy}{y} = \int (-8x)dx,$$

from which $\ln |y| = -4x^2 + C$.

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$. Then for $y \neq 0$

$$\int \frac{dy}{y} = \int (-8x)dx,$$

from which $\ln |y| = -4x^2 + C$.

Taking the exponent, we get $|y| = e^{-4x^2+C} = e^C e^{-4x^2}$, so

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$. Then for $y \neq 0$

$$\int \frac{dy}{y} = \int (-8x)dx,$$

from which $\ln |y| = -4x^2 + C$.

Taking the exponent, we get $|y| = e^{-4x^2+C} = e^C e^{-4x^2}$, so

$$y(x) = \pm e^C e^{-4x^2} = C_1 e^{-4x^2},$$

where $C_1 = \pm e^C$ is a non-zero constant (it generates all non zero solutions).

Separable equations: Example 1

Consider the equation $y' + 8xy = 0$ and bring it to the form

$$\frac{dy}{dx} = -8xy.$$

The equilibrium solution is $y = 0$. Then for $y \neq 0$

$$\int \frac{dy}{y} = \int (-8x) dx,$$

from which $\ln |y| = -4x^2 + C$.

Taking the exponent, we get $|y| = e^{-4x^2+C} = e^C e^{-4x^2}$, so

$$y(x) = \pm e^C e^{-4x^2} = C_1 e^{-4x^2},$$

where $C_1 = \pm e^C$ is a non-zero constant (it generates all non zero solutions). If we allow $C_1 = 0$ then the solution $y = C_1 e^{-4x^2}$ includes also the equilibrium solution.

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions?

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions? No, $e^y \neq 0$.

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions? No, $e^y \neq 0$.

Separate variables and integrate to obtain

$$\int e^{-y} dy = - \int 4xe^{-2x} dx.$$

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions? No, $e^y \neq 0$.

Separate variables and integrate to obtain

$$\int e^{-y} dy = - \int 4xe^{-2x} dx.$$

Integrate by part on the right-hand side:

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions? No, $e^y \neq 0$.

Separate variables and integrate to obtain

$$\int e^{-y} dy = - \int 4xe^{-2x} dx.$$

Integrate by part on the right-hand side:

$$-e^{-y} = 2xe^{-2x} - \int 2e^{-2x} dx =$$

Separable equations: Example 2

Consider the IVP $y' + 4xe^{y-2x} = 0$, $y(0) = 0$. The equation can be rewritten as

$$\frac{dy}{dx} = -4xe^{-2x}e^y.$$

Are there equilibrium solutions? No, $e^y \neq 0$.

Separate variables and integrate to obtain

$$\int e^{-y} dy = - \int 4xe^{-2x} dx.$$

Integrate by part on the right-hand side:

$$-e^{-y} = 2xe^{-2x} - \int 2e^{-2x} dx = (2x + 1)e^{-2x} + C.$$

Change the signs of both sides, take logarithms to produce the general solution

Separable equations: Example 2

$$y(x) = -\ln[-(2x + 1)e^{-2x} - C].$$

Since $y(0) = 0$, $C = -2$ and the solution of the IVP is

$$y(x) = -\ln[2 - (2x + 1)e^{-2x}].$$

Separable equations: Example 2

$$y(x) = -\ln[-(2x+1)e^{-2x} - C].$$

Since $y(0) = 0$, $C = -2$ and the solution of the IVP is

$$y(x) = -\ln[2 - (2x+1)e^{-2x}].$$

Exercise: Find the solution of the IVP

$$xy' = y + 2, \quad y(1) = -1.$$

Separable equations: Example 2

$$y(x) = -\ln[-(2x+1)e^{-2x} - C].$$

Since $y(0) = 0$, $C = -2$ and the solution of the IVP is

$$y(x) = -\ln[2 - (2x+1)e^{-2x}].$$

Exercise: Find the solution of the IVP

$$xy' = y + 2, \quad y(1) = -1.$$

Answer: $y = x - 2$.

Separable equations: Example 3

Consider the IVP $2(x+1)yy' - y^2 = 2, y(5) = 2$. No equilibrium solutions, and hence, for $x \neq -1$

$$\int \frac{2y \, dy}{y^2 + 2} = \int \frac{dx}{x + 1}.$$

Separable equations: Example 3

Consider the IVP $2(x+1)yy' - y^2 = 2, y(5) = 2$. No equilibrium solutions, and hence, for $x \neq -1$

$$\int \frac{2y \, dy}{y^2 + 2} = \int \frac{dx}{x + 1}.$$

$$\ln(y^2 + 2) = \ln|x + 1| + C, \quad C = \text{const.}$$

Separable equations: Example 3

Consider the IVP $2(x+1)yy' - y^2 = 2, y(5) = 2$. No equilibrium solutions, and hence, for $x \neq -1$

$$\int \frac{2y dy}{y^2 + 2} = \int \frac{dx}{x + 1}.$$

$$\ln(y^2 + 2) = \ln|x + 1| + C, \quad C = \text{const.}$$

Therefore,

$$y^2 = C_1(x + 1) - 2, \quad C_1 = \text{const} \neq 0.$$

Separable equations: Example 3

Consider the IVP $2(x+1)yy' - y^2 = 2, y(5) = 2$. No equilibrium solutions, and hence, for $x \neq -1$

$$\int \frac{2y dy}{y^2 + 2} = \int \frac{dx}{x+1}.$$

$$\ln(y^2 + 2) = \ln|x+1| + C, \quad C = \text{const.}$$

Therefore,

$$y^2 = C_1(x+1) - 2, \quad C_1 = \text{const} \neq 0.$$

Applying IC, we obtain $y^2 = x - 1$, or $y = \pm(x-1)^{1/2}$.

Separable equations: Example 3

Consider the IVP $2(x+1)yy' - y^2 = 2, y(5) = 2$. No equilibrium solutions, and hence, for $x \neq -1$

$$\int \frac{2y dy}{y^2 + 2} = \int \frac{dx}{x + 1}.$$

$$\ln(y^2 + 2) = \ln|x + 1| + C, \quad C = \text{const.}$$

Therefore,

$$y^2 = C_1(x + 1) - 2, \quad C_1 = \text{const} \neq 0.$$

Applying IC, we obtain $y^2 = x - 1$, or $y = \pm(x - 1)^{1/2}$. However, $y = -(x - 1)^{1/2}$ does not satisfy the IC.

Separable equations: Example 4

Consider the IVP

$$(5y^4 + 3y^2 + e^y)y' = \cos x, \quad y(0) = 0.$$

Applying separation of variables, we obtain **the family of all solution curves** for the differential equation.

Separable equations: Example 4

Consider the IVP

$$(5y^4 + 3y^2 + e^y)y' = \cos x, \quad y(0) = 0.$$

Applying separation of variables, we obtain **the family of all solution curves** for the differential equation.

$$y^5 + y^3 + e^y = \sin x + C, \quad C = \text{const.}$$

The IC yields

Separable equations: Example 4

Consider the IVP

$$(5y^4 + 3y^2 + e^y)y' = \cos x, \quad y(0) = 0.$$

Applying separation of variables, we obtain **the family of all solution curves** for the differential equation.

$$y^5 + y^3 + e^y = \sin x + C, \quad C = \text{const.}$$

The IC yields $C = 1$,

Separable equations: Example 4

Consider the IVP

$$(5y^4 + 3y^2 + e^y)y' = \cos x, \quad y(0) = 0.$$

Applying separation of variables, we obtain **the family of all solution curves** for the differential equation.

$$y^5 + y^3 + e^y = \sin x + C, \quad C = \text{const.}$$

The IC yields $C = 1$, so the solution curve passing through the point $(0, 0)$ has the equation

$$y^5 + y^3 + e^y = \sin x + 1.$$

Separable equations: Exercises

Solve the following IVP:

- ① $y' = -4xy^2, y(0) = 1$
- ② $y' = (3 - 2x)y, y(2) = e^6$
- ③ $y' = (x - 3)(y^2 + 1), y(0) = 1$
- ④ $3(x^2 + 2)y^2y' = 4x, y(1) = (\ln 9)^{1/3}$
- ⑤ $y' = xe^{2x}/(y^4 + 2y), y(0) = -1$

Linear Equations

A general form of a first-order linear DE is

$$y' + p(t)y = q(t),$$

where p and q are coefficient functions.

Linear Equations

A general form of a first-order linear DE is

$$y' + p(t)y = q(t),$$

where p and q are coefficient functions. To solve the equation,

- multiply the equation by an unknown nonzero function $\mu(t)$ (integrating factor)

$$\mu y' + \mu p y = \mu q,$$

Linear Equations

A general form of a first-order linear DE is

$$y' + p(t)y = q(t),$$

where p and q are coefficient functions. To solve the equation,

- multiply the equation by an unknown nonzero function $\mu(t)$ (integrating factor)

$$\mu y' + \mu p y = \mu q,$$

- choose μ so that the left-hand side is the derivative of the product μy :

Linear Equations

A general form of a first-order linear DE is

$$y' + p(t)y = q(t),$$

where p and q are coefficient functions. To solve the equation,

- multiply the equation by an unknown nonzero function $\mu(t)$ (integrating factor)

$$\mu y' + \mu p y = \mu q,$$

- choose μ so that the left-hand side is the derivative of the product μy :

$$\mu y' + \mu p y = (\mu y)' =$$

Linear Equations

A general form of a first-order linear DE is

$$y' + p(t)y = q(t),$$

where p and q are coefficient functions. To solve the equation,

- multiply the equation by an unknown nonzero function $\mu(t)$ (integrating factor)

$$\mu y' + \mu p y = \mu q,$$

- choose μ so that the left-hand side is the derivative of the product μy :

$$\mu y' + \mu p y = (\mu y)' = \mu y' + \mu' y \Rightarrow \mu' = \mu p$$

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow$$

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

$$\text{So, } \mu = Ce^{\int p dt}, \quad C = \text{const} \neq 0.$$

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

$$\text{So, } \mu = Ce^{\int p dt}, \quad C = \text{const} \neq 0.$$

- We need just one such function, let $C = 1$ and hence,

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

$$\text{So, } \mu = Ce^{\int p dt}, \quad C = \text{const} \neq 0.$$

- We need just one such function, let $C = 1$ and hence, $\mu = e^{\int p dt}$.

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

$$\text{So, } \mu = Ce^{\int p dt}, \quad C = \text{const} \neq 0.$$

- We need just one such function, let $C = 1$ and hence, $\mu = e^{\int p dt}$.
- Substitute μ into $\mu y' + \mu p y = \mu q \Rightarrow (\mu y)' = \mu q$ and,

Linear Equations

- find the solution of the equation $\mu' = \mu p$ by separation of variables

$$\int \frac{d\mu}{\mu} = \int p dt \quad \Rightarrow \quad \ln |\mu| = \int p dt,$$

$$\text{So, } \mu = Ce^{\int p dt}, \quad C = \text{const} \neq 0.$$

- We need just one such function, let $C = 1$ and hence, $\mu = e^{\int p dt}$.
- Substitute μ into $\mu y' + \mu p y = \mu q \Rightarrow (\mu y)' = \mu q$ and,
- finally find

$$y(t) = \frac{1}{\mu(t)} \left(\int \mu(t) q(t) dt + C \right).$$

Linear Equations

Can we include an IC $y(t_0) = y_0$ into the solution directly?

Linear Equations

Can we include an IC $y(t_0) = y_0$ into the solution directly? Yes.

Integrate $(\mu y)' = \mu q$ from t_0 to t , then

Linear Equations

Can we include an IC $y(t_0) = y_0$ into the solution directly? Yes.

Integrate $(\mu y)' = \mu q$ from t_0 to t , then

$$\mu(t)y(t) - \mu(t_0)y(t_0) = \int_{t_0}^t \mu(\tau)q(\tau) d\tau$$

and

$$y(t) = \frac{1}{\mu(t)} \left(\int_{t_0}^t \mu(\tau)q(\tau) d\tau + \mu(t_0)y(t_0) \right).$$

Linear Equations: Example 1

Consider the IVP

$$y' - 3y = 6, y(0) = -1.$$

Then

$$\mu(t) = e^{\int (-3) dt} = e^{-3t},$$

Linear Equations: Example 1

Consider the IVP

$$y' - 3y = 6, y(0) = -1.$$

Then

$$\mu(t) = e^{\int (-3) dt} = e^{-3t}, \quad y(t) = e^{3t} \left(\int 6e^{-3t} dt + C \right) = Ce^{3t} - 2.$$

Applying the IC, we obtain $C = 1$ and hence, $y(t) = e^{3t} - 2$.

Linear Equations: Example 2

Consider the IVP

$$(t^2 + 1)y' - ty = 2t(t^2 + 1)^2, y(0) = 2/3.$$

Linear Equations: Example 2

Consider the IVP

$$(t^2 + 1)y' - ty = 2t(t^2 + 1)^2, y(0) = 2/3.$$

Then $p(t) = -t/(t^2 + 1)$, $q(t) = 2t(t^2 + 1)$ and

Linear Equations: Example 2

Consider the IVP

$$(t^2 + 1)y' - ty = 2t(t^2 + 1)^2, y(0) = 2/3.$$

Then $p(t) = -t/(t^2 + 1)$, $q(t) = 2t(t^2 + 1)$ and

$$\mu(t) = \exp \left\{ - \int \frac{t}{t^2 + 1} dt \right\} = e^{-(1/2)\ln(t^2+1)} = (t^2 + 1)^{-1/2}.$$

Linear Equations: Example 2

Consider the IVP

$$(t^2 + 1)y' - ty = 2t(t^2 + 1)^2, y(0) = 2/3.$$

Then $p(t) = -t/(t^2 + 1)$, $q(t) = 2t(t^2 + 1)$ and

$$\mu(t) = \exp \left\{ - \int \frac{t}{t^2 + 1} dt \right\} = e^{-(1/2)\ln(t^2+1)} = (t^2 + 1)^{-1/2}.$$

Then the solution

$$y(t) = (t^2 + 1)^{1/2} \left\{ \int_0^t (\tau^2 + 1)^{-1/2} 2\tau(\tau^2 + 1) d\tau + \mu(0)y(0) \right\} =$$

Linear Equations: Example 2

Consider the IVP

$$(t^2 + 1)y' - ty = 2t(t^2 + 1)^2, y(0) = 2/3.$$

Then $p(t) = -t/(t^2 + 1)$, $q(t) = 2t(t^2 + 1)$ and

$$\mu(t) = \exp \left\{ - \int \frac{t}{t^2 + 1} dt \right\} = e^{-(1/2)\ln(t^2+1)} = (t^2 + 1)^{-1/2}.$$

Then the solution

$$\begin{aligned} y(t) &= (t^2 + 1)^{1/2} \left\{ \int_0^t (\tau^2 + 1)^{-1/2} 2\tau(\tau^2 + 1) d\tau + \mu(0)y(0) \right\} = \\ &= \frac{2}{3}(t^2 + 1)^2. \end{aligned}$$

Linear Equations: Exercises

Solve the following IVP:

- ① $(t-1)y' + y = (t-1)e^t$, $y(2) = 3$ Ans: $\frac{(t-2)e^t + 3}{t-1}$
- ② $y' + 4y + 16 = 0$, $y(0) = -2$ Ans: $y = 2e^{-4t} - 4$
- ③ $ty' + 4y = 6t^2$, $y(1) = 4$ Ans: $t^2 + 3t^{-4}$
- ④ $y' = (2+y)\sin t$, $y(\pi/2) = -3$ Ans: $-e^{-\cos t} - 2$
- ⑤ $2ty' - y = 2/\sqrt{t}$, $y(1) = 1$

Outline

- 1 Lecture 1: Introduction
 - Course Info
 - Differential Equations and Their Solutions
 - Classification of DE
- 2 Lecture 2: Separable and linear ordinary differential equations
 - Separable Equations
 - Linear equations
- 3 Lecture 3: Other first-order ordinary differential equations
 - Homogeneous Polar Equations
 - Bernoulli Equations
 - Riccati Equations
 - Exact Equations

Homogeneous Polar Equations

What is a homogeneous polar equation?

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ?

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ? $y' = v + xv'$
- The new unknown function v satisfies the DE

$$\frac{dv}{dx} = \frac{f(v) - v}{x}$$

What is the type of this equation?

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ? $y' = v + xv'$
- The new unknown function v satisfies the DE

$$\frac{dv}{dx} = \frac{f(v) - v}{x}$$

What is the type of this equation? Separable

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ? $y' = v + xv'$
- The new unknown function v satisfies the DE

$$\frac{dv}{dx} = \frac{f(v) - v}{x}$$

What is the type of this equation? Separable

- For $f(v) - v \neq 0$ $\int \frac{dv}{f(v) - v} = \int \frac{dx}{x}$

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ? $y' = v + xv'$
- The new unknown function v satisfies the DE

$$\frac{dv}{dx} = \frac{f(v) - v}{x}$$

What is the type of this equation? Separable

- For $f(v) - v \neq 0$ $\int \frac{dv}{f(v) - v} = \int \frac{dx}{x}$
- What happens when $f(v) - v = 0$?

Homogeneous Polar Equations

What is a homogeneous polar equation?

$$y'(x) = f\left(\frac{y}{x}\right), \quad x \neq 0,$$

where f is a given one-variable function.

- Make the substitution $y(x) = xv(x)$. What is the derivative of y ? $y' = v + xv'$
- The new unknown function v satisfies the DE

$$\frac{dv}{dx} = \frac{f(v) - v}{x}$$

What is the type of this equation? Separable

- For $f(v) - v \neq 0$ $\int \frac{dv}{f(v) - v} = \int \frac{dx}{x}$
- What happens when $f(v) - v = 0$? $v' = 0$ The equation has singular solutions of the form $y = cx$, $c = \text{const.}$

Homogeneous Polar Equations: Example 1

Consider the IVP

$$xy' = x + 2y, \quad y(1) = 3$$

Is it a homogeneous polar equation?

Homogeneous Polar Equations: Example 1

Consider the IVP

$$xy' = x + 2y, \quad y(1) = 3$$

Is it a homogeneous polar equation? Yes!

$$y' = 1 + 2\frac{y}{x}$$

Then $f(v) = 1 + 2v$ and $f(v) - v = v + 1$. For $v \neq -1$,

$$-1, \quad \int \frac{dv}{v+1} = \int \frac{dx}{x} \quad \Rightarrow \quad v + 1 = Cx, \quad C = \text{const} \neq 0.$$

$$y(x) = Cx^2 - x$$

The case $v = -1$ is equivalent to $y = -x$ and it is covered by the solution above if we allow $C = 0$. Applying the IC we obtain

$$y(x) = 4x^2 - x.$$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution?

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$.

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} =$$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined?

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes.

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE?

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow$$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow \int \frac{1 + 2v}{v} dv = \int \frac{dx}{x}$$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow \int \frac{1 + 2v}{v} dv = \int \frac{dx}{x}$$

$$2v + \ln|v| = \ln|x| + C \quad \text{or}$$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow \int \frac{1 + 2v}{v} dv = \int \frac{dx}{x}$$

$$2v + \ln|v| = \ln|x| + C \quad \text{or} \quad 2\frac{y}{x} + \ln\left|\frac{y}{x^2}\right| = C.$$

Apply the IC and find that $C =$

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow \int \frac{1 + 2v}{v} dv = \int \frac{dx}{x}$$

$$2v + \ln|v| = \ln|x| + C \quad \text{or} \quad 2\frac{y}{x} + \ln\left|\frac{y}{x^2}\right| = C.$$

Apply the IC and find that $C = 4 + \ln 2$. Therefore, the solution is

Homogeneous Polar Equations: Example 2

Consider the IVP

$$(x^2 + 2xy)y' = 2(xy + y^2), \quad y(1) = 2$$

Is there an equilibrium solution? $y = 0$. Does it satisfy the IC?

$$y' = \frac{2\frac{y}{x} + 2\left(\frac{y}{x}\right)^2}{1 + 2\frac{y}{x}} = \frac{2(v + v^2)}{1 + 2v} = f(v).$$

Is f well-defined? Yes. Does $y = -x/2$ satisfy the DE? No.

$$f(v) - v = \frac{v}{1 + 2v} \Rightarrow \int \frac{1 + 2v}{v} dv = \int \frac{dx}{x}$$

$$2v + \ln|v| = \ln|x| + C \quad \text{or} \quad 2\frac{y}{x} + \ln\left|\frac{y}{x^2}\right| = C.$$

Apply the IC and find that $C = 4 + \ln 2$. Therefore, the solution is

$$2\frac{y}{x} + \ln\left|\frac{y}{x^2}\right| = 4.$$

Homogeneous Polar Equations: Remark

A homogeneous polar equation can be written in the form

$$f_1(x, y)y' = f_2(x, y).$$

Homogeneous Polar Equations: Remark

A homogeneous polar equation can be written in the form

$$f_1(x, y)y' = f_2(x, y).$$

Let there exist a function $\bar{y}(x)$ with the property that

$$f_1(x, \bar{y}(x)) = f_2(x, \bar{y}(x)) = 0.$$

Homogeneous Polar Equations: Remark

A homogeneous polar equation can be written in the form

$$f_1(x, y)y' = f_2(x, y).$$

Let there exist a function $\bar{y}(x)$ with the property that

$$f_1(x, \bar{y}(x)) = f_2(x, \bar{y}(x)) = 0.$$

Is $\bar{y}(x)$ a solution?

Homogeneous Polar Equations: Remark

A homogeneous polar equation can be written in the form

$$f_1(x, y)y' = f_2(x, y).$$

Let there exist a function $\bar{y}(x)$ with the property that

$$f_1(x, \bar{y}(x)) = f_2(x, \bar{y}(x)) = 0.$$

Is $\bar{y}(x)$ a solution?

This type of 'solution' would be caused by the algebraic structure of the equation and not by its differential nature. In general, correctly formulated mathematical models are not expected to exhibit such anomalies.

Homogeneous Polar Equations: Remark

Consider the IVP

$$(xy - 3x^2)y' = 2y^2 - 5xy - 3x^2, \quad y(1) = 3.$$

Then

$$f_1(x, y) = xy - 3x^2 = x(y - 3x),$$

$$f_2(x, y) = 2y^2 - 5xy - 3x^2 = (x + 2y)(y - 3x),$$

and the function $y(x) = 3x$ satisfies it. We need to clean the DE algebraically, that is, assume that $y \neq 3x$ and divide both parts by $y - 3x$. We obtain the IVP $xy' = x + 2y$, $y(1) = 3$ with the solution $y = 4x^2 - x$.

The discarded function $y(x) = 3x$ also satisfies the prescribed IC, so, at first glance, it would appear that the IVP does not have a unique solution. This situation is unacceptable in mathematical modeling, where $y(x) = 3x$ is normally considered a spurious 'solution' and ignored.

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation?

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.
- Compute the derivative y' .

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.
- Compute the derivative y' .

$$\frac{1}{1-n} w^{n/(1-n)} w' + p w^{1/(1-n)} = q w^{n/(1-n)}.$$

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.
- Compute the derivative y' .

$$\frac{1}{1-n} w^{n/(1-n)} w' + p w^{1/(1-n)} = q w^{n/(1-n)}.$$

- Divide by $w^{n/(1-n)}$ and multiply by $1-n$ to obtain

$$w' + (1-n)pw = (1-n)q.$$

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.
- Compute the derivative y' .

$$\frac{1}{1-n} w^{n/(1-n)} w' + p w^{1/(1-n)} = q w^{n/(1-n)}.$$

- Divide by $w^{n/(1-n)}$ and multiply by $1-n$ to obtain

$$w' + (1-n)pw = (1-n)q.$$

Is it a linear equation?

Bernoulli Equations

The general form of a Bernoulli equation is

$$y' + p(t)y = q(t)y^n, \quad n \neq 1.$$

Is it a linear equation? No.

- Make the substitution $y(t) = (w(t))^{1/(1-n)}$.
- Compute the derivative y' .

$$\frac{1}{1-n} w^{n/(1-n)} w' + p w^{1/(1-n)} = q w^{n/(1-n)}.$$

- Divide by $w^{n/(1-n)}$ and multiply by $1-n$ to obtain

$$w' + (1-n)pw = (1-n)q.$$

Is it a linear equation? Yes. \Rightarrow Apply the known methods to solve it.

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, y(0) = -1$$

Is it a Bernoulli equation?

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, y(0) = -1$$

Is it a Bernoulli equation? Yes, $n = 2$, $p(t) = 3$, $q(t) = -6$.

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, y(0) = -1$$

Is it a Bernoulli equation? Yes, $n = 2$, $p(t) = 3$, $q(t) = -6$.

$$y = w^{-1} \Rightarrow y' = -w^{-2}w', w(0) = (y(0))^{-1} = -1.$$

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, \quad y(0) = -1$$

Is it a Bernoulli equation? Yes, $n = 2$, $p(t) = 3$, $q(t) = -6$.

$$y = w^{-1} \quad \Rightarrow \quad y' = -w^{-2}w', \quad w(0) = (y(0))^{-1} = -1.$$

We obtain the IVP

$$w' - 3w = 6, \quad w(0) = -1.$$

The solution is $w(t) = e^{3t} - 2$.

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, \quad y(0) = -1$$

Is it a Bernoulli equation? Yes, $n = 2$, $p(t) = 3$, $q(t) = -6$.

$$y = w^{-1} \Rightarrow y' = -w^{-2}w', \quad w(0) = (y(0))^{-1} = -1.$$

We obtain the IVP

$$w' - 3w = 6, \quad w(0) = -1.$$

The solution is $w(t) = e^{3t} - 2$.

What is the solution of the IVP for the Bernoulli equation?

Bernoulli Equations: Example 1

Consider the IVP

$$y' + 3y + 6y^2 = 0, \quad y(0) = -1$$

Is it a Bernoulli equation? Yes, $n = 2$, $p(t) = 3$, $q(t) = -6$.

$$y = w^{-1} \Rightarrow y' = -w^{-2}w', \quad w(0) = (y(0))^{-1} = -1.$$

We obtain the IVP

$$w' - 3w = 6, \quad w(0) = -1.$$

The solution is $w(t) = e^{3t} - 2$.

What is the solution of the IVP for the Bernoulli equation?

$$y(t) = (w(t))^{-1} = \frac{1}{e^{3t} - 2}.$$

Bernoulli Equations: Example 2

For the DE in the IVP

$$ty' + 8y = 12t^2\sqrt{y}, \quad y(1) = 16$$

we have

$$p(t) = 8/t, \quad q(t) = 12t, \quad n = 1/2.$$

Bernoulli Equations: Example 2

For the DE in the IVP

$$ty' + 8y = 12t^2\sqrt{y}, \quad y(1) = 16$$

we have

$$p(t) = 8/t, \quad q(t) = 12t, \quad n = 1/2.$$

What is the substitution?

Bernoulli Equations: Example 2

For the DE in the IVP

$$ty' + 8y = 12t^2\sqrt{y}, \quad y(1) = 16$$

we have

$$p(t) = 8/t, \quad q(t) = 12t, \quad n = 1/2.$$

What is the substitution? $y = w^2$.

Bernoulli Equations: Example 2

For the DE in the IVP

$$ty' + 8y = 12t^2\sqrt{y}, \quad y(1) = 16$$

we have

$$p(t) = 8/t, \quad q(t) = 12t, \quad n = 1/2.$$

What is the substitution? $y = w^2$. The new IVP is

$$tw' + 4w = 6t^2, \quad w(1) = 4.$$

Bernoulli Equations: Example 2

For the DE in the IVP

$$ty' + 8y = 12t^2\sqrt{y}, \quad y(1) = 16$$

we have

$$p(t) = 8/t, \quad q(t) = 12t, \quad n = 1/2.$$

What is the substitution? $y = w^2$. The new IVP is

$$tw' + 4w = 6t^2, \quad w(1) = 4.$$

Exercise: Solve this IVP and the IVP for the Bernoulli equation.

Answer: $y(t) = (t^2 + 3t^{-4})^2$.

Riccati Equations

The general form of a Riccati equation is

$$y' = q_0(t) + q_1(t)y + q_2(t)y^2,$$

where q_0 , q_1 , and q_2 are given functions, with q_0 , $q_2 \neq 0$.

Riccati Equations

The general form of a Riccati equation is

$$y' = q_0(t) + q_1(t)y + q_2(t)y^2,$$

where q_0 , q_1 , and q_2 are given functions, with q_0 , $q_2 \neq 0$.

- Assume that we know a particular solution y_1 of the R. equation.

Riccati Equations

The general form of a Riccati equation is

$$y' = q_0(t) + q_1(t)y + q_2(t)y^2,$$

where q_0 , q_1 , and q_2 are given functions, with q_0 , $q_2 \neq 0$.

- Assume that we know a particular solution y_1 of the R. equation.
- Make the substitution $y = y_1 + \frac{1}{w}$ to obtain

$$y' = y_1' - \frac{w'}{w^2} = q_0 + q_1(y_1 + \frac{1}{w}) + q_2(y_1 + \frac{1}{w})^2. \text{ What is } y_1'?$$

Riccati Equations

The general form of a Riccati equation is

$$y' = q_0(t) + q_1(t)y + q_2(t)y^2,$$

where q_0 , q_1 , and q_2 are given functions, with q_0 , $q_2 \neq 0$.

- Assume that we know a particular solution y_1 of the R. equation.
- Make the substitution $y = y_1 + \frac{1}{w}$ to obtain

$$y' = y_1' - \frac{w'}{w^2} = q_0 + q_1(y_1 + \frac{1}{w}) + q_2(y_1 + \frac{1}{w})^2. \text{ What is } y_1'?$$

- $q_0(t) + q_1(t)y_1 + q_2(t)y_1^2 - \frac{w'}{w^2} = q_0(t) + q_1(t)y_1 +$
 $+ \frac{q_1}{w} + q_2y_1^2 + 2\frac{q_2y_1}{w} + \frac{q_2}{w^2}$

Riccati Equations

The general form of a Riccati equation is

$$y' = q_0(t) + q_1(t)y + q_2(t)y^2,$$

where q_0 , q_1 , and q_2 are given functions, with q_0 , $q_2 \neq 0$.

- Assume that we know a particular solution y_1 of the R. equation.
- Make the substitution $y = y_1 + \frac{1}{w}$ to obtain

$$y' = y_1' - \frac{w'}{w^2} = q_0 + q_1(y_1 + \frac{1}{w}) + q_2(y_1 + \frac{1}{w})^2. \text{ What is } y_1'?$$

- $q_0(t) + q_1(t)y_1 + q_2(t)y_1^2 - \frac{w'}{w^2} = q_0(t) + q_1(t)y_1 + \frac{q_1}{w} + q_2y_1^2 + 2\frac{q_2y_1}{w} + \frac{q_2}{w^2}$

We obtain the linear equation $w' + (q_1 + 2q_2y_1)w = -q_2$.

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ?

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.
Guess a particular solution.

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2, q_1 = 2(t^{-1} + t), q_2 = -1$.
Guess a particular solution. $y_1(t) = t$.

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.

Guess a particular solution. $y_1(t) = t$. Then

$$w' + 2t^{-1}w = 1.$$

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.

Guess a particular solution. $y_1(t) = t$. Then

$$w' + 2t^{-1}w = 1.$$

The solution is $w(t) = \frac{1}{3}t + Ct^{-2}$, $C = \text{const.}$

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.

Guess a particular solution. $y_1(t) = t$. Then

$$w' + 2t^{-1}w = 1.$$

The solution is $w(t) = \frac{1}{3}t + Ct^{-2}$, $C = \text{const.}$ Why?

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.

Guess a particular solution. $y_1(t) = t$. Then

$$w' + 2t^{-1}w = 1.$$

The solution is $w(t) = \frac{1}{3}t + Ct^{-2}$, $C = \text{const.}$ Why?

$$y = y_1 + \frac{1}{w} = t + \frac{3t^2}{t^3 + 3C}.$$

Riccati Equations: Example 1

Consider the IVP for the Riccati equation

$$y' = -1 - t^2 + 2(t^{-1} + t)y - y^2, \quad y(1) = 10/7$$

What is q_0, q_1, q_2 ? $q_0 = -1 - t^2$, $q_1 = 2(t^{-1} + t)$, $q_2 = -1$.
Guess a particular solution. $y_1(t) = t$. Then

$$w' + 2t^{-1}w = 1.$$

The solution is $w(t) = \frac{1}{3}t + Ct^{-2}$, $C = \text{const.}$ Why?

$$y = y_1 + \frac{1}{w} = t + \frac{3t^2}{t^3 + 3C}.$$

$$y(1) = 10/7 \quad \Rightarrow \quad C = 2 \quad \Rightarrow \quad y(t) = \frac{t^4 + 3t^2 + 6t}{t^3 + 6}.$$

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution $y_1(t) = \cos t$.

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution $y_1(t) = \cos t$. Then substitution is $y = \cos t + 1/w$, and the linear first-order equation takes the form

$$w' - (\tan t)w = \sec t.$$

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution $y_1(t) = \cos t$. Then substitution is $y = \cos t + 1/w$, and the linear first-order equation takes the form

$$w' - (\tan t)w = \sec t.$$

What is its general solution?

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution $y_1(t) = \cos t$. Then substitution is $y = \cos t + 1/w$, and the linear first-order equation takes the form

$$w' - (\tan t)w = \sec t.$$

What is its general solution? $w(t) = \frac{t + C}{\cos t}.$

Riccati Equations: Example 2

The IVP

$$y' = -\cos t + (2 - \tan t)y - (\sec t)y^2, \quad y(0) = 0$$

admits the particular solution $y_1(t) = \cos t$. Then substitution is $y = \cos t + 1/w$, and the linear first-order equation takes the form

$$w' - (\tan t)w = \sec t.$$

What is its general solution? $w(t) = \frac{t + C}{\cos t}$.

Therefore,

$$y(t) = \left(1 + \frac{1}{t + C}\right) \cos t, \quad C = -1, \quad y(t) = \frac{t \cos t}{t - 1}.$$

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

- If such a function $f(x, y)$ exists then the exact equation has the form $df(x, y) = 0$ with the general solution $f(x, y) = C$, $C = \text{const.}$

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

- If such a function $f(x, y)$ exists then the exact equation has the form $df(x, y) = 0$ with the general solution $f(x, y) = C$, $C = \text{const.}$
- If $f(x, y)$ exists then $df = f_x dx + f_y dy$ and hence,

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

- If such a function $f(x, y)$ exists then the exact equation has the form $df(x, y) = 0$ with the general solution $f(x, y) = C$, $C = \text{const.}$
- If $f(x, y)$ exists then $df = f_x dx + f_y dy$ and hence,
- $f_x = P$, $f_y = Q$.

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

- If such a function $f(x, y)$ exists then the exact equation has the form $df(x, y) = 0$ with the general solution $f(x, y) = C$, $C = \text{const.}$
- If $f(x, y)$ exists then $df = f_x dx + f_y dy$ and hence,
- $f_x = P$, $f_y = Q$.
- $P_y = Q_x \iff$ this is the necessary and sufficient condition for the equation to be exact.

Exact Equations

The DE

$$P(x, y) + Q(x, y)y' = 0$$

is called an **exact equation** when $Pdx + Qdy$ is the differential of a function $f(x, y)$.

- If such a function $f(x, y)$ exists then the exact equation has the form $df(x, y) = 0$ with the general solution $f(x, y) = C$, $C = \text{const.}$
- If $f(x, y)$ exists then $df = f_x dx + f_y dy$ and hence,
- $f_x = P$, $f_y = Q$.
- $P_y = Q_x \iff$ this is the necessary and sufficient condition for the equation to be exact.

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$P(x, y)$, $Q(x, y)$?

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$$P(x, y), Q(x, y)? \quad P(x, y) = y^2 - 4xy^3 + 2, \quad Q(x, y) = 2xy - 6x^2y^2$$

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$P(x, y), Q(x, y)?$ $P(x, y) = y^2 - 4xy^3 + 2, Q(x, y) = 2xy - 6x^2y^2$

Is the equation exact?

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$P(x, y), Q(x, y)?$ $P(x, y) = y^2 - 4xy^3 + 2, Q(x, y) = 2xy - 6x^2y^2$

Is the equation exact? $P_y = 2y - 12xy^2 = Q_x \Rightarrow$ the equation is exact

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$$P(x, y), Q(x, y)? \quad P(x, y) = y^2 - 4xy^3 + 2, \quad Q(x, y) = 2xy - 6x^2y^2$$

Is the equation exact? $P_y = 2y - 12xy^2 = Q_x \Rightarrow$ the equation is exact

Then there is a function $f = f(x, y)$ such that

$$f_x(x, y) = P(x, y) = y^2 - 4xy^3 + 2,$$

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$$P(x, y), Q(x, y)? \quad P(x, y) = y^2 - 4xy^3 + 2, \quad Q(x, y) = 2xy - 6x^2y^2$$

Is the equation exact? $P_y = 2y - 12xy^2 = Q_x \Rightarrow$ the equation is exact

Then there is a function $f = f(x, y)$ such that

$$f_x(x, y) = P(x, y) = y^2 - 4xy^3 + 2, \quad f_y(x, y) = Q(x, y) = 2xy - 6x^2y^2.$$

Exact Equations: Example

Consider the IVP

$$y^2 - 4xy^3 + 2 + (2xy - 6x^2y^2)y' = 0, \quad y(1) = 1$$

$$P(x, y), Q(x, y)? \quad P(x, y) = y^2 - 4xy^3 + 2, \quad Q(x, y) = 2xy - 6x^2y^2$$

Is the equation exact? $P_y = 2y - 12xy^2 = Q_x \Rightarrow$ the equation is exact

Then there is a function $f = f(x, y)$ such that

$$f_x(x, y) = P(x, y) = y^2 - 4xy^3 + 2, \quad f_y(x, y) = Q(x, y) = 2xy - 6x^2y^2.$$

We can integrate either A. the first equation with respect to x or
B. the second equation with respect to y .

Exact Equations: Example

A.

$$f(x, y) = \int f_x(x, y) dx =$$

Exact Equations: Example

A.

$$f(x, y) = \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx =$$

Exact Equations: Example

A.

$$\begin{aligned}f(x, y) &= \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx = \\&= xy^2 - 2x^2y^3 + 2x + g(y),\end{aligned}$$

where $g(y)$ is an arbitrary function of y .

Exact Equations: Example

A.

$$\begin{aligned} f(x, y) &= \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx = \\ &= xy^2 - 2x^2y^3 + 2x + g(y), \end{aligned}$$

where $g(y)$ is an arbitrary function of y .

$$f_y(x, y) = 2xy - 6x^2y^2 + g'(y) = 2xy - 6x^2y^2 \quad \Rightarrow$$

Exact Equations: Example

A.

$$\begin{aligned} f(x, y) &= \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx = \\ &= xy^2 - 2x^2y^3 + 2x + g(y), \end{aligned}$$

where $g(y)$ is an arbitrary function of y .

$$f_y(x, y) = 2xy - 6x^2y^2 + g'(y) = 2xy - 6x^2y^2 \quad \Rightarrow g = \text{Const.}$$

Exact Equations: Example

A.

$$\begin{aligned}f(x, y) &= \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx = \\&= xy^2 - 2x^2y^3 + 2x + g(y),\end{aligned}$$

where $g(y)$ is an arbitrary function of y .

$$f_y(x, y) = 2xy - 6x^2y^2 + g'(y) = 2xy - 6x^2y^2 \Rightarrow g = \text{Const.}$$

The general solution has the form $xy^2 - 2x^2y^3 + 2x = C$.

What is the value of C ?

Exact Equations: Example

A.

$$\begin{aligned}f(x, y) &= \int f_x(x, y) dx = \int (y^2 - 4xy^3 + 2) dx = \\&= xy^2 - 2x^2y^3 + 2x + g(y),\end{aligned}$$

where $g(y)$ is an arbitrary function of y .

$$f_y(x, y) = 2xy - 6x^2y^2 + g'(y) = 2xy - 6x^2y^2 \Rightarrow g = \text{Const.}$$

The general solution has the form $xy^2 - 2x^2y^3 + 2x = C$.

What is the value of C ? $C = 1$.

$$xy^2 - 2x^2y^3 + 2x = 1$$

Exercise: Try the alternative B.

Exercises

Solve the following IVP:

① $6xy^{-1} + 8x^{-3}y^3 + (4y - 3x^2y^{-2} - 12x^{-2}y^2)y' = 0, y(1) = 1/2$

Ans: $3x^2y^{-1} - 4x^{-2}y^3 + 2y^2 = 6$

② $x \sin(2y) - 3x^2 + (y + x^2 \cos(2y))y' = 0, y(1) = \pi$

Ans: $y^2 + x^2 \sin 2y - 2x^3 = \pi^2 - 2$

③ $y' + y = -y^3, y(0) = 1$ Ans: $y(t) = (2e^{2t} - 1)^{-1/2}$

④ $y' - 3y = -e^{-4t}y^2, y(0) = -1/2$

⑤ $xy' = 3y - x, y(1) = 1$ Ans: $y(x) = (x^3 + x)/2$

⑥ $(x + y)y' = 2x - y, y(1) = -1 - \sqrt{2}$

⑦ $y' = t^{-2} + 3t^{-1} - (4t^{-1} + 3)y + 2y^2, y(1) = 5/2, y_1(t) = 1/t$

Ans: $y(t) = (3t + 2)/(2t)$

⑧ $y' = 2 - 4t - 4t^2e^{-t} + (2 + 4te^{-t})ye^{-t}y^2, y(1) = 2 + e, y_1(t) = 2t$