

ECG Modeling for Simulation of Arrhythmias in Time-Varying Conditions: Default Parameter Values

The default value for the ECG simulator parameters that are introduced in the manuscript are listed here. Where not specified otherwise, the simulator parameters introduced in the previous version of the simulator are left unchanged. Parameters provided with ranges are uniformly distributed and a new parameter is sampled on a beat basis where not specified otherwise. Additionally, the probability of APB type, VPB type, and bigeminy over trigeminy (involving, respectively, p_{APBi} , p_{VPBi} and p_B) is sampled at each transition of the Markov chain to the respective AT, VPB and BT states.

Atrial premature beats

| Parameter | Value | Meaning |
|------------------|--------------|--|
| $\beta_{APB1,p}$ | [0.55, 0.95] | RR interval prematurity factor of APB with sinus node reset |
| $\beta_{APB2,p}$ | [0.55, 0.95] | RR interval prematurity factor of APB with delayed sinus node reset |
| $\beta_{APB2,f}$ | [1.10, 1.35] | RR interval compensatory pause factor of APB with delayed sinus node reset |
| $\beta_{APB3,p}$ | [0.55, 0.95] | RR interval prematurity factor of APB with compensatory pause |
| $\beta_{APB4,p}$ | [0.55, 0.65] | RR interval prematurity factor of interpolated APB |
| p_{APB1} | 0.3 | Probability of APB with sinus node reset |
| p_{APB2} | 0.3 | Probability of APB with delayed sinus node reset |
| p_{APB3} | 0.3 | Probability of APB with compensatory pause |

| | | |
|-------------------|-----|---------------------------------|
| p_{APB4} | 0.1 | Probability of interpolated APB |
|-------------------|-----|---------------------------------|

Atrial tachycardia

| Parameter | Value | Meaning |
|------------------------|----------------|--|
| p_{APB} | 0.95 | Probability of APBs in AT episode duration PMF |
| p_{AC} | 0.040 | Probability of atrial couplets in AT episode duration PMF |
| b_{AT} | 1.00 | Exponential decay in AT episode duration PMF for $l \geq 3$ |
| $\beta_{\text{AT,p}}$ | [0.55, 0.95] | RR interval prematurity factor of AT episode, sampled only at AT episode onset |
| $\beta_{\text{AT,f}}$ | [0.70, 1.50] | RR interval compensatory pause factor for AT episode, sampled only at AT episode end |
| β_{AT} | [0.5, 0.91] | RR interval shortening factor for RR intervals inside the AT episode, sampled only at AT episode onset |
| Δd_{RR} | [-100, 100] ms | RR interval variability factor for RR intervals inside the AT episode |

Ventricular premature beats

| Parameter | Value | Meaning |
|-------------------------|--------------|--|
| $\beta_{\text{VPB1,p}}$ | [0.55, 0.9] | RR interval prematurity factor of VPB with full compensatory pause |
| $\beta_{\text{VPB2,p}}$ | [0.55, 0.9] | RR interval prematurity factor of VPB with noncompensatory pause |
| $\beta_{\text{VPB3,p}}$ | [0.55, 0.65] | RR interval prematurity factor of interpolated VPB |
| p_{VPB1} | 0.47 | Probability of VPB with full compensatory pause |
| p_{VPB2} | 0.47 | Probability of VPB with noncompensatory pause |
| p_{VPB3} | 0.06 | Probability of interpolated VPB |

Bigeminy and trigeminy

| Parameter | Value | Meaning |
|----------------|-------------|--|
| p_B | 0.72 | Probability of bigeminy over trigeminy |
| a_{BT} | 0.84 | Exponential amplitude factor in BT episode duration PMF |
| $\beta_{BT,p}$ | [0.55, 0.9] | RR interval prematurity factor of ventricular beat during bigeminy or trigeminy |
| $\beta_{BT,f}$ | [1.1, 1.3] | RR interval compensatory pause factor of ventricular beat during bigeminy or trigeminy |

Atrial fibrillation

| Parameter | Value | Meaning |
|-----------|-------|--|
| b_{AF} | 0.012 | Exponential decay in AF episode duration PMF |

Muscular noise

| Parameter | Value | Meaning |
|-------------------|----------------|---|
| ν | [0.99, 0.9995] | Filter parameter which models the standard deviation of the muscular noise $w(n)$ |
| σ_w^2 | 10 μV | Variance of variation of the simulated muscular noise $x_{\sigma_w}(n)$ |
| $m_{\sigma_w}(n)$ | 20 μV | The muscular noise intensity |
| $\sigma_{w,min}$ | 0 μV | Minimum value of the muscular noise standard deviation $w(n)$ |

Motion artifacts

| Parameter | Value | Meaning |
|------------|----------------|--|
| N | 2000 ms | Length of the impulse response $h(n)$ |
| α_1 | [0.7, 99] | Coefficient of the exponentially increasing part of the impulse response |
| α_2 | [0.7, 99] | Coefficient of the exponentially decreasing part of the impulse response |
| K | [500, 1500] ms | Impulse response turning point |

Respiratory cycle

| Parameter | Value | Meaning |
|-----------------------------|-------------|---|
| $\gamma_{in} (\gamma_{ex})$ | 3 | Inspiration/expiration steepness during the respiratory cycle |
| δ_{in} | [1.8 2.2] s | Inspiration approximate duration |
| δ_{ex} | [6.3 7.7] s | Expiration approximate duration |
| $\alpha_{o,p}$ | [0.9 1.1] | Amplitude of the template respiratory cycle |
| T_r | 10 s | Template respiratory cycle duration |
| ξ_o | 5 degrees | Angular signal $\varphi(t)$ maximum variation |