Tutorium 8th Session

Long lived comet:

Hammersley and Handscomb (1964) presents a Monte Carlo calculation for the lifetime of a long lived comet. In their model, a comet with energy x < 0 takes time $(-x)^{-\frac{3}{2}}$ to complete one orbit, while one with x > 0 just leaves the solar system. Due to gravitational interaction with planets, the energy level changes randomly. Their model is that x changes to x + Z, where $Z \sim \mathcal{N}(0, \sigma^2)$. By choosing the units of energy appropriately, they can use $\sigma = 1$. A comet that starts with energy x_0 will stay in the solar system for time

$$T = \sum_{j=0}^{m-1} (-x_j)^{-\frac{2}{3}},$$

where $x_{j+1} = x_j + z_j$ and $m = \min\{j|x_j > 0\}$ indexes the first orbit to obtain a positive energy. Because the number of m of orbits to count is itself random it is hard to study the distribution of T analytically. However, the quantity T is well suited to Monte Carlo sampling and we can estimate $\mathbb{P}(T < t)$, i.e. the cumulative distribution function.

Consider a sample of N = 10'000 comets starting with $x_0 = -1$ and continuing until they either get positive energy (leaving the solar system) or complete 10^5 orbits (to make the problem computationally tractable).

- Compute the mean and the standard deviation of the sample. Repeat the experiments more times. What do you observe?
- Plot the survival distribution function, i.e. $\mathbb{P}(T > t)$.
- Compute the Chebyshev confidence interval for the probability that a comet will make more than 10^k , for $k \in \{1, 2, 3, 4\}$.
- For those comets observed to complete more that 10⁴ orbits, show the histogram of the number of orbits observed.
- For those comets observed to leave the solar system, show the histogram of the number of orbits observed.

Hammersley, J. M. (1961). On the statistical loss of long period comets from the solar system, ii. In Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, volume 3, pages 17–78.

Hammersley, J. M. and Handscomb, D. C. (1964). *Monte Carlo methods*. Methuen, London.