

Ordinary Differential Equations

Lorenzo Niccoli

Contents

1	Solving Ordinary Differential Equations	2
1.1	Euler algorithm	3
2	Higher-Order Methods	3
2.1	Order of integration methods	3

Warnings

1. This document is a draft version and may contain errors or incomplete information.
2. This is not intended to be a comprehensive treatment of the topic, but rather a brief introduction to some basic concepts and methods. The focus is on practical implementation and numerical methods rather than rigorous mathematical theory.

1 Solving Ordinary Differential Equations

The first set of problems we address are ordinary differential equation of the form:

$$\frac{dy}{dt} = f(y, t) \quad y(t_0) = y_0 \quad (1)$$

where y is the unknown function, t is the independent variable, and $f(y, t)$ is a given function. The initial condition $y(t_0) = y_0$ specifies the value of the solution at time t_0 .

This is a quite general formulation that includes many problems of interest in science and engineering. This equation models, for example, simple physical problems such as radiocative decay. As an example, consider the following differential equation:

$$\frac{dy}{dt} = -ky \quad y(0) = y_0 \quad (2)$$

where k is a positive constant. This equation describes an exponential decay process, such as the cooling of a hot object in a colder environment (Newton's law of cooling). The analytical solution to this equation is given by:

$$y(t) = y_0 e^{-kt} \quad (3)$$

Many physical phenomena can be modeled using ordinary differential equations. Some examples are listed in Table 1.

However, not all ODEs can be solved analytically, and numerical methods are often required to obtain approximate solutions. We will explore differnt numerical methods for solving ODEs in the following sections, starting with the simplest one, the Euler method and continuing with sligtly more advanced techniques such as the Runge-Kutta methods.

Table 1: Some physical phenomena modeled by ordinary differential equations (ODEs).

Physical phenomena	ODEs
Radioactive decay	$\frac{dN}{dt} = -\lambda N$
Exponential growth	$\frac{dy}{dt} = ky$
Newton cooling law	$\frac{dT}{dt} = -k(T - T_{\text{amb}})$
Charge/Discharge plate (RC)	$\frac{dV}{dt} = \frac{1}{RC}(V_0 - V)$
Logistic growth	$\frac{dP}{dt} = rP(1 - P/K)$
First order kinetics	$\frac{d[A]}{dt} = -k[A]$

1.1 Euler algorithm

The simplest numerical method for solving ordinary differential equations is the Euler method. It is based on the idea of approximating the solution by a series of small steps, using a Taylor expansion up to the first order.

$$y(t_0 + \Delta t) \approx y(t_0) + \Delta t \frac{dy}{dt} = y_0 + \Delta t f(y_0, t_0) + O(\Delta t^2) \quad (4)$$

Iterating this equation and introducing a discrete time variable $t_n = t_0 + n\Delta t$, we obtain the Euler algorithm:

$$y_{n+1} = y_n + \Delta t f(y_n, t_n) + O(\Delta t^2) \quad (5)$$

In contrast to the analytical solution which was easy for the simple example given above, the Euler method retains its simplicity nomatter which differential equations it is applied to. In this repository, there is implemented a simple C++ code that uses the Euler method to solve Newton's law of cooling. The code is in the folder `newton_eq_euler`.

2 Higher-Order Methods

2.1 Order of integration methods

The truncation of the Taylor expansion in the Euler method leads to a local error of order $O(\Delta t^2)$ per step, which accumulates to a global error of order $O(\Delta t)$ over the entire integration interval. In any simulation we need to control this error in order to obtain results which is not influenced by the time interval Δt .

There are two options to reduce the error: either decrease the time step Δt or use a **higher-order integration method**.

A method which introduces an error of order $O(\Delta t^n)$ per single step is said to be locally of n-th order. Iterating a locally n-th order method over N steps leads to a global error of order $O(\Delta t^{n-1})$.

The total error over the time T is then:

$$\text{Error} \propto N \Delta t^n = \frac{T}{\Delta t} \Delta t^n = T \Delta t^{n-1} \quad (6)$$

and the method is said to be globally of (n-1)-th order.

For example, the Euler method is locally of the second order and globally of the first order.