

Power Spectrum Maths

Lorne Whiteway, UCL

7 January 2019

1 Introduction

Notes concerning the calculation of the 2D power spectrum of a projected field given the 3D power spectrum of the unprojected field.

2 Preliminaries

Let \mathbf{x} denote a vector in three-space, x its corresponding length, and $\hat{\mathbf{x}}$ the corresponding unit length vector (i.e. point on S^2).

We deal with random variables on 3-space or on the sphere. Let angle brackets $\langle \rangle$ denote the (ensemble) average i.e. the expectation of the corresponding random variable. All random variables are statistically homogeneous and isotropic, so their expectation values cannot depend on position or direction.

2.1 3D spectral analysis

Conventions for Fourier representation of functions defined on 3-space:

$$f(\mathbf{k}) = (2\pi)^{-3/2} A^{-1} \int f(\mathbf{x}) \exp(-i\mathbf{k} \cdot \mathbf{x}) d^3x \quad (1)$$

and

$$f(\mathbf{x}) = (2\pi)^{-3/2} A \int f(\mathbf{k}) \exp(i\mathbf{k} \cdot \mathbf{x}) d^3k. \quad (2)$$

Here A is an arbitrary constant that defines the Fourier convention being used. The BOSS paper ([LMA⁺19]) uses $A = (2\pi)^{-3/2}$. See [MVH97] Appendix A for an alternative, more general parameterisation of Fourier conventions.

For a given random field f consider $\langle f(\mathbf{k}) f^*(\mathbf{k}') \rangle$. Because of statistical homogeneity and isotropy this will vanish unless $\mathbf{k} = \mathbf{k}'$ and in this case it will depend only on k . Thus we define the power spectrum P via

$$P(k) = A^2 \langle |f(\mathbf{k})|^2 \rangle \quad (3)$$

2.2 Spherical harmonic functions

Normalisation for the spherical harmonic functions Y_ℓ^m :

$$\int Y_\ell^m(\hat{\mathbf{n}}) Y_{\ell'}^{m'*}(\hat{\mathbf{n}}) d^2n = \delta_{\ell\ell'} \delta_{mm'} \quad (4)$$

2.3 2D spectral analysis

Convention for representing functions defined on the sphere using spherical harmonics:

$$f(\hat{\mathbf{n}}) = \sum_{\ell=0}^{\infty} \sum_{m=-\ell}^{\ell} a_{\ell m} Y_{\ell}^m(\hat{\mathbf{n}}) \quad (5)$$

where

$$a_{\ell m} = \int f(\hat{\mathbf{n}}) Y_{\ell}^{m*}(\hat{\mathbf{n}}) d^2 n. \quad (6)$$

To establish equation(6), multiply equation(5) by $Y_{\ell'}^{m'*}(\hat{\mathbf{n}})$ and integrate over the sphere; then use equation(4).

Let f be a statistically isotropic field on the sphere with zero expectation everywhere and consider $\langle a_{\ell m} a_{\ell' m'}^* \rangle$. This will vanish unless $\ell = \ell'$ and $m = m'$ and in this case it will depend only on ℓ . Thus we define the angular power spectrum C via

$$C_{\ell} = \langle |a_{\ell m}|^2 \rangle. \quad (7)$$

2.4 Plane Wave Expansion

The plane wave expansion is:

$$\exp(i\mathbf{k} \cdot \mathbf{x}) = \sum_{\ell=0}^{\infty} (2\ell + 1) i^{\ell} j_{\ell}(kx) P_{\ell}(\hat{\mathbf{k}} \cdot \hat{\mathbf{x}}). \quad (8)$$

Here j_{ℓ} is a spherical Bessel function and P_{ℓ} is a Legendre polynomial. This expansion provides the link between spectral analysis in 3-space and spectral analysis on the sphere.

2.5 Spherical harmonic addition theorem

The spherical harmonic addition theorem states:

$$P_{\ell}(\hat{\mathbf{x}} \cdot \hat{\mathbf{y}}) = \frac{4\pi}{2\ell + 1} \sum_{m=-\ell}^{\ell} Y_{\ell m}(\hat{\mathbf{y}}) Y_{\ell m}^*(\hat{\mathbf{x}}) \quad (9)$$

This is a generalisation of the formula for the cosine of the sum of two angles.

3 Application to density fields

The 3D density contrast field is

$$\delta^{3D}(\mathbf{x}) = \frac{\rho(\mathbf{x}) - \bar{\rho}}{\bar{\rho}} \quad (10)$$

where ρ is the density and $\bar{\rho}$ its spatial average. This is a (homogeneous and isotropic) random field in 3 space; its expectation value is zero everywhere. What we observe is one realisation of this random field. Apply the spectral analysis (above) to define the power spectrum P of this field.

Now define the projected density contrast field to be

$$\delta^{2D}(\hat{\mathbf{n}}) = \int_0^{\infty} \phi(y) \delta^{3D}(y\hat{\mathbf{n}}) dy \quad (11)$$

where y is comoving distance and ϕ is a weight function that we assume to be normalised i.e. its integral is one. (For example if ϕ is a top-hat then it serves to pick out a particular spherical shell for the projection). Now δ^{2D} is a isotropic random field on the sphere with expectation value zero everywhere. Apply the spectral analysis (above) to define the power spectrum C of this field. How are P and C related?

We see

$$\begin{aligned}
\delta^{2D}(\hat{\mathbf{n}}) &= \int_0^\infty \phi(y) \delta^{3D}(y\hat{\mathbf{n}}) dy \\
&= (2\pi)^{-3/2} A \int_0^\infty \phi(y) \int \delta^{3D}(\mathbf{k}) \exp(i\mathbf{k} \cdot y\hat{\mathbf{n}}) d^3k dy \\
&= (2\pi)^{-3/2} A \int_0^\infty \phi(y) \int \delta^{3D}(\mathbf{k}) \sum_{\ell=0}^\infty (2\ell+1) i^\ell j_\ell(ky) P_\ell(\hat{\mathbf{k}} \cdot \hat{\mathbf{n}}) d^3k dy \\
&= (2\pi)^{-3/2} A \int_0^\infty \phi(y) \int \delta^{3D}(\mathbf{k}) \sum_{\ell=0}^\infty (2\ell+1) i^\ell j_\ell(ky) \frac{4\pi}{2\ell+1} \sum_{m=-\ell}^\ell Y_{\ell m}(\hat{\mathbf{n}}) Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k dy \\
&= \sum_{\ell=0}^\infty \sum_{m=-\ell}^\ell \left(\sqrt{2/\pi} A \int_0^\infty \phi(y) \int \delta^{3D}(\mathbf{k}) i^\ell j_\ell(ky) Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k dy \right) Y_{\ell m}(\hat{\mathbf{n}}).
\end{aligned} \tag{12}$$

Thus

$$\begin{aligned}
a_{\ell m} &= \sqrt{2/\pi} A \int_0^\infty \phi(y) \int \delta^{3D}(\mathbf{k}) i^\ell j_\ell(ky) Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k dy \\
&= \sqrt{2/\pi} A \int \left(\int_0^\infty \phi(y) j_\ell(ky) dy \right) \delta^{3D}(\mathbf{k}) i^\ell Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k \\
&= \sqrt{2/\pi} A \int W_\ell(k) \delta^{3D}(\mathbf{k}) i^\ell Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k
\end{aligned} \tag{13}$$

where

$$W_\ell(k) = \int_0^\infty \phi(y) j_\ell(ky) dy. \tag{14}$$

Therefore

$$\begin{aligned}
C_\ell &= \langle |a_{\ell m}|^2 \rangle = \langle a_{\ell m} a_{\ell m}^* \rangle \\
&= \frac{2}{\pi} A^2 \left\langle \left(\int W_\ell(k) \delta^{3D}(\mathbf{k}) i^\ell Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k \right) \left(\int W_\ell(k') \delta^{3D*}(\mathbf{k}') (-i)^\ell Y_{\ell m}(\hat{\mathbf{k}}') d^3k' \right) \right\rangle \\
&= \frac{2}{\pi} A^2 \int \int W_\ell(k) W_\ell(k') \langle \delta^{3D}(\mathbf{k}) \delta^{3D*}(\mathbf{k}') \rangle Y_{\ell m}^*(\hat{\mathbf{k}}) Y_{\ell m}(\hat{\mathbf{k}}') d^3k d^3k' \\
&= \frac{2}{\pi} \int \int W_\ell(k) W_\ell(k') P(k) \delta(\mathbf{k} - \mathbf{k}') Y_{\ell m}^*(\hat{\mathbf{k}}) Y_{\ell m}(\hat{\mathbf{k}}') d^3k d^3k' \\
&= \frac{2}{\pi} \int W_\ell(k)^2 P(k) |Y_{\ell m}^*(\hat{\mathbf{k}})|^2 d^3k \\
&= \frac{2}{\pi} \int k^2 W_\ell(k)^2 P(k) dk \int |Y_{\ell m}^*(\hat{\mathbf{k}})|^2 d^2k \\
&= \frac{2}{\pi} \int k^2 W_\ell(k)^2 P(k) dk.
\end{aligned} \tag{15}$$

Define the dimensionless power spectrum via:

$$\Delta^2(k) = \frac{k^3 P(k)}{2\pi^2}; \tag{16}$$

then

$$C_\ell = 4\pi \int W_\ell(k)^2 \Delta^2(k) dk/k. \quad (17)$$

4 Cross correlations

Assume now that we have two weight functions ϕ^i and ϕ^j and hence two window functions $W_\ell^i(k)$ and $W_\ell^j(k)$. We can then define the cross-correlation

$$\begin{aligned} C_\ell^{ij} &= \langle a_{\ell m}^i a_{\ell m}^{j*} \rangle \\ &= \frac{2}{\pi} A^2 \left\langle \left(\int W_\ell^i(k) \delta^{3D}(\mathbf{k}) i^\ell Y_{\ell m}^*(\hat{\mathbf{k}}) d^3k \right) \left(\int W_\ell^j(k') \delta^{3D*}(\mathbf{k}') (-i)^\ell Y_{\ell m}(\hat{\mathbf{k}}') d^3k' \right) \right\rangle \\ &= \frac{2}{\pi} A^2 \int \int W_\ell^i(k) W_\ell^j(k') \langle \delta^{3D}(\mathbf{k}) \delta^{3D*}(\mathbf{k}') \rangle Y_{\ell m}^*(\hat{\mathbf{k}}) Y_{\ell m}(\hat{\mathbf{k}}') d^3k d^3k' \\ &= \frac{2}{\pi} \int \int W_\ell^i(k) W_\ell^j(k') P(k) \delta(\mathbf{k} - \mathbf{k}') Y_{\ell m}^*(\hat{\mathbf{k}}) Y_{\ell m}(\hat{\mathbf{k}}') d^3k d^3k' \\ &= \frac{2}{\pi} \int W_\ell^i(k) W_\ell^j(k) P(k) |Y_{\ell m}^*(\hat{\mathbf{k}})|^2 d^3k \\ &= \frac{2}{\pi} \int k^2 W_\ell^i(k) W_\ell^j(k) P(k) dk \int |Y_{\ell m}^*(\hat{\mathbf{k}})|^2 d^2k \\ &= \frac{2}{\pi} \int k^2 W_\ell^i(k) W_\ell^j(k) P(k) dk. \end{aligned} \quad (18)$$

This cross-correlations are therefore real, but not necessarily positive.

5 References

- [LMA⁺19] Arthur Loureiro, Bruno Moraes, Filipe B Abdalla, Andrei Cuceu, Michael McLeod, Lorne Whiteway, Sreekumar T Balan, Aurélien Benoit-Lévy, Ofer Lahav, Marc Manera, et al. Cosmological Measurements from Angular Power Spectra Analysis of BOSS DR12 Tomography. *Monthly Notices of the Royal Astronomical Society*, 2019.
- [MVH97] S Matarrese, Licia Verde, and AF Heavens. Large-scale bias in the universe: bispectrum method. *Monthly Notices of the Royal Astronomical Society*, 290(4):651–662, 1997.