

Concise Total Synthesis of Salimabromide

Hai-Hua Lu,^{*} Kang-Ji Gan,[†] Fu-Qiang Ni,[†] Zhihan Zhang, and Yao ZhuCite This: *J. Am. Chem. Soc.* 2022, 144, 18778–18783

Read Online

ACCESS |



Metrics & More



Article Recommendations



Supporting Information

ABSTRACT: We achieved a concise total synthesis of salimabromide by using a novel intramolecular radical cyclization to simultaneously construct the unique benzo-fused [4.3.1] carbon skeleton and the vicinal quaternary stereocenters. Other notable transformations include a tandem Michael/Mukaiyama aldol reaction to introduce most of the molecule's structural elements, along with hidden information for late-stage transformations, an intriguing tandem oxidative cyclization of a diene to form the bridged butyrolactone and enone moieties spontaneously, and a highly enantioselective hydrogenation of a cycloheptenone derivative (97% ee) that paved the way for the asymmetric synthesis of salimabromide.

In sharp contrast to terrestrial myxobacteria,¹ marine myxobacteria (*Enhygromyxa*, *Plesiocystis*, *Pseudoenhygromyxa*, *Haliangium*) have few known secondary metabolites.^{2,3} Among the known natural products originating from marine myxobacteria, salimabromide, which was discovered by König and co-workers in 2013,³ is the first to have been obtained from the *Plesiocystis/Enhygromyxa* clade of obligatory marine myxobacteria. Preliminary studies showed that it is a potent inhibitor of *Arthrobacter crystallopoietes*. Unfortunately, further evaluation of its biological activity was hampered by its extremely low natural abundance (only 0.5 mg was isolated from 64 L of culture), as well as the difficulty of cultivating marine myxobacteria compared with terrestrial myxobacteria.

Salimabromide has a unique benzo-fused [4.3.1] carbon skeleton bearing four contiguous stereocenters, including vicinal quaternary carboncenters. Interestingly, the aromatic ring is dibrominated; and the unusually bridged cycloheptenone moiety is fused to a butyrolactone ring. These structural features make salimabromide highly compact and conformationally rigid, as well as an appealing synthetic challenge. In 2015, Menche and Schmalzbauer reported the first stereoselective construction of its tricyclic core.^{4a} In 2018, Magauer and co-workers successfully completed the first racemic total synthesis of salimabromide.⁵ Soon after, Menche and co-workers achieved its first enantioselective synthesis and showcased the quite unusual almost racemic nature of natural salimabromide.^{4b,6} Intriguingly, both research groups employed an in situ ketiminium-[2+2]-cycloaddition of a tetralin-tethered amide to build the central bridged carbon framework and a Baeyer–Villiger oxidation to form the butyrolactone part (Figure 1).⁷ In contrast, we envisioned that an intramolecular radical cyclization of diene 3 mediated by hydrogen atom transfer (HAT) could be used to simultaneously build the key benzo-fused [4.3.1] carbon skeleton and the vicinal quaternary centers and that a subsequent novel tandem oxidative cyclization would complete a concise synthesis of salimabromide.

We commenced the synthesis by carrying out a tandem Michael/Mukaiyama aldol reaction⁸ of cycloheptadienone 4

(Scheme 1). To our delight, the silyl enol ether intermediate generated by the initial conjugate addition of Grignard reagent 5a⁹ with 4, reacted smoothly with 6a in the presence of zinc bromide as a Lewis acid to give desired adduct 8 in 63% yield, accompanied by a 15% yield of hydrolyzed product 7, which could easily be transformed to 8. Next, an oxidative rearrangement of the tertiary allylic alcohol¹⁰ resulting from addition of methyl lithium to cycloheptenone 8 was used to quickly access β -methyl substituted cycloheptenone 3a, which was required for the planned HAT-mediated cyclization.

Since the pioneering studies by Baran and co-workers,¹¹ the HAT-promoted Giese reaction has been thoroughly investigated^{12–14} and has found a wide array of synthetic applications because of its practicality.^{15,16} Inspired by this prior work, we subjected diene 3b to HAT conditions with various catalysts (Fe, Co, Mn, etc.)^{15a} in combination with a number of reductants (silanes)^{14a} and solvents (Table 1; for complete optimization efforts, see Supporting Information). However, the desired radical cyclization proved to be challenging: the undesired [1,5]-HAT process¹⁷ always predominated, producing a large amount of undesired alkene 9b. We attributed these results both to the difficulty of forming vicinal quaternary centers^{18,19} because of steric hindrance during the cyclization and to the proximity of the initially generated tertiary radical to the hydrogen atom at the γ -position of the enone moiety, which was supported by DFT calculations (Figure 2). The best results were eventually obtained with iron catalysis. Despite the relatively low yield of desired adduct 10a (28%), the reaction could easily be scaled up to produce gram quantities in a single operation.

Having obtained ample quantities of 10a, which contains all the hidden information required for subsequent trans-

Received: August 6, 2022

Published: October 4, 2022






Figure 1. Previous work and our retrosynthetic analysis.




Scheme 1. Total Synthesis of Salimabromide [(±)-1].^a^aSee the Supporting Information for detailed procedures and characterization data.

Figure 2. Comparison of optimized transition states leading to 9a (TS1) and 10a (TS2) from 3a based on DFT calculations.

formations,^{20,21} we moved forward to complete the synthesis of salimabromide. Dehydrogenation of 10a quickly provided


enone 11, which was reduced and dehydrated to produce diene 12 initially. It is interesting to note that the dehydration was best performed with the Burgess reagent, and we found that this reduction-dehydration process could be carried out in one pot. To obtain a better overall yield, an alternative sequence (triflation-reduction) was later developed. At this stage, the lactone ring needs to be formed. To our delight, treatment of 12 with the Jones reagent directly and efficiently afforded butyrolactone ring-fused enone product 13. Based on pioneering studies by McDonald and Towne who used hydroxy-directed alkene difunctionalizations for polyether synthesis,²² we assumed that this tandem oxidative cyclization was directed by an acid intermediate generated in situ. This hypothesis was supported by preliminary experimental studies.²³ Finally, dibromination by utilizing the Brønsted acid-catalyzed halogenation of arenes recently disclosed by Jiao

Table 1. Selected Optimization Data of the Giese–Baran Cyclization^a

entry	metal salt	silane	solvent	yield (%) ^b	
				9b	10b
1 ^c	Mn(dpm) ₃	PhSiH ₃	ⁱ PrOH	4	17
2 ^c	Co(acac) ₂	PhSiH ₃	ⁱ PrOH	ND	ND
3	Fe(acac) ₃	PhSiH ₃	ⁱ PrOH	11	22
4	Fe(acac) ₃	PhSiH ₃	EtOH	21	26
5	Fe(acac) ₃	PhSiH ₃	TFE	NR	NR
6	Fe(acac) ₃	PhSiH ₃	dioxane	33	15
7	Fe(acac) ₃	PhSiH ₃	DCE	NR	NR
8 ^d	Fe(acac) ₃	PhSiH ₂ (O ^{<i>i</i>} Pr) ₂	EtOH	4	21
9 ^d	Fe(acac) ₃	Et ₃ SiH	EtOH	NR	NR
10 ^d	Fe(acac) ₃	TMDSO	EtOH	NR	NR
11 ^d	Fe(acac) ₃	Ph ₂ SiH ₂	EtOH	17	20
12 ^e	Fe(acac) ₃	PhSiH ₃	EtOH	30	28

^aReaction conditions: **3b** (0.25 mmol), metal salt (0.3 equiv), and silane (3.0 equiv) in solvents under N₂ at 60 °C for 1–3 h. ^bDetermined by crude NMR. ^cTBHP (2.0 equiv) was added. ^dConducted at 40 °C. ^eKH₂PO₄ (2.0 equiv) was added. ND = Not Determined (messy). NR = No Reaction.

Scheme 2. Formal Enantioselective Total Synthesis of (+)-Salimabromide [(+)-1].^a



^aSee the [Supporting Information](#) for detailed procedures and characterization data and X-ray data of (+)-8 (CCDC 2205845; key: C, gray; H, white; O, red).

and co-workers was found to be superior to Magauer's protocol⁵ and provided salimabromide in a better yield.²⁴

We, then, sought to accomplish an enantioselective synthesis of salimabromide (**Scheme 2**). Initially, we attempted to realize an enantioselective version of the tandem Michael/Mukaiyama

aldol reaction of cycloheptadienone **4** to produce chiral **8**. Unfortunately, the initial conjugate addition proved extremely challenging with an aryl Grignard reagent **5** or an aluminum or boronic acid reagent, and only racemic adducts were obtainable in our hands.^{25–27} Inspired by a recent advance

achieved by Zhang and co-workers on enantioselective hydrogenation of endocyclic enones,²⁸ we turned our attention to the enantioselective hydrogenation of cycloheptenone derivative **14**. To our delight, when we used (*R,S*)-ZhaoPhos as a ligand, reduction product (–)-**15** was obtained in 86% yield with 97% ee, for which the absolute configuration was determined by single-crystal X-ray analysis of its later intermediate (+)-**8**. Subsequent Suzuki coupling and dehydrogenation provided chiral enone (–)-**7**. Its transformation to (+)-**8** thus constitutes a formal asymmetric synthesis of (+)-salimabromide. Because (*S,R*)-ZhaoPhos is also easily accessible, we expect that the same sequence could be used to obtain (–)-salimabromide from the enantiomer of (–)-**15**.

In summary, we have achieved a concise synthesis of the marine natural product salimabromide by utilizing a novel intramolecular HAT-mediated radical cyclization to simultaneously construct the unique benzo-fused [4.3.1] carbon skeleton and the vicinal quaternary centers. In addition, the discovery and application of an unprecedented acid-directed tandem oxidative cyclization to form the bridged butyrolactone and enone moieties further enhanced the overall efficiency of the synthesis. Moreover, a highly enantioselective hydrogenation of a cycloheptenone intermediate (97% ee) paved the way for an enantioselective synthesis of (+)-salimabromide and would likely permit the synthesis of its enantiomer, (–)-salimabromide. We expect that the transformations described herein will have implications for future developments in the field of organic synthesis.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.2c08337>.

Experimental procedures and characterization data for all the products (PDF)

Accession Codes

CCDC 2205845 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

■ AUTHOR INFORMATION

Corresponding Author

Hai-Hua Lu – Key Laboratory of Precise Synthesis of Functional Molecules of Zhejiang Province, Department of Chemistry, School of Science and Research Center for Industries of the Future, Westlake University, Hangzhou 310030, China; Institute of Natural Sciences, Westlake Institute for Advanced Study, Hangzhou 310024, China; orcid.org/0000-0003-3604-7145; Email: luhaihua@westlake.edu.cn

Authors

Kang-Ji Gan – Key Laboratory of Precise Synthesis of Functional Molecules of Zhejiang Province, Department of Chemistry, School of Science and Research Center for Industries of the Future, Westlake University, Hangzhou 310030, China; Institute of Natural Sciences, Westlake Institute for Advanced Study, Hangzhou 310024, China;

Department of Chemistry, Zhejiang University, Hangzhou 310058, China

Fu-Qiang Ni – Key Laboratory of Precise Synthesis of Functional Molecules of Zhejiang Province, Department of Chemistry, School of Science and Research Center for Industries of the Future, Westlake University, Hangzhou 310030, China; Institute of Natural Sciences, Westlake Institute for Advanced Study, Hangzhou 310024, China

Zhihan Zhang – CCNU-uOttawa Joint Research Centre, Key Laboratory of Pesticide & Chemical Biology, Ministry of Education, College of Chemistry, Central China Normal University, Wuhan, Hubei 430079, China

Yao Zhu – Key Laboratory of Precise Synthesis of Functional Molecules of Zhejiang Province, Department of Chemistry, School of Science and Research Center for Industries of the Future, Westlake University, Hangzhou 310030, China; Institute of Natural Sciences, Westlake Institute for Advanced Study, Hangzhou 310024, China

Complete contact information is available at: <https://pubs.acs.org/10.1021/jacs.2c08337>

Author Contributions

[†]K.J.G. and F.Q.N. contributed equally to this work.

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We are grateful to the National Natural Science Foundation of China (no. 22271236) and the Natural Science Foundation of Zhejiang Province (no. LXZ22B020001) for support of this research. We thank Mr. Menglong Zhao and Professors Jialin Wen and Xumu Zhang (Southern University of Science and Technology) for kindly providing us their (*R*, *S*)-ZhaoPhos ligand. We acknowledge the Instrumentation and Service Centre for Molecular Sciences at Westlake University for help with measurement and data interpretation.

■ ABBREVIATIONS

PCC, pyridinium chlorochromate.; LHMDs, lithium bis-(trimethylsilyl)amide.; Burgess reagent, methyl *N*-(triethylammoniosulfonyl)carbamate; Rh(*nbd*)₂BF₄, bis-(norbornadiene)rhodium(I) tetrafluoroborate.; DCE, 1,2-dichloroethane.; TFE, 2,2,2-trifluoroethanol.; HFIP, 1,1,1,3,3,3-hexafluoro-2-propanol.; DIPEA, *N,N*-diisopropylethylamine.; TMDSO, tetramethyldisiloxane.; DBDMH, 1,3-dibromo-5,5-dimethylhydantoin.; *y*, yield; *dr*, diastereomeric ratio; *ee*, enantiomeric excess

■ REFERENCES

- (1) Herrmann, J.; Fayad, A. A.; Müller, R. Natural Products from Myxobacteria: Novel Metabolites and Bioactivities. *Nat. Prod. Rep.* **2017**, *34*, 135–160.
- (2) (a) Fudouzi, R.; Iizuka, T.; Sato, S.; Ando, T.; Shimba, N.; Yamanaka, S. Haliangicin, a Novel Antifungal Metabolite Produced by a Marine Myxobacterium. *J. Antibiot.* **2001**, *54*, 153–156. (b) Felder, S.; Kehraus, S.; Neu, E.; Bierbaum, G.; Schäberle, T. F.; König, G. M. Salimyoxins and Enhygrolides: Antibiotic, Sponge-Related Metabolites from the Obligate Marine Myxobacterium *Enhygromyxa salina*. *ChemBioChem.* **2013**, *14*, 1363–1371. (c) Dávila-Céspedes, A.; Hufendiek, P.; Crüsemann, M.; Schäberle, T. F.; König, G. M. Marine-derived Myxobacteria of the Suborder Nannocystineae: An Underexplored Source of Structurally Intriguing and Biologically Active Metabolites. *Beilstein J. Org. Chem.* **2016**, *12*, 969–984.

- (3) Felder, S.; Dreisigacker, S.; Kehraus, S.; Neu, E.; Bierbaum, G.; Wright, P. R.; Menche, D.; Schäberle, T. F.; König, G. M. Salimabromide: Unexpected Chemistry from the Obligate Marine Myxobacterium *Enhygromyxa salina*. *Chem. Eur. J.* **2013**, *19*, 9319–9324.
- (4) (a) Schmalzbauer, B.; Menche, D. Concise Synthesis of the Tricyclic Core of Salimabromide. *Org. Lett.* **2015**, *17*, 2956–2959. (b) Palm, A.; Knopf, C.; Schmalzbauer, B.; Menche, D. Enantioselective Total Synthesis of (+)-Salimabromide Reveals Almost Racemic Nature of Natural Salimabromide. *Org. Lett.* **2019**, *21*, 1939–1942.
- (5) (a) Schmid, M.; Grossmann, A. S.; Wurst, K.; Magauer, T. Total Synthesis of Salimabromide: A Tetracyclic Polyketide from a Marine Myxobacterium. *J. Am. Chem. Soc.* **2018**, *140*, 8444–8447. (b) Schmid, M.; Grossmann, A. S.; Mayer, P.; Müller, T.; Magauer, T. Ring-Expansion Approaches for the Total Synthesis of Salimabromide. *Tetrahedron* **2019**, *75*, 3195–3215.
- (6) Scott, K. A.; Ropek, N.; Melillo, B.; Schreiber, S. L.; Cravatt, B. F.; Vinogradova, E. V. Stereochemical Diversity as a Source of Discovery in Chemical Biology. *Curr. Res. Chem. Biol.* **2022**, *2*, 100028.
- (7) Notably, Menche and co-workers developed an excellent one-pot allylic/Baeyer–Villiger oxidation to complete an efficient synthesis of salimabromide (ref 4b).
- (8) (a) For selected reviews, see: Kan, S. B. J.; Ng, K. K.-H.; Paterson, I. The Impact of the Mukaiyama Aldol Reaction in Total Synthesis. *Angew. Chem., Int. Ed.* **2013**, *52*, 9097–9108. (b) Matsuo, J.; Murakami, M. The Mukaiyama Aldol Reaction: 40 Years of Continuous Development. *Angew. Chem., Int. Ed.* **2013**, *52*, 9109–9118.
- (9) The corresponding 2-bromo-4-ethyl-1-propenylbenzene could be conveniently prepared in two steps, see [Supporting Information](#) (page S6).
- (10) (a) Babler, J. H.; Coghlan, M. J. A Facile Method for the Bishomologation of Ketones to α,β -Unsaturated Aldehydes: Application to the Synthesis of the Cyclohexanoid Components of the Boll Weevil Sex Attractant. *Synth. Commun.* **1976**, *6*, 469–474. (b) Dauben, W. G.; Michno, D. M. Direct Oxidation of Tertiary Allylic Alcohols. A Simple and Effective Method for Alkylative Carbonyl Transposition. *J. Org. Chem.* **1977**, *42*, 682–685.
- (11) (a) Lo, J. C.; Yabe, Y.; Baran, P. S. A Practical and Catalytic Reductive Olefin Coupling. *J. Am. Chem. Soc.* **2014**, *136*, 1304–1307. (b) Lo, J. C.; Kim, D.; Pan, C.-M.; Edwards, J. T.; Yabe, Y.; Gui, J.; Qin, T.; Gutiérrez, S.; Giacoboni, J.; Smith, M. W.; Holland, P. L.; Baran, P. S. Fe-Catalyzed C–C Bond Construction from Olefins via Radicals. *J. Am. Chem. Soc.* **2017**, *139*, 2484–2503.
- (12) Giese, B.; Meister, J. Die Addition von Kohlenwasserstoffen an Olefine Eine neue Synthetische Methode. *Chem. Ber.* **1977**, *110*, 2588–2600.
- (13) Kim, D.; Rahaman, S. M. W.; Mercado, B. Q.; Poli, R.; Holland, P. Roles of Iron Complexes in Catalytic Radical Alkene Cross-Coupling: A Computational and Mechanistic Study. *J. Am. Chem. Soc.* **2019**, *141*, 7473–7485.
- (14) (a) Obradors, C. L.; Martinez, R.; Shenvi, R. A. Ph(*i*-PrO)SiH₂: An Exceptional Reductant for Metal-Catalyzed Hydrogen Atom Transfers. *J. Am. Chem. Soc.* **2016**, *138*, 4962–4971. (b) Green, S. A.; Crossley, S. W. M.; Matos, J. L. M.; Vásquez-Céspedes, S.; Shevick, S. L.; Shenvi, R. A. The High Chemofidelity of Metal-Catalyzed Hydrogen Atom Transfer. *Acc. Chem. Res.* **2018**, *51*, 2628–2640. (c) Demoret, R. M.; Baker, M. A.; Ohtawa, M.; Chen, S.; Lam, C. C.; Khom, S.; Roberto, M.; Forli, S.; Houk, K. N.; Shenvi, R. A. Synthetic, Mechanistic, and Biological Interrogation of Ginkgo biloba Chemical Space En Route to (–)-Bilobalide. *J. Am. Chem. Soc.* **2020**, *142*, 18599–18618.
- (15) (a) For recent reviews on metal-hydride HAT reactions (and references therein), see: Crossley, S. W. M.; Martinez, R. M.; Obradors, C.; Shenvi, R. A. Mn, Fe, and Co-Catalyzed Radical Hydrofunctionalizations of Olefins. *Chem. Rev.* **2016**, *116*, 8912–9000. (b) Green, S. A.; Crossley, S. W. M.; Matos, J. L. M.; Vásquez-Céspedes, S.; Shevick, S. L.; Shenvi, R. A. The High Chemofidelity of Metal-Catalyzed Hydrogen Atom Transfer. *Acc. Chem. Res.* **2018**, *51*, 2628–2640. (c) Shevick, S. L.; Wilson, C. V.; Kotesova, S.; Kim, D.; Holland, P. L.; Shenvi, R. A. Catalytic Hydrogen Atom Transfer to Alkenes: A Roadmap for Metal Hydrides and Radicals. *Chem. Sci.* **2020**, *11*, 12401–12422. (d) Thomas, W. P.; Pronin, S. V. New Methods and Strategies in the Synthesis of Terpenoid Natural Products. *Acc. Chem. Res.* **2021**, *54*, 1347–1359. (e) Schatz, D. J.; Kuenstner, E. J.; George, D. T.; Pronin, S. V. Synthesis of Rearranged Indole Diterpenes of the Paxilline type. *Nat. Prod. Rep.* **2022**, *39*, 946–968. (f) Wu, J.; Ma, Z. Metal-Hydride Hydrogen Atom Transfer (MHAT) Reactions in Natural Product Synthesis. *Org. Chem. Front.* **2021**, *8*, 7050–7056.
- (16) (a) For selected recent applications of metal-hydride HAT reactions in natural product synthesis, see: Foy, N. J.; Pronin, S. V. Synthesis of Pleuromutilin. *J. Am. Chem. Soc.* **2022**, *144*, 10174–10179. (b) Thomas, W. P.; Pronin, S. V. A Concise Enantioselective Approach to Quassinoids. *J. Am. Chem. Soc.* **2022**, *144*, 118–122. (c) Fang, X.; Zhang, N.; Chen, S.-C.; Luo, T. Scalable Total Synthesis of (–)-Triptonide: Serendipitous Discovery of a Visible-Light-Promoted Olefin Coupling Initiated by Metal-Catalyzed Hydrogen Atom Transfer (MHAT). *J. Am. Chem. Soc.* **2022**, *144*, 2292–2300. (d) Zhao, Y.; Hu, J.; Chen, R.; Xiong, F.; Xie, H.; Ding, H. Divergent Total Syntheses of (–)-Crinipellins Facilitated by a HAT-Initiated Dowd–Beckwith Rearrangement. *J. Am. Chem. Soc.* **2022**, *144*, 2495–2500. (e) Lusi, R. F.; Sennari, G.; Sarpong, R. Total Synthesis of Nine Longiborneol Sesquiterpenoids Using a Functionalized Camphor Strategy. *Nat. Chem.* **2022**, *14*, 450–456. (f) Yu, K.; Yao, F.; Zeng, Q.; Xie, H.; Ding, H. Asymmetric Total Syntheses of (+)-Davisinol and (+)-18-Benzoyldavisinol: A HAT-Initiated Transannular Redox Radical Approach. *J. Am. Chem. Soc.* **2021**, *143*, 10576–10581. (g) Vrubliauskas, D.; Gross, B. M.; Vanderwal, C. D. Stereocontrolled Radical Bicyclizations of Oxygenated Precursors Enable Short Syntheses of Oxidized Abietane Diterpenoids. *J. Am. Chem. Soc.* **2021**, *143*, 2944–2952. (h) Chen, P.; Wang, C.; Yang, R.; Xu, H.; Wu, J.; Jiang, H.; Chen, K.; Ma, Z. Asymmetric Total Synthesis of Dankasterones A and B and Periconiastone A Through Radical Cyclization. *Angew. Chem., Int. Ed.* **2021**, *60*, 5512–5518. (i) Xin, Z.; Wang, H.; He, H.; Zhao, X.; Gao, S. Asymmetric Total Synthesis of Norzoanthamine. *Angew. Chem., Int. Ed.* **2021**, *60*, 12807–12812. (j) Ji, Y.; Xin, Z.; He, H.; Gao, S. Total Synthesis of Viridin and Viridiol. *J. Am. Chem. Soc.* **2019**, *141*, 16208–16212. (k) Xu, G.; Wu, J.; Li, L.; Lu, Y.; Li, C. Total Synthesis of (–)-Daphnezomines A and B. *J. Am. Chem. Soc.* **2020**, *142*, 15240–15245. (l) Hu, Y.; Bai, M.; Yang, Y.; Tian, J.; Zhou, Q. Rapid Access to Tetracyclic Core of Wortmannin via an Intramolecular Reductive Olefin Coupling Strategy. *Org. Lett.* **2020**, *22*, 6308–6312. (m) Hu, P. F.; Chi, H. M.; DeBacker, K. C.; Gong, X.; Keim, J. H.; Hsu, I. T.; Snyder, S. A. Quaternary-Centre-Guided Synthesis of Complex Polycyclic Terpenes. *Nature* **2019**, *569*, 703–707. (n) Liu, J.; Ma, D. A Unified Approach for the Assembly of Atisine- and Hetidine-type Diterpenoid Alkaloids: Total Syntheses of Azitine and the Proposed Structure of Navirine C. *Angew. Chem., Int. Ed.* **2018**, *57*, 6676–6680. (o) Kerschgens, I.; Rovira, A. R.; Sarpong, R. Total Synthesis of (–)-Xishacorene B from (R)-Carvone Using a C–C Activation Strategy. *J. Am. Chem. Soc.* **2018**, *140*, 9810–9813. (p) Lu, Z.; Zhang, X.; Guo, Z.; Chen, Y.; Mu, T.; Li, A. Total Synthesis of Aplysiasecosterol A. *J. Am. Chem. Soc.* **2018**, *140*, 9211–9218. (q) Xu, G.; Elkin, M.; Tantillo, D. J.; Newhouse, T. R.; Maimone, T. J. Traversing Biosynthetic Carbocation Landscapes in the Total Synthesis of Andrastin and Terretinin Meroterpenes. *Angew. Chem., Int. Ed.* **2017**, *56*, 12498–12502. (r) Deng, H.; Cao, W.; Liu, R.; Zhang, Y.; Liu, B. Asymmetric Total Synthesis of Hispidanin A. *Angew. Chem., Int. Ed.* **2017**, *56*, 5849–5852. (s) Ruider, S. A.; Sandmeier, T.; Carreira, E. M. Total Synthesis of (±)-Hippolachnin A. *Angew. Chem., Int. Ed.* **2015**, *54*, 2378–2382.
- (17) This 1,5-HAT process was supported by deuterium control experiments, and the yield of the desired cyclization product was greatly improved with the deuterated substrate (see [Supporting](#)

Information, pages S25–30; we thank one of the reviewers for this insightful idea, and are currently doing further synthetic studies). For previous interesting applications of 1,5-HAT in total syntheses, see: (a) Farney, E. P.; Feng, S. S.; Schäfers, F.; Reisman, S. E. Total Synthesis of (+)-Pleuromutilin. *J. Am. Chem. Soc.* **2018**, *140*, 1267–1270. (b) Qu, Y.; Wang, Z.; Zhang, Z.; Zhang, W.; Huang, J.; Yang, Z. Asymmetric Total Synthesis of (+)-Waihoensene. *J. Am. Chem. Soc.* **2020**, *142*, 6511–6515. (c) Yang, F.; Porco, J. A., Jr. Unified, Asymmetric Total Synthesis of the Asnovolins and Related Spiromeroterpenoids: A Fragment Coupling Approach. *J. Am. Chem. Soc.* **2022**, *144*, 12970–12978. (d) Vasilev, V. H.; Spessert, L.; Yu, K.; Maimone, T. J. Total Synthesis of Resiniferatoxin. *J. Am. Chem. Soc.* **2022**, *144*, 16332–16337. For recent inspiring applications of deuterium kinetic isotope effect in total syntheses, see: (e) Miyashita, M.; Sasaki, M.; Hattori, I.; Sakai, M.; Tanino, K. Total Synthesis of Norzoanthamine. *Science* **2004**, *305*, 495–499. (f) Quasdorf, K. W.; Hutters, A. D.; Lodewyk, M. W.; Tantillo, D. J.; Garg, N. K. Total Synthesis of Oxidized Welwitindolinones and (–)-N-Methylwelwitindolinone C Isonitrile. *J. Am. Chem. Soc.* **2012**, *134*, 1396–1399. (g) Kanda, Y.; Nakamura, H.; Umehiya, S.; Puthukanoori, R. K.; Murthy Appala, V. R.; Gaddamanugu, G. K.; Paraselli, B. R.; Baran, P. S. Two-Phase Synthesis of Taxol. *J. Am. Chem. Soc.* **2020**, *142*, 10526–10533. (h) Chen, H.; Li, Z.; Shao, P.; Yuan, H.; Chen, S.-C.; Luo, T. Total Synthesis of (+)-Mutilin: A Transannular [2 + 2] Cycloaddition/Fragmentation Approach. *J. Am. Chem. Soc.* **2022**, *144*, 15462–15467.

(18) Christofers, J.; Baro, A., Eds. *Quaternary Stereocenters: Challenges and Solutions for Organic Synthesis*; Wiley-VCH, 2005.

(19) (a) Eggert, A.; Etling, C.; Lübken, D.; Saxarra, M.; Kalesse, M. Contiguous Quaternary Carbons: A Selection of Total Syntheses. *Molecules* **2020**, *25*, 3841. (b) Xin, Z.; Wang, H.; He, H.; Gao, S. Recent Advances in the Total Synthesis of Natural Products Bearing the Contiguous All-Carbon Quaternary Stereocenters. *Tetrahedron Lett.* **2021**, *71*, 153029.

(20) Cao, M.-Y.; Ma, B.-J.; Gu, Q.-X.; Fu, B.; Lu, H.-H. Concise Enantioselective Total Synthesis of Daphenylline Enabled by an Intramolecular Oxidative Dearomatization. *J. Am. Chem. Soc.* **2022**, *144*, 5750–5755.

(21) (a) Corey, E. J.; Cheng, X.-M. *The Logic of Chemical Synthesis*; Wiley, 1989. (b) Hoffmann, R. W. *Elements of Synthesis Planning*; Springer, 2009. (c) Trost, B. M. The Atom Economy—A Search for Synthetic Efficiency. *Science* **1991**, *254*, 1471–1477. (d) Wilson, R. M.; Danishefsky, S. J. Pattern Recognition in Retrosynthetic Analysis: Snapshots in Total Synthesis. *J. Org. Chem.* **2007**, *72*, 4293–4305. (e) Young, I. S.; Baran, P. S. Protecting-Group-Free Synthesis as an Opportunity for Invention. *Nat. Chem.* **2009**, *1*, 193–205. (f) Burns, N. Z.; Baran, P. S.; Hoffmann, R. W. Redox Economy in Organic Synthesis. *Angew. Chem., Int. Ed.* **2009**, *48*, 2854–2867. (g) Gaich, T.; Baran, P. S. Aiming for the Ideal Synthesis. *J. Org. Chem.* **2010**, *75*, 4657–4673. (h) Peters, D. S.; Pitts, C. R.; McClymont, K. S.; Stratton, T. P.; Bi, C.; Baran, P. S. Ideality in Context: Motivations for Total Synthesis. *Acc. Chem. Res.* **2021**, *54*, 605–617. (i) Wender, P. A. Toward the Ideal Synthesis and Transformative Therapies: The Roles of Step Economy and Function Oriented Synthesis. *Tetrahedron* **2013**, *69*, 7529–7550. (j) Hayashi, Y. Time Economy in Total Synthesis. *J. Org. Chem.* **2021**, *86*, 1–23.

(22) (a) McDonald, F. E.; Towne, T. B. *syn*-Oxidative Polycyclization of Hydroxy Polyenes: A New Approach to Polyether Synthesis. *J. Am. Chem. Soc.* **1994**, *116*, 7921–7922. (b) Towne, T. B.; McDonald, F. E. *syn*-Oxidative Polycyclizations of Hydroxypolyenes: Highly Stereoselective and Potentially Biomimetic Syntheses of *all-trans*-Polytetrahydrofurans. *J. Am. Chem. Soc.* **1997**, *119*, 6022–6028.

(23) The desired product **13** could be obtained by means of the corresponding acid intermediate under the same reaction conditions; further mechanistic studies are underway (see [Supporting Information](#), pages S30–31).

(24) Wang, W.; Yang, X.; Dai, R.; Yan, Z.; Wei, J.; Dou, X.; Qiu, X.; Zhang, H.; Wang, C.; Liu, Y.; Song, S.; Jiao, N. Catalytic Electrophilic

Halogenation of Arenes with Electron-Withdrawing Substituents. *J. Am. Chem. Soc.* **2022**, *144*, 13415–13425.

(25) (a) Feringa, B. L.; Badorrey, R.; Peña, D.; Harutyunyan, S. R.; Minnaard, A. J. Copper-Catalyzed Asymmetric Conjugate Addition of Grignard Reagents to Cyclic Enones. *Proc. Natl. Acad. Sci. U. S. A.* **2004**, *101*, 5834–5838. (b) López, F.; Harutyunyan, S. R.; Minnaard, A. J.; Feringa, B. L. Copper-Catalyzed Enantioselective Conjugate Addition of Grignard Reagents to Acyclic Enones. *J. Am. Chem. Soc.* **2004**, *126*, 12784–12785.

(26) To our knowledge, enantioselective conjugate addition reactions involving arylmetal reagents remain challenging, and little progress has been achieved to date. For selected reviews, see: (a) Alexakis, A.; Bäckvall, J. E.; Krause, N.; Pàmies, O.; Diéguez, M. Enantioselective Copper-Catalyzed Conjugate Addition and Allylic Substitution Reactions. *Chem. Rev.* **2008**, *108*, 2796–2823. (b) Jerphagnon, T.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. Recent Advances in Enantioselective Copper-catalyzed 1,4-Addition. *Chem. Soc. Rev.* **2009**, *38*, 1039–1075. (c) Thaler, T.; Knochel, P. Copper-Catalyzed Asymmetric Michael Addition of Magnesium, Zinc, and Aluminum Organometallic Reagents: Efficient Synthesis of Chiral Molecules. *Angew. Chem., Int. Ed.* **2009**, *48*, 645–648. (d) Hayashi, M.; Matsubara, R. Recent Topics on Catalytic Asymmetric 1,4-Addition. *Tetrahedron Lett.* **2017**, *58*, 1793–1805.

(27) Zhang, S.-S.; Hu, T.-J.; Li, M.-Y.; Song, Y.-K.; Yang, X.-D.; Feng, C.-G.; Lin, G.-Q. Asymmetric Alkenylation of Enones and Imines Enabled by A Highly Efficient Aryl to Vinyl 1,4-Rhodium Migration. *Angew. Chem., Int. Ed.* **2019**, *58*, 3387–3391.

(28) Lang, Q.; Yang, H.; Gu, G.; Feng, Q.; Wen, J.; Zhang, X. Enantioselective Hydrogenation of Endocyclic Enones: the Solution to a Historical Problem. *Chin. J. Chem.* **2021**, *39*, 933–936.

Recommended by ACS

Syntheses of Bufospirostenin A and Ophiopogonol A by a Conformation-Controlled Transannular Prins Cyclization

Peicheng Yang, Jinghan Gui, *et al.*

SEPTEMBER 20, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Total Synthesis of Matrine Alkaloids

Nicholas L. Magann, Michael S. Sherburn, *et al.*

OCTOBER 19, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Total Synthesis of (+)-Mutilin: A Transannular [2+2] Cycloaddition/Fragmentation Approach

Han Chen, Tuoping Luo, *et al.*

AUGUST 22, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Enantioselective Total Synthesis of (+)-Aberrarone

Willi M. Amberg and Erick M. Carreira

AUGUST 19, 2022
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >