

Topological Persistence (in a nutshell)

X topological space

$$f : X \rightarrow \mathbb{R}$$






persistence



$\mathrm{Dg} f$

signature: *persistence diagram*


encodes the topological structure of the pair (X, f)



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family



Topological Persistence (in a nutshell)

Inside the black box:


- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family
- Finite set of intervals (barcode) encodes births/deaths of topological features




Topological Persistence (in a nutshell)

Inside the black box:

- Nested family (*filtration*) of sublevel-sets $f^{-1}((-\infty, t])$ for t ranging over \mathbb{R}
- Track the evolution of the topology (homology) throughout the family
- Finite set of intervals (barcode) encodes births/deaths of topological features





- Alternate representation as a (multi-) set of points in the plane (*diagram*).



Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$



$$x \mapsto \min_{p \in P} \|x - p\|_2$$



Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$



$$x \mapsto \min_{p \in P} \|x - p\|_2$$



Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$



$$x \mapsto \min_{p \in P} \|x - p\|_2$$



Example: Distance Function



$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$

$$x \mapsto \min_{p \in P} \|x - p\|_2$$





Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$
$$x \mapsto \min_{p \in P} \|x - p\|_2$$





Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$
$$x \mapsto \min_{p \in P} \|x - p\|_2$$





Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$
$$x \mapsto \min_{p \in P} \|x - p\|_2$$




Example: Distance Function

$$f_P : \mathbb{R}^2 \rightarrow \mathbb{R}$$
$$x \mapsto \min_{p \in P} \|x - p\|_2$$



Example: Distance Function



Topological Persistence (in a nutshell)

3 pillars:

1. Decomposition theorems (existence of barcodes / diagrams)
2. algorithm (computation of barcodes / diagrams)
3. stability theorem (use of barcodes as signatures in applications)

Mathematical viewpoint: homology + quivers

Filtration: $F_1 \subseteq F_2 \subseteq F_3 \subseteq F_4 \subseteq F_5 \cdots$


Example 1: *offsets filtration* (nested family of unions of balls, cf. previous slide)

Mathematical viewpoint: homology + quivers

Filtration: $F_1 \subseteq F_2 \subseteq F_3 \subseteq F_4 \subseteq F_5 \cdots$

Example 1: *offsets filtration* (nested family of unions of balls, cf. previous slide)

Example 2: *simplicial filtration* (nested family of simplicial complexes)




Mathematical viewpoint: homology + quivers

Filtration: $F_1 \subseteq F_2 \subseteq F_3 \subseteq F_4 \subseteq F_5 \cdots$

Example 1: *offsets filtration* (nested family of unions of balls, cf. previous slide)


Example 2: *simplicial filtration* (nested family of simplicial complexes)

Example 3: *sublevel-sets filtration* (family of sublevel sets of a function $f : X \rightarrow \mathbb{R}$)



Mathematical viewpoint: homology + quivers

Filtration: $F_1 \subseteq F_2 \subseteq F_3 \subseteq F_4 \subseteq F_5 \cdots$



(homology functor)



topological level

algebraic level

Persistence module: $H_*(F_1) \rightarrow H_*(F_2) \rightarrow H_*(F_3) \rightarrow H_*(F_4) \rightarrow H_*(F_5) \cdots$

Mathematical viewpoint: homology + quivers

Example:



(1-homology functor)



$$\mathbf{k} \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} \mathbf{k}^2 \xrightarrow{\begin{pmatrix} 0 & 1 \end{pmatrix}} \mathbf{k} \xrightarrow{\begin{pmatrix} 0 \\ 1 \end{pmatrix}} \mathbf{k}^2 \xrightarrow{\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}} \mathbf{k}^2 \dots$$

Mathematical viewpoint: homology + quivers

Theorem. Let M be a persistence module over an index set $T \subseteq \mathbb{R}$. Then, M decomposes as a direct sum of *interval modules* $\mathbf{k}_{[b,d]}$:

$$\underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t < [b,d]} \xrightarrow{0} \underbrace{\mathbf{k} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \mathbf{k}}_{[b,d]} \xrightarrow{0} \underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t > [b,d]}$$

$$M \simeq \bigoplus_{j \in J} \mathbf{k}_{[b_j, d_j]}$$



(the barcode is a complete descriptor of the algebraic structure of M)

Mathematical viewpoint: homology + quivers

Theorem. Let M be a persistence module over an index set $T \subseteq \mathbb{R}$. Then, M decomposes as a direct sum of *interval modules* $\mathbf{k}_{[b,d]}$:

$$\underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t < [b,d]} \xrightarrow{0} \underbrace{\mathbf{k} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \mathbf{k}}_{[b,d]} \xrightarrow{0} \underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t > [b,d]}$$

in the following cases:

- T is finite [Gabriel 1972] [Auslander 1974],
- M is *pointwise finite-dimensional* (every space M_t has finite dimension) [Webb 1985] [Crawley-Boevey 2012].

Moreover, when it exists, the decomposition is **unique** up to isomorphism and permutation of the terms [Azumaya 1950].

(Note: this is independent of the choice of field \mathbf{k} .)

Mathematical viewpoint: homology + quivers

Theorem. Let M be a persistence module over an index set $T \subseteq \mathbb{R}$. Then, M decomposes as a direct sum of *interval modules* $\mathbf{k}_{[b,d]}$:

$$\underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t < [b,d]} \xrightarrow{0} \underbrace{\mathbf{k} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \mathbf{k}}_{[b,d]} \xrightarrow{0} \underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t > [b,d]}$$

in the following cases:

- T is finite [Gabriel 1972] [Auslander 1974],
- M is *pointwise finite-dimensional* (every space M_t has finite dimension) [Webb 1985] [Crawley-Boevey 2012].

► **Not sufficient for our purposes:**

∃ compact sets whose offsets do not induce *pfd* modules.



Mathematical viewpoint: homology + quivers

Theorem. Let M be a persistence module over an index set $T \subseteq \mathbb{R}$. Then, M decomposes as a direct sum of *interval modules* $\mathbf{k}_{[b,d]}$:

$$\underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t < [b,d]} \xrightarrow{0} \underbrace{\mathbf{k} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \mathbf{k}}_{[b,d]} \xrightarrow{0} \underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t > [b,d]}$$

in the following cases:

- T is finite [Gabriel 1972] [Auslander 1974],
- M is *pointwise finite-dimensional* (every space M_t has finite dimension) [Webb 1985] [Crawley-Boevey 2012].

► **Not sufficient for our purposes:**

∃ compact sets whose offsets do not induce *pfd* modules.



Mathematical viewpoint: homology + quivers

Theorem. Let M be a persistence module over an index set $T \subseteq \mathbb{R}$. Then, M decomposes as a direct sum of *interval modules* $\mathbf{k}_{[b,d]}$:

$$\underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t < [b,d]} \xrightarrow{0} \underbrace{\mathbf{k} \xrightarrow{\text{id}} \dots \xrightarrow{\text{id}} \mathbf{k}}_{[b,d]} \xrightarrow{0} \underbrace{0 \xrightarrow{0} \dots \xrightarrow{0} 0}_{t > [b,d]}$$

in the following cases:


- T is finite [Gabriel 1972] [Auslander 1974],
- M is *pointwise finite-dimensional* (every space M_t has finite dimension) [Webb 1985] [Crawley-Boevey 2012].
- M is *q-tame* ($\text{rank } m_s^t < \infty$ for all $s < t \in T$).

→ barcode is well-defined, even though M may not be interval-decomposable


[Chazal, Cohen-Steiner, Glisse, Guibas, O. 2009] [Chazal, de Silva, Glisse, O. 2016]

Example: Distance Function


$(\mathbb{R} \bmod \mathbb{Z})^2$




$$(u, v) \mapsto \frac{1}{\sqrt{2}} (\cos(2\pi u), \sin(2\pi u), \cos(2\pi v), \sin(2\pi v))$$



$\subset S^3 \subset \mathbb{R}^4$




Example: Distance Function




(spiral winding around the flat torus)


$$(u, v) \mapsto \frac{1}{\sqrt{2}} (\cos(2\pi u), \sin(2\pi u), \cos(2\pi v), \sin(2\pi v))$$





$$\subset \mathbb{S}^3 \subset \mathbb{R}^4$$



Example: Distance Function



(spiral winding around the flat torus)



Stability Properties

