



A numerical solution for variable order fractional functional differential equation



Yun-Tao Jia, Min-Qiang Xu*, Ying-Zhen Lin

School of Science, Zhuhai Campus, Beijing Institute of Technology, Zhuhai, Guangdong, 519085, China

ARTICLE INFO

Article history:

Received 18 July 2016

Received in revised form 28 August 2016

Accepted 28 August 2016

Available online 6 September 2016

Keywords:

Variable fractional order

Functional differential equation

Simplified reproducing kernel method

ABSTRACT

This paper is concerned with an efficient numerical scheme for variable order fractional functional boundary value problems. The algorithm relies on the simplified reproducing kernel method (SRKM). The convergence of the method is proven, followed by estimates on approximate solution. A numerical example with exact solutions is studied to demonstrate the performance of the method. Results obtained by the method indicate the algorithm performs extremely well in terms of accuracy, efficiency and simplicity.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

A great quantity of nature phenomena can be modeled by variable fractional differential equations. The study of such problems has attracted much attention recently. Results about existence and uniqueness results for boundary value problems of fractional differential equations have been obtained in [1,2]. On the other side, several numerical methods, including finite difference method, matrix approach and reproducing kernel method, for solving variable-order fractional differential equations were proposed in [3–8]. At the mean time, Sun and Chen described some valuable applications of variable fractional derivative in [9–11]. Functional differential equations arose in a variety of applications, such as electrodynamics, astrophysics, nonlinear dynamical systems, and probability theory on algebraic structure. Consequently, several research papers were issued to investigate the theory and solutions of some functional differential equations (see [12–14] and references therein).

However, limited work has been done in the study on variable order fractional functional equation. In this letter, based on previous work [15–20], we focus on providing a numerical scheme to solve the following

* Corresponding author.

E-mail address: mqxu.math@126.com (M. Xu).

variable fractional functional boundary value problems in reproducing kernel space.

$$\begin{cases} D^{\alpha(x)}u(x) + a(x)u'(x) + b(x)u(x) + c(x)u(\tau(x)) = f(x), & x \in [0, 1], \\ u(0) = \lambda_0, & u(1) = \lambda_1 \end{cases} \quad (1)$$

where $a(x), b(x), c(x) \in C^2[0, 1]$, $\alpha(x), \tau(x), f(x) \in C[0, 1]$, $1 \leq \alpha(x) < 2$, $0 \leq \tau(x) \leq 1$ and $D^{\alpha(x)}$ denotes the variable order Caputo derivative.

We organize the rest of paper as follows. In Section 2, we present some reproducing kernel space. In Section 3, the SRKM is applied to solve Eq. (1) and the uniform convergence analysis is carried out as well. A numerical experiment is studied to demonstrate the efficiency, reliability and simplicity of the algorithm in Section 4. Finally, we summarize this paper in Section 5.

2. Several reproducing kernel spaces

Definition 2.1. The variable fractional order $\alpha(x)$ of function $u(x)$ in Caputo's sense, is defined as

$$D^{\alpha(x)}u(x) = \frac{1}{\Gamma(2 - \alpha(x))} \int_0^x (x - t)^{1 - \alpha(x)} u''(t) dt,$$

where $\Gamma(\cdot)$ is Gamma function.

Definition 2.2. $W_2^m[0, 1] = \{u(x) | u^{m-1}(x) \text{ is an absolutely continuous real value function in } [0, 1], u^{(m)}(x) \in L^2[0, 1]\}$ with the inner products and norms as follows:

$$\begin{aligned} \langle u(x), v(x) \rangle_m &= \sum_{k=0}^{m-1} u^{(k)}(0)v^{(k)}(0) + \int_0^1 u^{(m)}(x)v^{(m)}(x)dx, \quad \forall u, v \in W_2^m[0, 1], \\ \|u\|_m^2 &= \sqrt{\langle u, u \rangle_m}, \quad \forall u \in W_2^m[0, 1]. \end{aligned} \quad (2)$$

Clearly, the inner spaces $W_2^3[0, 1]$ and $W_2^1[0, 1]$ are reproducing kernel spaces with kernel functions $R_x(y)$ and $r_x(y)$, respectively, (see [21]), and the expression of $R_x(y)$ is

$$R_x(y) = \begin{cases} \frac{1}{120}(120 - 5xy^4 + y^5 + 10x^2y^2(3 + y) + 120xy), & y > x, \\ \frac{1}{120}(120 + x^5 - 5x^4y + 10x^2y^2(3 + x) + 120xy), & y \leq x. \end{cases}$$

By Eq. (1), we define a linear operator $\mathcal{L} : W_2^3 \rightarrow W_2^1$, for all $u \in W_2^3$ as follows

$$\mathcal{L}u(x) = D^{\alpha(x)}u(x) + a(x)u'(x) + b(x)u(x) + c(x)u(\tau(x)), \quad (3)$$

then Eq. (1) can be converted into the following equivalent operator equation

$$\begin{cases} \mathcal{L}u = f(x), \\ u(0) = \lambda_0, & u(1) = \lambda_1. \end{cases} \quad (4)$$

Lemma 2.3. \mathcal{L} is a bounded linear operator.

Proof. By the reproducing property of $R_x(y)$, we have,

$$u^{(i)}(x) = \langle u, \partial_x^i R_x \rangle_3, \quad i = 0, 1, 2, 3, \quad u^{(j)}(\tau(x)) = \langle u, \partial_x^j R_{\tau(x)} \rangle_3, \quad j = 0, 1. \quad (5)$$

A direct application of the *Cauchy–Schwarz* inequality to Eq. (5) yields that

$$|u^{(i)}(x)| \leq \|u\|_3 \cdot \|\partial_x^i R_x\|_3, \quad |u^{(j)}(\tau(x))| \leq \|u\|_3 \cdot \|\partial_x^j R_{\tau(x)}\|_3.$$

By the fact that $\|\partial_x^i R_x\|_3$ and $\|\partial_x^j R_{\tau(x)}\|_3$ are boundedness on $[0, 1]$, there exist positive constants M_i, N_j such that

$$|u^{(i)}(x)| \leq M_i \|u\|_3, \quad |u^{(j)}(\tau(x))| \leq N_j \|u\|_3. \quad (6)$$

A simple calculation together with Eq. (6) yields that

$$\begin{aligned} \int_0^1 (b(x)u(x))'^2 dx &= \int_0^1 [b^2(x)u^2(x) + b^2(x)u'^2(x) + 2b(x)b'(x)u(x)u'(x)] dx \\ &\leq \widetilde{M} \left[\int_0^1 u^2(x) dx + \int_0^1 u'^2(x) dx + \int_0^1 u(x)u'(x) dx \right] \\ &\leq \widetilde{A}_0 \|u\|_3^2. \end{aligned}$$

Therefore, there holds

$$\|b(x)u(x)\|_1^2 = b^2(0)u^2(0) + \int_0^1 (b(x)u(x))'^2 dx \leq A_0 \|u\|_3^2, \quad (7)$$

and by the same argument,

$$\|a(x)u'(x)\|_1^2 \leq A_1 \|u\|_3^2, \quad \|c(x)u(\tau(x))\|_1^2 \leq A_2 \|u\|_3^2. \quad (8)$$

Next, the use of integration by parts gives

$$D^{\alpha(x)}u(x) = \frac{x^{2-\alpha(x)}}{\Gamma(3-\alpha(x))} u''(0) + \frac{1}{\Gamma(3-\alpha(x))} \int_0^x (x-t)^{2-\alpha(x)} u'''(t) dt.$$

Since

$$\begin{aligned} \left\| \int_0^x (x-t)^{2-\alpha(x)} u'''(t) dt \right\|_1^2 &= \int_0^1 \left[\int_0^x (x-t)^{2-\alpha(x)} u'''(t) dt \right]'^2 dx \\ &= \int_0^1 \left[\int_0^x (x-t)^{2-\alpha(x)} \left(-\alpha'(x) \ln(x-t) + \frac{2-\alpha(x)}{x-t} \right) u'''(t) dt \right]^2 dx \\ &\leq \|u\|_3^2 \int_0^1 \left[\frac{\alpha'(x)}{(3-\alpha(x))^2} x^{3-\alpha(x)} - \frac{\alpha'(x)}{3-\alpha(x)} x^{3-\alpha(x)} \ln(x) + x^{2-\alpha(x)} \right]^2 dx \\ &\leq \widetilde{A}_3 \|u\|_3^2, \end{aligned}$$

we immediately have

$$\|D^{\alpha(x)}u(x)\|_1^2 \leq A_3 \|u\|_3^2. \quad (9)$$

Summarizing the estimate Eqs. (7)–(9), there exists a positive constant A such that

$$\|\mathcal{L}u(x)\|_1 \leq A \|u\|_3.$$

Namely, \mathcal{L} is a bounded linear operator. \square

3. SRKM and convergence analysis

3.1. The approximate solution of Eq. (4)

In view of Lemma 2.3, it is clear that $\mathcal{L} : W_2^3[0, 1] \rightarrow W_2^1[0, 1]$ is a bounded linear operator, put $\psi_i(x) = \mathcal{L}^* r_x(x_i)$, where \mathcal{L}^* is the conjugate operator of \mathcal{L} and $\{x_i\}_{i=1}^\infty$ is a dense subset of $[0, 1]$. Considering the boundary conditions, we skillfully construct $\varphi_1(x)$ and $\varphi_2(x)$ like collocation method in [22,23] as follows:

$$\varphi_1(x) = R_x(0), \quad \varphi_2(x) = R_x(1).$$

Theorem 3.1. For each fixed n , $\{\varphi_1, \varphi_2\} \cup \{\psi_j\}_1^n$ are linearly independent in $W_2^3[0, 1]$.

The proof can be found in [19].

Let

$$S = \overline{\text{span}\{\{\varphi_1, \varphi_2\} \cup \{\psi_j\}_1^\infty\}}, \quad S_n = \overline{\text{span}\{\{\varphi_1, \varphi_2\} \cup \{\psi_j\}_1^n\}}$$

and we denote \mathcal{P} and \mathcal{P}_n as the projection operators from W_2^3 to S and S_n , respectively.

Theorem 3.2. If $u(x)$ is the solution of Eq. (4), then $y = \mathcal{P}u$ is also the solution of Eq. (4) in S .

Proof.

$$\begin{aligned} \mathcal{L}(\mathcal{P}u)(x_i) &= \langle \mathcal{P}u, \mathcal{L}^* r_{x_i} \rangle_3 = \langle \mathcal{P}u, \psi_i \rangle_3 = \langle u, \mathcal{P}\psi_i \rangle_3 = \langle u, \psi_i \rangle_3 \\ &= \langle u, \mathcal{L}^* r_{x_i} \rangle_3 = \langle \mathcal{L}u, r_{x_i} \rangle_1 = \mathcal{L}u(x_i) = f(x_i). \end{aligned}$$

Since $\{x_i\}_{i=1}^\infty$ is a dense subset of $[0, 1]$, we have $\mathcal{L}(\mathcal{P}u) = f$. On the other hand,

$$\begin{aligned} \mathcal{P}u(0) &= \langle \mathcal{P}u, \varphi_1(x) \rangle = \langle u, \mathcal{P}R_x(0) \rangle = \langle u, R_x(0) \rangle = u(0) = \lambda_0, \\ \mathcal{P}u(1) &= \langle \mathcal{P}u, \varphi_2(x) \rangle = \langle u, \mathcal{P}R_x(1) \rangle = \langle u, R_x(1) \rangle = u(1) = \lambda_1. \end{aligned}$$

Namely, $\mathcal{P}u$ is the solution of Eq. (4). \square

Considering numerical computation, we define the n -term approximation to $y(x)$ by

$$y_n(x) = (\mathcal{P}_n y)(x).$$

Therefore, the simplified representation of the reproducing kernel method reads as: find $y_n \in S_n$ such that

$$\begin{cases} \langle y_n, \psi_i \rangle = f(x_i), & i = 1, 2, \dots, n, \\ \langle y_n, \varphi_1 \rangle = \lambda_0, & \langle y_n, \varphi_2 \rangle = \lambda_1. \end{cases} \quad (10)$$

As $y_n \in S_n$, y_n has the representation

$$y_n(x) = \sum_{i=1}^2 \beta_i \varphi_i(x) + \sum_{j=1}^n \alpha_j \psi_j(x), \quad (11)$$

where $\beta_1, \beta_2, \alpha_j, j = 1, 2, \dots, n$ are constants to be determined.

Substituting Eq. (11) into Eq. (10), we derive a linear system. Thereupon, the approximate solution $y_n(x)$ given in Eq. (11) can be obtained by solving the linear system.

Table 1Comparison of absolute errors for [Example 4.1](#).

x	$E_{10}(x)$ in [8]	Our method	$E_{20}(x)$ in [8]	Our method
0.1	1.43×10^{-6}	1.05×10^{-7}	1.92×10^{-7}	1.17×10^{-8}
0.2	2.28×10^{-6}	1.80×10^{-7}	3.19×10^{-7}	1.77×10^{-8}
0.3	2.84×10^{-6}	2.32×10^{-7}	4.01×10^{-7}	2.17×10^{-8}
0.4	3.11×10^{-6}	2.60×10^{-7}	4.44×10^{-7}	2.39×10^{-8}
0.5	3.13×10^{-6}	2.66×10^{-7}	4.53×10^{-7}	2.45×10^{-8}
0.6	2.94×10^{-6}	2.51×10^{-7}	4.30×10^{-7}	2.34×10^{-8}
0.7	2.53×10^{-6}	2.24×10^{-7}	3.80×10^{-7}	2.07×10^{-8}
0.8	1.88×10^{-6}	1.60×10^{-7}	3.11×10^{-7}	1.59×10^{-8}
0.9	9.97×10^{-7}	6.54×10^{-8}	2.08×10^{-7}	1.11×10^{-8}

Table 2Maximum absolute errors for [Example 4.1](#).

E'_{10}	$D^{3/2}E_{10}$	E'_{20}	$D^{3/2}E_{20}$
1.15×10^{-6}	3.50×10^{-6}	1.01×10^{-8}	2.83×10^{-8}

3.2. Convergence analysis

Theorem 3.3. If $y_n(x) \rightarrow y(x)$ in S , then $y_n \rightarrow y$ uniformly on $[0, 1]$. Furthermore, $D^{\alpha(x)}y_n \rightarrow D^{\alpha(x)}y$ uniformly on $[0, 1]$.

Proof. Note that

$$\begin{aligned} y_n(x) &= \langle y_n, R_x \rangle_3, & D^{\alpha(x)}y_n(x) &= \langle y_n, D^{\alpha(x)}R_x \rangle_3, \\ y(x) &= \langle y, R_x \rangle_3, & D^{\alpha(x)}y(x) &= \langle y, D^{\alpha(x)}R_x \rangle_3. \end{aligned}$$

By applying *Cauchy-Schwarz* inequality and the continuity of $\|D^{\alpha(x)}R_x\|_3$, there holds

$$\begin{aligned} |y_n(x) - y(x)| &= |\langle y_n - y, R_x \rangle| \leq \|y_n - y\|_3 \cdot \|R_x\|_3 \leq M\|y_n - y\|_3 \rightarrow 0. \\ |D^{\alpha(x)}(y_n(x) - y(x))| &= |\langle y_n - y, D^{\alpha(x)}R_x \rangle| \leq \|y_n - y\|_3 \cdot \|D^{\alpha(x)}R_x\|_3 \leq N\|y_n - y\|_3 \rightarrow 0. \end{aligned}$$

Namely, y_n and $D^{\alpha(x)}y_n$ uniformly convergent to y and $D^{\alpha(x)}y$, respectively. \square

The following corollary is a direct consequence of [Theorems 3.2](#) and [3.3](#).

Corollary 3.4. Let $u(x)$ be the solution of Eq. (4), then $y_n = \mathcal{P}_n y \rightarrow y = \mathcal{P}u$, moreover, $y_n \rightarrow y$ uniformly.

Theorem 3.5. The approximate solution $y_n(x)$ for Eq. (4) has second order convergence, that is, there exists a positive constant C such that

$$|y_n(x) - y(x)| \leq \frac{C}{n^2} \|f\|_1.$$

For the proof, one may refer to [\[20\]](#).

4. Numerical examples

Example 4.1. Consider the following variable fractional BVPs [\[8\]](#) of the form

$$\begin{cases} D^{\alpha(x)}u(x) + \cos x u'(x) + 4u(x) + 5u(x^2) = f(x), & x \in [0, 1], \\ u(0) = 0, & u(1) = 1, \end{cases}$$

where $\alpha(x) = \frac{5+\sin(x)}{4}$, $f(x) = \frac{2x^{2-\alpha(x)}}{\Gamma(3-\alpha(x))} + 5x^4 + 4x^2 + 2x \cos(x)$. The exact solution is $u(x) = x^2$. Applying SRKM, we take $x_i = \frac{i}{n}$, $i = 1, 2, \dots, n$, $n = 10, 20$. The absolute error $E_n(x) = |y_n(x) - u(x)|$ obtained is compared with [8] in Table 1. Moreover, we report maximum errors $E'_n = |y'_n(x) - u'(x)|_{max}$ and $D^{3/2}E_n = |D^{3/2}y_n(x) - D^{3/2}u(x)|_{max}$ in Table 2.

5. Conclusion

In this paper, an effective numerical algorithm based on SRKM is proposed for variable order fractional differential equations with boundary conditions. Numerical example shows that our algorithm is simple, accurate and effective.

References

- [1] R.P. Agarwal, B. Ahmad, Existence theory for anti-periodic boundary value problems of fractional equations and inclusion, *Comput. Math. Appl.* 62 (2011) 1200–1214.
- [2] R.P. Agarwal, M. Benchohra, H. Hamani, A survey on existence results for boundary value problems of nonlinear fractional equations and inclusion, *Acta Appl. Math.* 109 (2010) 973–1033.
- [3] Q.M. Al-Mdallal, M.I. Syam, M.N. Anwar, A collocation-shooting method for solving fractional boundary value problems, *Commun. Nonlinear Sci. Numer. Simul.* 15 (2010) 3814–3822.
- [4] M. Rehman, R.A. Khan, The Legendre wavelet method for solving fractional differential equations, *Commun. Nonlinear Sci. Numer. Simul.* 16 (2011) 4163–4173.
- [5] M. Refai, M.A. Haiji, Monotone iterative sequences for wavelet method for nonlinear boundary value problems of fractional order, *Nonlinear Anal.* 74 (2011) 3531–3539.
- [6] A. Pedas, E. Tamme, Piecewise polynomial collocation for linear boundary value problems of fractional differential equations, *J. Comput. Appl. Math.* 236 (2012) 3349–3359.
- [7] Y. Jia, Y. Lin, A new algorithm for nonlinear fractional BVPs, *Appl. Math. Lett.* 57 (2016) 121–125.
- [8] X. Li, B. Wu, A numerical technique for variable fractional functional boundary value problems, *Appl. Math. Lett.* 43 (2015) 108–113.
- [9] H.G. Sun, W. Chen, S. Hu, Y.Q. Chen, On mean square displacement behaviors of anomalous diffusions with variable and random orders, *Phys. Lett. A* 374 (2010) 906–910.
- [10] W. Chen, J.J. Zhang, J.Y. Zhang, A variable-order time-fractional derivative model for chloride ion sub-diffusion in concrete structures, *Fract. Calc. Appl. Anal.* 16 (1) (2013) 76–92.
- [11] H.G. Sun, Y. Zhang, W. Chen, D.M. Reeves, Use of a variable-index fractional-derivative model to capture transient dispersion in heterogeneous media, *J. Contam. Hydrol.* 157 (2014) 47–58.
- [12] A. Saadatmandi, M. Dehghan, Variational iteration method for solving a generalized pantograph equation, *Comput. Math. Appl.* 58 (2009) 2190–2196.
- [13] W.S. Wang, T. Qin, Stability of one-leg θ -methods for nonlinear neutral differential equations with proportional delay, *Appl. Math. Comput.* 213 (2009) 177–183.
- [14] H. Kocak, A. Yildirim, Series solution for a delay differential equation arising in electrodynamics, *Commun. Numer. Methods Eng.* 25 (2009) 1084–1096.
- [15] M.Q. Xu, Y.Z. Lin, Simplified reproducing kernel method for fractional differential equations with delay, *Appl. Math. Lett.* 52 (2016) 156–161.
- [16] Y.T. Jia, Y.Z. Lin, A numerical algorithm for singular multi-point BVPs using the reproducing kernel method, *Pure Appl. Math.* 21 (2014) 51–60.
- [17] M.Q. Xu, Y.Z. Lin, A new algorithm for nonlinear fourth order multi-point boundary value problems, *Appl. Math. Comput.* 1 (2016) 163–168.
- [18] Y.Z. Lin, M.G. Cui, A new method to solving the damped nonlinear Klein-Gordon equation, *Sci. China A* 51 (2) (2008) 561–572.
- [19] R.M. Zhang, Y.Z. Lin, A novel method for nonlinear boundary value problems, *J. Comput. Appl. Math.* 282 (2015) 77–82.
- [20] Z.H. Zhao, Y.Z. Lin, Convergence order of the reproducing kernel method for solving boundary value problems, *Math. Model. Anal.* 21 (2016) 466–477.
- [21] B. Wu, Y. Lin, *Application of the Reproducing Kernel Space*, Science Press, 2012.
- [22] R. Salehi, M. Dehghan, A generalized moving least square reproducing kernel method, *J. Comput. Appl. Math.* 249 (2013) 120–132.
- [23] M. Dehghan, R. Salehi, A meshless based numerical technique for traveling solitary wave solution of Boussinesq equation, *Appl. Math. Model.* 36 (5) (2012) 1939–1956.