



Short communication

The reciprocal of the geometric mean of many positive numbers is a Stieltjes transform

Feng Qi^{a,b,c,*}, Bai-Ni Guo^d^a Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China^b College of Mathematics, Inner Mongolia University for Nationalities, Tongliao City, Inner Mongolia Autonomous Region, 028043, China^c Department of Mathematics, College of Science, Tianjin Polytechnic University, Tianjin City, 300387, China^d School of Mathematics and Informatics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China

ARTICLE INFO

Article history:

Received 7 July 2016

MSC:

primary 30E20

secondary 26E60

44A15

44A20

Keywords:

Integral representation

Reciprocal of the geometric mean

Stieltjes transform

Completely monotonic function

Logarithmically completely monotonic function

Cauchy integral formula

ABSTRACT

In the paper, by the Cauchy integral formula in the theory of complex functions, an integral representation for the reciprocal of the geometric mean of many positive numbers is established. As a result, the reciprocal of the geometric mean of many positive numbers is verified to be a Stieltjes transform and, consequently, a (logarithmically) completely monotonic function.

© 2016 Elsevier B.V. All rights reserved.

1. Preliminaries

We recall some definitions and notion.

Recall from [1, Chapter IV] that an infinitely differentiable function f on an interval I is said to be completely monotonic on I if it satisfies $(-1)^{n-1}f^{(n-1)}(x) \geq 0$ for $x \in I$ and $n \in \mathbb{N}$, where \mathbb{N} stands for the set of all positive integers. Theorem 12b in [1] reads that a necessary and sufficient condition that $f(x)$ should be completely monotonic for $0 < x < \infty$ is that

$$f(x) = \int_0^\infty e^{-xt} d\alpha(t),$$

where $\alpha(t)$ is non-decreasing and the integral converges for $0 < x < \infty$.

In [2–4], it was defined implicitly and explicitly that an infinitely differentiable and positive function f is said to be logarithmically completely monotonic on an interval I if $(-1)^k[\ln f(x)]^{(k)} \geq 0$ on I for all $k \in \mathbb{N}$. In [5, Theorem 1.1],

* Corresponding author at: Institute of Mathematics, Henan Polytechnic University, Jiaozuo City, Henan Province, 454010, China.

E-mail addresses: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com (F. Qi), bai.ni.guo@gmail.com, bai.ni.guo@hotmail.com (B.-N. Guo).

URLs: <https://qifeng618.wordpress.com> (F. Qi), http://www.researchgate.net/profile/Bai-Ni_Guo/ (B.-N. Guo).

[6, Theorem 4], [3, Theorem 1], and [4, Theorem 4], it was found and verified once again that a logarithmically completely monotonic function must be completely monotonic, but not conversely.

In [7, Definition 2.1], it was defined that a Stieltjes transform is a function $f : (0, \infty) \rightarrow [0, \infty)$ which can be written in the form

$$f(x) = \frac{a}{x} + b + \int_0^\infty \frac{1}{u+x} d\mu(u), \quad (1.1)$$

where a, b are nonnegative constants and μ is a nonnegative measure on $(0, \infty)$ such that the integral $\int_0^\infty \frac{1}{1+s} d\mu(s) < \infty$. In [5, Theorem 1.2], it was proved that a positive Stieltjes transform must be a logarithmically completely monotonic function on $(0, \infty)$, but not conversely.

An infinitely differentiable function $f : I \rightarrow [0, \infty)$ is called a Bernstein function on an interval I if f' is completely monotonic on I . The Bernstein functions on $(0, \infty)$ can be characterized by [7, Theorem 3.2] which states that a function $f : (0, \infty) \rightarrow [0, \infty)$ is a Bernstein function if and only if it admits the representation

$$f(x) = a + bx + \int_0^\infty (1 - e^{-xt}) d\mu(t), \quad (1.2)$$

where $a, b \geq 0$ and μ is a measure on $(0, \infty)$ satisfying $\int_0^\infty \min\{1, t\} d\mu(t) < \infty$. The formula (1.2) is called the Lévy–Khintchine representation of f . In [8, pp. 161–162, Theorem 3] and [7, Proposition 5.25], it was proved that the reciprocal of a Bernstein function is logarithmically completely monotonic.

If the Lévy measure μ in (1.2) has a completely monotonic density $m(t)$ with respect to the Lebesgue measure, that is,

$$f(x) = a + bx + \int_0^\infty (1 - e^{-xt}) m(t) dt, \quad a, b \geq 0,$$

where $m(t)$ is a completely monotonic function on $(0, \infty)$ and satisfies $\int_0^\infty \min\{1, t\} m(t) dt < \infty$, then f is said to be a complete Bernstein function on $(0, \infty)$. Theorem 7.3 in [7] reads that a (non-trivial) function f is a complete Bernstein function if and only if $\frac{1}{f}$ is a (non-trivial) Stieltjes function.

2. Motivations and main results

Theorem 4.2 in [9] states that the principal branch of the geometric mean

$$G_{a,b}(z) = \sqrt{(a+z)(b+z)}$$

for $b > a > 0$ and $z \in \mathbb{C} \setminus [-b, -a]$ has the Lévy–Khintchine representation

$$G_{a,b}(z) = G_{a,b}(0) + z + \frac{b-a}{2\pi} \int_0^\infty \frac{\rho((b-a)s)}{s} e^{-as} (1 - e^{-zs}) ds,$$

where $\rho(s)$ is defined by

$$\rho(s) = \int_0^{1/2} \left(\sqrt{\frac{1}{u}} - 1 - \frac{1}{\sqrt{1/u - 1}} \right) [1 - e^{-(1-2u)s}] e^{-su} du > 0.$$

Consequently, the geometric mean $G_{a,b}(t-a)$ is a complete Bernstein function on $(0, \infty)$ and $\frac{1}{G_{a,b}(t-a)}$ is a Stieltjes transform.

For $b > a$ and $z \in \mathbb{C} \setminus [-b, -a]$, Lemma 2.4 in [10] reads that the reciprocal of $G_{a,b}(z)$ can be represented as

$$h_{a,b}(z) = \frac{1}{G_{a,b}(z)} = \frac{1}{\sqrt{(z+a)(z+b)}} = \frac{1}{\pi} \int_a^b \frac{1}{\sqrt{(t-a)(b-t)}} \frac{1}{t+z} dt.$$

Consequently, the function $h_{a,b}(x-a)$ is a Stieltjes transform and, consequently, a (logarithmically) completely monotonic function on $(0, \infty)$.

Let $n \in \mathbb{N}$ and $\mathbf{a} = (a_1, a_2, \dots, a_n)$ be a positive sequence, that is, $a_k > 0$ for $1 \leq k \leq n$, and no any two are same. For $z \in \mathbb{C} \setminus [-\max\{a_k, 1 \leq k \leq n\}, -\min\{a_k, 1 \leq k \leq n\}]$ and $n \geq 2$, let $\mathbf{1} = \overbrace{(1, 1, \dots, 1)}^n$ and

$$G_n(z\mathbf{1} + \mathbf{a}) = \left[\prod_{k=1}^n (a_k + z) \right]^{1/n}.$$

Let σ be a permutation of the set $\{1, 2, \dots, n\}$ such that $\sigma(\mathbf{a}) = (a_{\sigma(1)}, a_{\sigma(2)}, \dots, a_{\sigma(n)})$ is a rearrangement of \mathbf{a} in an ascending order $a_{\sigma(1)} < a_{\sigma(2)} < \dots < a_{\sigma(n)}$. Then it was obtained in [11, Theorem 1.1] that the principal branch of the geometric mean $G_n(z\mathbf{1} + \mathbf{a})$ has the Lévy–Khintchine representation

$$G_n(z\mathbf{1} + \mathbf{a}) = G_n(\mathbf{a}) + z + \int_0^\infty Q_{n,\mathbf{a}}(u) (1 - e^{-zu}) du,$$




Fig. 1. The arguments of $(x + ir) - (-a_k)$ and $(x + ir) - (-a_{k+1})$.

where

$$Q_{n,\mathbf{a}}(u) = \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin \frac{\ell\pi}{n} \int_{a_{\sigma(\ell)}}^{a_{\sigma(\ell+1)}} \left| \prod_{k=1}^n (a_k - t) \right|^{1/n} e^{-ut} dt.$$

For more information on this topic, please refer to the papers [12–18] and closely-related references therein.

The main aim of this paper is to establish by the Cauchy integral formula in the theory of complex functions an integral representation for the reciprocal $\frac{1}{G_n(z\mathbf{1}+\mathbf{a})}$ of the geometric mean $G_n(z\mathbf{1}+\mathbf{a})$ of $n > 2$ positive numbers a_k for $1 \leq k \leq n$. As a result, the reciprocal $\frac{1}{G_n(z\mathbf{1}+\mathbf{a})}$ of the geometric mean $G_n(z\mathbf{1}+\mathbf{a})$ is derived to be a Stieltjes transform and, consequently, a (logarithmically) completely monotonic function.

Our main results can be stated as the following theorem.

Theorem 2.1. For $a_k < a_{k+1}$, the principal branch of the reciprocal $\frac{1}{G_n(z\mathbf{1}+\mathbf{a})}$ of the geometric mean $G_n(z\mathbf{1}+\mathbf{a})$ can be represented as

$$\frac{1}{G_n(z\mathbf{1}+\mathbf{a})} = \frac{1}{\sqrt[n]{\prod_{k=1}^n (z + a_k)}} = \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin\left(\frac{\ell\pi}{n}\right) \int_{a_\ell}^{a_{\ell+1}} \frac{1}{\sqrt[n]{\prod_{k=1}^n |t - a_k|}} \frac{1}{t + z} dt, \quad (2.1)$$

where $z \in \mathbb{C} \setminus [-a_n, -a_1]$. Consequently, the function $\frac{1}{G_n((x-a_1)\mathbf{1}+\mathbf{a})}$ is a Stieltjes transform and a (logarithmically) completely monotonic function on $(0, \infty)$.

3. Proof of Theorem 2.1

For $r > 0$ and by virtue of Fig. 1, we see that

(1) when $x \leq -a_k$,

$$\frac{\pi}{2} \leq \arg(x + ir + a_k) < \pi;$$

(2) when $-a_{k+1} \leq x \leq -a_k$,

$$0 < \arg(x + ir + a_{k+1}) \leq \frac{\pi}{2}.$$

A straightforward computation gives

$$\begin{aligned} \frac{1}{G_n((x+ir)\mathbf{1}+\mathbf{a})} &= \frac{1}{\sqrt[n]{\prod_{k=1}^n (x + a_k + ir)}} \\ &= \frac{1}{\exp\left\{\frac{1}{n} \sum_{k=1}^n [\ln |x + a_k + ir| + i \arg(x + a_k + ir)]\right\}} \\ &\rightarrow \frac{1}{\exp\left[\frac{1}{n} \sum_{k=1}^n \ln |x + a_k|\right] \exp\left[\frac{i}{n} \sum_{k=1}^n \lim_{r \rightarrow 0^+} \arg(x + a_k + ir)\right]} \end{aligned}$$

$$\begin{aligned}
&= \begin{cases} \frac{1}{\sqrt[n]{\prod_{k=1}^n |x + a_k|}} \exp(i\pi), & x < -a_n \\ \frac{1}{\sqrt[n]{\prod_{k=1}^n |x + a_k|}} \exp\left(\frac{i}{n} \ell \pi\right), & -a_{\ell+1} < x < -a_\ell, 1 \leq \ell \leq n-1 \end{cases} \\
&= \begin{cases} \frac{1}{\sqrt[n]{\prod_{k=1}^n |x + a_k|}}, & x < -a_n \\ \frac{\cos\left(\frac{\ell}{n} \pi\right) - i \sin\left(\frac{\ell}{n} \pi\right)}{\sqrt[n]{\prod_{k=1}^n |x + a_k|}}, & -a_{\ell+1} < x < -a_\ell, 1 \leq \ell \leq n-1 \end{cases}
\end{aligned}$$

as $r \rightarrow 0^+$. Consequently, it follows that

$$\lim_{r \rightarrow 0^+} \Im \left[\frac{1}{G_n((x + ir)\mathbf{1} + \mathbf{a})} \right] = \begin{cases} 0, & x < -a_n; \\ -\frac{\sin\left(\frac{\ell}{n} \pi\right)}{\sqrt[n]{\prod_{k=1}^n |x + a_k|}}, & -a_{\ell+1} < x < -a_\ell, 1 \leq \ell \leq n-1. \end{cases} \quad (3.1)$$

For any fixed point $z_0 = x_0 + iy_0 \in \mathbb{C} \setminus (-\infty, -a_1]$, choose $r > 0$ and $R > a_n > 0$ such that

$$\begin{cases} 0 < r < |y_0| \leq |z_0| < R, & y_0 \neq 0; \\ 0 < r < x_0 + a_1 < R, & y_0 = 0; \end{cases}$$

and consider the positively oriented contour $C(r, R)$ in $\mathbb{C} \setminus (-\infty, -a_1]$, which consists of the half circle $C(a_1, r) : z + a_1 = re^{i\theta}$ for $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and two half lines $\ell_\pm : z = x \pm ir$ for $x \leq -a_1$ until they cut the circle $C(R) : z = Re^{i\theta}$ for

$$\theta \in \left(-\pi + \arctan \frac{r}{\sqrt{R^2 - r^2}}, \pi - \arctan \frac{r}{\sqrt{R^2 - r^2}} \right)$$

at the points $-\sqrt{R^2 - r^2} \pm ir$.

Along the positively oriented contour $C(r, R)$ in $\mathbb{C} \setminus (-\infty, -a_1]$, applying the Cauchy integral formula yields

$$\begin{aligned}
h_{n,\mathbf{a}}(z_0) &\triangleq \frac{1}{G_n(z_0\mathbf{1} + \mathbf{a})} = \frac{1}{2\pi i} \oint_{C(r,R)} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi \\
&= \frac{1}{2\pi i} \left[\int_{C(R)} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi + \int_{C(a,r)} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi + \int_{\ell_+} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi + \int_{\ell_-} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi \right].
\end{aligned}$$

By a straightforward computation, we have

$$\begin{aligned}
\int_{C(R)} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi &= \int_{-\pi + \arctan \frac{r}{\sqrt{R^2 - r^2}}}^{\pi - \arctan \frac{r}{\sqrt{R^2 - r^2}}} \frac{h_{n,\mathbf{a}}(Re^{i\theta})}{Re^{i\theta} - z_0} iRe^{i\theta} d\theta \\
&\rightarrow i \int_{-\pi}^{\pi} \lim_{R \rightarrow \infty} \frac{1}{1 - z_0/Re^{i\theta}} \lim_{R \rightarrow \infty} h_{n,\mathbf{a}}(Re^{i\theta}) d\theta, \quad R \rightarrow \infty \\
&= i \int_{-\pi}^{\pi} \lim_{R \rightarrow \infty} \frac{1}{G_n(Re^{i\theta}\mathbf{1} + \mathbf{a})} d\theta \\
&= 0, \\
\int_{C(a,r)} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi &= \int_{\pi/2}^{-\pi/2} \frac{ire^{i\theta}}{re^{i\theta} - a_1 - z_0} \frac{1}{G_n((re^{i\theta} - a_1)\mathbf{1} + \mathbf{a})} d\theta \\
&\rightarrow 0, \quad r \rightarrow 0^+,
\end{aligned}$$

and

$$\begin{aligned}
 & \int_{\ell_+} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi + \int_{\ell_-} \frac{h_{n,\mathbf{a}}(\xi)}{\xi - z_0} d\xi = \int_{-R(r)}^{-a_1} \frac{h_{n,\mathbf{a}}(x+ir)}{x+ir-z_0} dx + \int_{-a_1}^{-R(r)} \frac{h_{n,\mathbf{a}}(x-ir)}{x-ir-z_0} dx \\
 &= \int_{-R(r)}^{-a_1} \left[\frac{h_{n,\mathbf{a}}(x+ir)}{x+ir-z_0} - \frac{h_{n,\mathbf{a}}(x-ir)}{x-ir-z_0} \right] dx \\
 &= \int_{-R(r)}^{-a_1} \frac{(x-z_0)[h_{n,\mathbf{a}}(x+ir) - h_{n,\mathbf{a}}(x-ir)] - ir[h_{n,\mathbf{a}}(x+ir) + h_{n,\mathbf{a}}(x-ir)]}{(x-z_0)^2 + r^2} dx \\
 &= \int_{-R(r)}^{-a_1} \frac{(x-z_0)[h_{n,\mathbf{a}}(x+ir) - h_{n,\mathbf{a}}(\overline{x+ir})] - ir[h_{n,\mathbf{a}}(x+ir) + h_{n,\mathbf{a}}(\overline{x+ir})]}{(x-z_0)^2 + r^2} dx \\
 &= \int_{-R(r)}^{-a_1} \frac{(x-z_0)[h_{n,\mathbf{a}}(x+ir) - \overline{h_{n,\mathbf{a}}(x+ir)}] - ir[h_{n,\mathbf{a}}(x+ir) + \overline{h_{n,\mathbf{a}}(x+ir)}]}{(x-z_0)^2 + r^2} dx \\
 &= \int_{-R(r)}^{-a_1} \frac{2i(x-z_0)\Im[h_{n,\mathbf{a}}(x+ir)] - 2ir\Re[h_{n,\mathbf{a}}(x+ir)]}{(x-z_0)^2 + r^2} dx \\
 &\rightarrow 2i \int_{-\infty}^{-a_1} \frac{1}{x-z_0} \lim_{r \rightarrow 0^+} \Im[h_{n,\mathbf{a}}(x+ir)] dx \\
 &= -2i \sum_{\ell=1}^{n-1} \int_{-a_{\ell+1}}^{-a_\ell} \frac{1}{x-z_0} \frac{\sin\left(\frac{\ell}{n}\pi\right)}{\sqrt[n]{\prod_{k=1}^n |x+a_k|}} dx \quad \text{by the limit (3.1)} \\
 &= 2i \sum_{\ell=1}^{n-1} \int_{a_{\ell+1}}^{a_\ell} \frac{1}{-t-z_0} \frac{\sin\left(\frac{\ell}{n}\pi\right)}{\sqrt[n]{\prod_{k=1}^n |a_k-t|}} dt \\
 &= 2i \sum_{\ell=1}^{n-1} \sin\left(\frac{\ell}{n}\pi\right) \int_{a_\ell}^{a_{\ell+1}} \frac{1}{t+z_0} \frac{1}{\sqrt[n]{\prod_{k=1}^n |a_k-t|}} dt
 \end{aligned}$$

as $r \rightarrow 0^+$ and $R \rightarrow \infty$, where $R(r) = \sqrt{R^2 - r^2}$. In conclusion, we obtain

$$h_{n,\mathbf{a}}(z_0) = \frac{1}{G_n(z_0 \mathbf{1} + \mathbf{a})} = \frac{1}{\pi} \sum_{\ell=1}^{n-1} \sin\left(\frac{\ell}{n}\pi\right) \int_{a_\ell}^{a_{\ell+1}} \frac{1}{t+z_0} \frac{1}{\sqrt[n]{\prod_{k=1}^n |a_k-t|}} dt.$$

The integral representation (2.1) is proved.

The rest results can be derived readily from comparing the integral representations (1.1) and (2.1) and making use of the relations, which are mentioned in Section 1 of this paper, among the Stieltjes transforms, logarithmically completely monotonic functions, and completely monotonic functions. The proof of Theorem 2.1 is complete.

References

- [1] D.V. Widder, *The Laplace Transform*, in: Princeton Mathematical Series, vol. 6, Princeton University Press, Princeton, N. J., 1941.
- [2] R.D. Atanassov, U.V. Tsoukrovski, Some properties of a class of logarithmically completely monotonic functions, *C. R. Acad. Bulgare Sci.* 41 (2) (1988) 21–23.
- [3] F. Qi, C.-P. Chen, A complete monotonicity property of the gamma function, *J. Math. Anal. Appl.* 296 (2004) 603–607. Available online at <http://dx.doi.org/10.1016/j.jmaa.2004.04.026>.
- [4] F. Qi, B.-N. Guo, Complete monotonicities of functions involving the gamma and digamma functions, *RGMI Res. Rep. Coll.* 7 (1) (2004) 63–72. Art. 8. Available online at <http://rgmia.org/v7n1.php>.
- [5] C. Berg, Integral representation of some functions related to the gamma function, *Mediterr. J. Math.* 1 (4) (2004) 433–439. Available online at <http://dx.doi.org/10.1007/s00009-004-0022-6>.
- [6] B.-N. Guo, F. Qi, A property of logarithmically absolutely monotonic functions and the logarithmically complete monotonicity of a power-exponential function, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* 72 (2) (2010) 21–30.
- [7] R.L. Schilling, R. Song, Z. Vondraček, *Bernstein Functions—Theory and Applications*, second ed., in: de Gruyter Studies in Mathematics, vol. 37, Walter de Gruyter, Berlin, Germany, 2012, Available online at <http://dx.doi.org/10.1515/9783110269338>.
- [8] C.-P. Chen, F. Qi, H.M. Srivastava, Some properties of functions related to the gamma and psi functions, *Integral Transforms Spec. Funct.* 21 (2) (2010) 153–164. Available online at <http://dx.doi.org/10.1080/10652460903064216>.

- [9] F. Qi, X.-J. Zhang, W.-H. Li, The harmonic and geometric means are Bernstein functions, *Bol. Soc. Mat. Mexicana* 22 (3) (2016) (in press). Available online at <http://dx.doi.org/10.1007/s40590-016-0085-y>.
- [10] F. Qi, B.-N. Guo, V. Čerňanová, X.-T. Shi, Explicit expressions, Cauchy products, integral representations, convexity, and inequalities of central Delannoy numbers, ResearchGate Working Paper, 2016. Available online at <http://dx.doi.org/10.13140/RG.2.1.4889.6886>.
- [11] F. Qi, X.-J. Zhang, W.-H. Li, Lévy-Khintchine representation of the geometric mean of many positive numbers and applications, *Math. Inequal. Appl.* 17 (2) (2014) 719–729. Available online at <http://dx.doi.org/10.7153/mia-17-53>.
- [12] Á. Besenyei, On complete monotonicity of some functions related to means, *Math. Inequal. Appl.* 16 (1) (2013) 233–239. Available online at <http://dx.doi.org/10.7153/mia-16-17>.
- [13] B.-N. Guo, F. Qi, On the degree of the weighted geometric mean as a complete Bernstein function, *Afr. Mat.* 26 (7) (2015) 1253–1262. Available online at <http://dx.doi.org/10.1007/s13370-014-0279-2>.
- [14] F. Qi, Integral representations and properties of Stirling numbers of the first kind, *J. Number Theory* 133 (7) (2013) 2307–2319. Available online at <http://dx.doi.org/10.1016/j.jnt.2012.12.015>.
- [15] F. Qi, B.-N. Guo, The reciprocal of the weighted geometric mean is a Stieltjes transform, ResearchGate Working Paper, 2016. Available online at <http://dx.doi.org/10.13140/RG.2.1.3024.9847>.
- [16] F. Qi, X.-J. Zhang, W.-H. Li, An elementary proof of the weighted geometric mean being a Bernstein function, *Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys.* 77 (1) (2015) 35–38.
- [17] F. Qi, X.-J. Zhang, W.-H. Li, An integral representation for the weighted geometric mean and its applications, *Acta Math. Sin. (Engl. Ser.)* 30 (1) (2014) 61–68. Available online at <http://dx.doi.org/10.1007/s10114-013-2547-8>.
- [18] F. Qi, X.-J. Zhang, W.-H. Li, Lévy-Khintchine representations of the weighted geometric mean and the logarithmic mean, *Mediterr. J. Math.* 11 (2) (2014) 315–327. Available online at <http://dx.doi.org/10.1007/s00009-013-0311-z>.