




ELSEVIER

Contents lists available at ScienceDirect

Linear Algebra and its Applications

www.elsevier.com/locate/laa

Leonard pairs and quantum algebra $U_q(sl_2)$ 

Man Sang, Suogang Gao, Bo Hou*

College of Mathematics and Information Science, Hebei Normal University,
Shijiazhuang, 050024, PR China

ARTICLE INFO

Article history:

Received 28 January 2016

Accepted 29 August 2016

Available online 1 September 2016

Submitted by R. Brualdi

MSC:

17B37

05E30

33C45

33D45

Keywords:

Leonard pair

Quantum algebra

Equitable generator

LB-TD pair

ABSTRACT

Let \mathbb{K} denote an algebraically closed field of characteristic zero. Let V denote a vector space over \mathbb{K} with finite positive dimension. A *Leonard pair* on V is an ordered pair of linear transformations in $\text{End}(V)$ such that for each of these transformations there exists a basis for V with respect to which the matrix representing that transformation is diagonal and the matrix representing the other transformation is irreducible tridiagonal. Fix a nonzero scalar $q \in \mathbb{K}$ which is not a root of unity. Consider the quantum algebra $U_q(sl_2)$ with equitable generators $x^{\pm 1}, y, z$. Let d denote a nonnegative integer and let $V_{d,1}$ denote an irreducible $U_q(sl_2)$ -module of dimension $d+1$ and of type 1. In this paper, we determine all linear transformations A in $\text{End}(V_{d,1})$ such that on $V_{d,1}$, the pair A, x^{-1} , the pair A, y and the pair A, z are all Leonard pairs.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Leonard pairs were introduced by Terwilliger [10] to extend the algebraic approach of Bannai and Ito [4] to a result of D. Leonard concerning the sequences of orthogonal polynomials with discrete support for which there is a dual sequence of orthogonal

* Corresponding author.

E-mail address: houbu1969@163.com (B. Hou).

polynomials. Because these polynomials frequently arise in connection with the finite-dimensional representations of good Lie algebras and quantum groups, it is natural to find Leonard pairs associated with these algebraic objects. Leonard pairs of Krawtchouk type have been described in [8,11] using split basis and normalized semisimple generators of sl_2 . Leonard pairs of q -Krawtchouk type have been described in [11] using split basis of $U_q(sl_2)$. Recently, Alnajjar and Curtin [1] gave general construction of Leonard pairs of Racah, Hahn, dual Hahn and Krawtchouk type using equitable basis of sl_2 . Alnajjar [2,3] gave general construction of Leonard pairs of q -Racah, q -Hahn, dual q -Hahn, q -Krawtchouk, dual q -Krawtchouk, quantum q -Krawtchouk, and affine q -Krawtchouk type using equitable generators of $U_q(sl_2)$. Equitable presentations for sl_2 and $U_q(sl_2)$ were introduced in [5] and [7], respectively.

In this paper we describe a relationship between Leonard pairs and quantum algebra $U_q(sl_2)$ that appears to be new. Let \mathbb{K} denote an algebraically closed field of characteristic zero. Fix a nonzero scalar $q \in \mathbb{K}$ which is not a root of unity. Consider the quantum algebra $U_q(sl_2)$ with equitable generators $x^{\pm 1}$, y , z . Let d denote a nonnegative integer and let $V_{d,1}$ denote an irreducible $U_q(sl_2)$ -module of dimension $d+1$ and of type 1. We determine all linear transformations A in $\text{End}(V_{d,1})$ such that on $V_{d,1}$, the pair A, x^{-1} , the pair A, y and the pair A, z are all Leonard pairs.

2. Preliminaries

In this section we recall the definitions and some related facts concerning Leonard pairs and the quantum algebra $U_q(sl_2)$.

Throughout this paper \mathbb{K} will denote an algebraically closed field of characteristic zero.

2.1. Leonard pairs

In this subsection we recall some terms of Leonard pairs.

Let d be a nonnegative integer. Let \mathbb{K}^{d+1} denote the \mathbb{K} -vector space consisting of the column vectors of length $d+1$, and let $\text{Mat}_{d+1}(\mathbb{K})$ denote the \mathbb{K} -algebra consisting of the $(d+1) \times (d+1)$ matrices. The algebra $\text{Mat}_{d+1}(\mathbb{K})$ acts on \mathbb{K}^{d+1} by left multiplication.

Let V denote a \mathbb{K} -vector space of dimension $d+1$. Let $\text{End}(V)$ denote the \mathbb{K} -algebra consisting of all linear transformations from V to V . Let $\{v_i\}_{i=0}^d$ denote a basis for V . For $A \in \text{End}(V)$ and $X \in \text{Mat}_{d+1}(\mathbb{K})$, we say X represents A with respect to $\{v_i\}_{i=0}^d$ whenever $Av_j = \sum_{i=0}^d X_{ij}v_i$ for $0 \leq j \leq d$.

Let X be a square matrix. X is said to be *upper* (resp. *lower*) *bidiagonal* whenever every nonzero entry appears on or immediately above (resp. below) the main diagonal. X is said to be *tridiagonal* whenever every nonzero entry appears on, immediately above, or immediately below the main diagonal. Assume X is tridiagonal. Then X is said to be *irreducible* whenever all entries immediately above and below the main diagonal are nonzero.

Definition 2.1. [10, Definition 1.1] By a *Leonard pair* on V , we mean an ordered pair $A : V \rightarrow V$ and $A^* : V \rightarrow V$ that satisfy both (i), (ii) below.

- (i) There exists a basis for V with respect to which the matrix representing A is irreducible tridiagonal, and the matrix representing A^* is diagonal.
- (ii) There exists a basis for V with respect to which the matrix representing A^* is irreducible tridiagonal, and the matrix representing A is diagonal.

Let A, A^* be a Leonard pair on V . Obviously, A^*, A is also a Leonard pair on V .

Let A, A^* be a Leonard pair on V . By [10, Lemma 1.3] each of A, A^* has mutually distinct $d + 1$ eigenvalues. Let $\{\theta_i\}_{i=0}^d$ denote an ordering of the eigenvalues of A . For $0 \leq i \leq d$ pick an eigenvector $v_i \in V$ of A associated with θ_i . Then the ordering $\{\theta_i\}_{i=0}^d$ is said to be *standard* whenever the basis $\{v_i\}_{i=0}^d$ satisfies Definition 2.1(ii). A standard ordering of A^* is similarly defined. Note that for a standard ordering $\{\theta_i\}_{i=0}^d$ of the eigenvalues of A , the ordering $\{\theta_{d-i}\}_{i=0}^d$ is also standard and no further ordering is standard. A similar result applies to A^* . Let $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^d$) be a standard ordering of the eigenvalues of A (resp. A^*). By [10, Theorem 1.9] the expressions

$$\frac{\theta_{i-2} - \theta_{i+1}}{\theta_{i-1} - \theta_i}, \quad \frac{\theta_{i-2}^* - \theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*} \quad (1)$$

are equal and independent of i for $2 \leq i \leq d - 1$. Let β be one less the common value of (1). We call β the *fundamental parameter* of A, A^* . Let q be a nonzero scalar such that $\beta = q^2 + q^{-2}$. We call q a *quantum parameter* of A, A^* [9].

Lemma 2.2. Let A, A^* be a Leonard pair on V with quantum parameter q that is not a root of unity. Let $\{\theta_i\}_{i=0}^d$ (resp. $\{\theta_i^*\}_{i=0}^d$) be a standard ordering of the eigenvalues of A (resp. A^*). Then there exist scalars $\alpha, \alpha^*, a, a', b, b'$ such that

$$\theta_i = \alpha + aq^{2i-d} + a'q^{d-2i} \quad (0 \leq i \leq d), \quad (2)$$

$$\theta_i^* = \alpha^* + bq^{2i-d} + b'q^{d-2i} \quad (0 \leq i \leq d). \quad (3)$$

Proof. Immediate from [10, Lemma 9.2]. \square

Definition 2.3. By a *Leonard pair* in $\text{Mat}_{d+1}(\mathbb{K})$, we mean an ordered pair A, A^* in $\text{Mat}_{d+1}(\mathbb{K})$ that acts on \mathbb{K}^{d+1} as a Leonard pair.

Definition 2.4. [9, Definition 1.4] An ordered pair of matrices A, A^* in $\text{Mat}_{d+1}(\mathbb{K})$ is said to be *LB-TD* whenever A is lower bidiagonal with subdiagonal entries all 1 and A^* is irreducible tridiagonal.

Definition 2.5. [9, Definition 1.5] A Leonard pair on V is said to have *LB-TD form* whenever there exists a basis for V with respect to which the matrices representing A, A^* form an LB-TD pair in $\text{Mat}_{d+1}(\mathbb{K})$.

Consider the following LB-TD pair in $\text{Mat}_{d+1}(\mathbb{K})$:

$$A = \begin{pmatrix} \theta_0 & & & & 0 \\ 1 & \theta_1 & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ & & & \ddots & \theta_{d-1} \\ 0 & & & & 1 & \theta_d \end{pmatrix}, \quad A^* = \begin{pmatrix} x_0 & y_1 & & & \\ z_1 & x_1 & y_2 & & \\ & z_2 & \ddots & \ddots & \\ & & \ddots & \ddots & \\ & & & \ddots & x_{d-1} & y_d \\ & & & & z_d & x_d \end{pmatrix}. \quad (4)$$

Lemma 2.6. [9, Proposition 1.7] Fix a nonzero scalar q that is not a root of unity. Let $\alpha, \alpha^*, a, a', b, b', c$ be scalars with $c \neq 0$. Define scalars $\{\theta_i\}_{i=0}^d, \{x_i\}_{i=0}^d, \{y_i\}_{i=1}^d, \{z_i\}_{i=1}^d$ by

$$\theta_i = \alpha + aq^{2i-d} + a'q^{d-2i}, \quad (5)$$

$$x_i = \alpha^* + (b + b')q^{d-2i} + a'cq^{d-2i}(q^{d+1} + q^{-d-1} - q^{d-2i-1} - q^{d-2i+1}), \quad (6)$$

$$y_i = (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})(b - a'cq^{d-2i+1})(b' - a'cq^{d-2i+1})c^{-1}, \quad (7)$$

$$z_i = -cq^{d-2i+1}. \quad (8)$$

Then the matrices A, A^* in (4) form an LB-TD Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$ if and only if the scalars a, a', b, b', c satisfy the following inequalities:

$$a \notin \{a'q^{2d-2}, a'q^{2d-4}, \dots, a'q^{2-2d}\}, \quad (9)$$

$$b \notin \{b'q^{2d-2}, b'q^{2d-4}, \dots, b'q^{2-2d}\}, \quad (10)$$

$$bc^{-1}, b'c^{-1} \notin \{aq^{d-1}, aq^{d-3}, \dots, aq^{1-d}\} \cup \{a'q^{d-1}, a'q^{d-3}, \dots, a'q^{1-d}\}. \quad (11)$$

Lemma 2.7. [9, Theorem 1.10] Consider sequences of scalars $\{\theta_i\}_{i=0}^d, \{x_i\}_{i=0}^d, \{y_i\}_{i=1}^d, \{z_i\}_{i=1}^d$ such that $y_i z_i \neq 0$ for $1 \leq i \leq d$, and consider the matrices A, A^* in (4). Assume that A, A^* is a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$ with quantum parameter q that is not a root of unity. Then, after replacing q with q^{-1} if necessary, there exist scalars $\alpha, \alpha^*, a, a', b, b', c$ with $c \neq 0$ that satisfy (5)–(11).

Definition 2.8. [13, Definition 22.1] By a *parameter array* over \mathbb{K} of diameter d , we mean a sequence of scalars

$$(\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d, \{\varphi_i\}_{i=1}^d, \{\phi_i\}_{i=1}^d)$$

taken from \mathbb{K} that satisfies the following conditions:

- (PA1) $\theta_i \neq \theta_j$, $\theta_i^* \neq \theta_j^*$ if $i \neq j$ ($0 \leq i, j \leq d$).
 (PA2) $\varphi_i \neq 0$, $\phi_i \neq 0$ ($1 \leq i \leq d$).
 (PA3) $\varphi_i = \phi_1 \sum_{h=0}^{i-1} \frac{\theta_h - \theta_{d-h}}{\theta_0 - \theta_d} + (\theta_i^* - \theta_0^*)(\theta_{i-1} - \theta_d)$ ($1 \leq i \leq d$).
 (PA4) $\phi_i = \varphi_1 \sum_{h=0}^{i-1} \frac{\theta_h - \theta_{d-h}}{\theta_0 - \theta_d} + (\theta_i^* - \theta_0^*)(\theta_{d-i+1} - \theta_0)$ ($1 \leq i \leq d$).
 (PA5) The expressions

$$\frac{\theta_{i-2} - \theta_{i+1}}{\theta_{i-1} - \theta_i}, \quad \frac{\theta_{i-2}^* - \theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*}$$

are equal and independent of i for $2 \leq i \leq d-1$.

Lemma 2.9. [10, Corollary 14.2] Let B and B^* denote matrices in $\text{Mat}_{d+1}(\mathbb{K})$ of the form

$$B = \begin{pmatrix} \theta_0 & & & & 0 \\ 1 & \theta_1 & & & \\ & 1 & \theta_2 & & \\ & & \ddots & \ddots & \\ 0 & & & 1 & \theta_d \end{pmatrix}, \quad B^* = \begin{pmatrix} \theta_0^* & \varphi_1 & & & 0 \\ & \theta_1^* & \varphi_2 & & \\ & & \theta_2^* & \ddots & \\ & & & \ddots & \varphi_d \\ 0 & & & & \theta_d^* \end{pmatrix}.$$

Then the following are equivalent:

- (i) B, B^* is a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$.
 (ii) There exist scalars $\phi_1, \phi_2, \dots, \phi_d$ in \mathbb{K} such that the sequence $(\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d, \{\varphi_i\}_{i=1}^d, \{\phi_i\}_{i=1}^d)$ is a parameter array.

Lemma 2.10. [12, Theorem 1.5] Let A, A^* denote a Leonard pair on V . Then there exists a sequence of scalars $\beta, \gamma, \gamma^*, \varrho, \varrho^*, \omega, \eta, \eta^*$ taken from \mathbb{K} such that both

$$A^2 A^* - \beta A A^* A + A^* A^2 - \gamma(A A^* + A^* A) - \varrho A^* = \gamma^* A^2 + \omega A + \eta I, \quad (12)$$

$$A^{*2} A - \beta A^* A A^* + A A^{*2} - \gamma^*(A^* A + A A^*) - \varrho^* A = \gamma A^{*2} + \omega A^* + \eta^* I. \quad (13)$$

The sequence is uniquely determined by the pair A, A^* provided the dimension of V is at least 4.

We refer to (12) and (13) as the Askey–Wilson relations. We call the sequence $\beta, \gamma, \gamma^*, \varrho, \varrho^*, \omega, \eta, \eta^*$ satisfying (12) and (13) an Askey–Wilson parameter sequence for A, A^* .

Lemma 2.11. [9, Lemma 5.3] With reference to Lemma 2.2, assume $\alpha = 0$ and $\alpha^* = 0$. Then there exists some scalar ξ in \mathbb{K} such that the Askey–Wilson parameter sequence for A, A^* is as follows.

$$\begin{aligned}
\gamma &= 0, & \gamma^* &= 0, \\
\varrho &= -aa'(q^2 - q^{-2})^2, & \varrho^* &= -bb'(q^2 - q^{-2})^2, \\
\omega &= (q - q^{-1})^2((q^{d+1} + q^{-d-1})\xi - (a + a')(b + b')), \\
\eta &= -(q - q^{-1})(q^2 - q^{-2})((a + a')\xi - aa'(b + b')(q^{d+1} + q^{-d-1})), \\
\eta^* &= -(q - q^{-1})(q^2 - q^{-2})((b + b')\xi - bb'(a + a')(q^{d+1} + q^{-d-1})).
\end{aligned}$$

2.2. Quantum algebra $U_q(sl_2)$

For the rest of this paper fix a nonzero scalar $q \in \mathbb{K}$ which is not a root of unity. In this subsection we recall some facts concerning irreducible finite-dimensional $U_q(sl_2)$ -modules.

Definition 2.12. [6] The quantum algebra $U_q(sl_2)$ is the \mathbb{K} -algebra with generators $e, f, k^{\pm 1}$ satisfying the following conditions:

$$kk^{-1} = k^{-1}k = 1, \quad ke = q^2ek, \quad kf = q^{-2}fk, \quad ef - fe = \frac{k - k^{-1}}{q - q^{-1}}.$$

We call $e, f, k^{\pm 1}$ the Chevalley generators for $U_q(sl_2)$.

Lemma 2.13. [7, Theorem 2.1] The quantum algebra $U_q(sl_2)$ is isomorphic to the unital associative \mathbb{K} -algebra with generators $x^{\pm 1}, y, z$ and the relations $xx^{-1} = 1, x^{-1}x = 1,$

$$\frac{qxy - q^{-1}yx}{q - q^{-1}} = 1, \quad \frac{qyz - q^{-1}zy}{q - q^{-1}} = 1, \quad \frac{qzx - q^{-1}xz}{q - q^{-1}} = 1.$$

We call $x^{\pm 1}, y, z$ the equitable generators for $U_q(sl_2)$.

Lemma 2.14. [7, Lemma 4.2] For each $\epsilon \in \{1, -1\}$, there is an irreducible finite-dimensional $U_q(sl_2)$ -module $V_{d,\epsilon}$ with basis v_0, v_1, \dots, v_d and action

$$\begin{aligned}
\epsilon xv_i &= q^{d-2i}v_i \quad (0 \leq i \leq d), \\
\epsilon yv_i &= q^{2i-d}v_i + (q^{-d} - q^{2i+2-d})v_{i+1} \quad (0 \leq i \leq d-1), \\
\epsilon yv_d &= q^d v_d, \\
\epsilon zv_i &= (q^d - q^{2i-2-d})v_{i-1} + q^{2i-d}v_i \quad (1 \leq i \leq d), \\
\epsilon zv_0 &= q^{-d}v_0.
\end{aligned}$$

We call ϵ the type of the module. Since the module $V_{d,-1}$ can be treated similar to module $V_{d,1}$, we shall prove our results only for the module $V_{d,1}$.

Definition 2.15. Let v_0, v_1, \dots, v_d be the basis for $V_{d,1}$ from [Lemma 2.14](#). Denote the basis v_0, v_1, \dots, v_d by $[x]_{\text{row}}$.

Definition 2.16. [[14, Definition 10.2](#)] Let K_q denote the diagonal matrix in $\text{Mat}_{d+1}(\mathbb{K})$ with (i, i) -entry q^{d-2i} for $0 \leq i \leq d$.

Definition 2.17. [[14, Definition 10.7](#)] Let E_q denote the upper bidiagonal matrix in $\text{Mat}_{d+1}(\mathbb{K})$ with (i, i) -entry q^{2i-d} for $0 \leq i \leq d$ and $(i-1, i)$ -entry $q^d - q^{2i-2-d}$ for $1 \leq i \leq d$.

Definition 2.18. [[14, Definition 10.4](#)] We define a matrix $S \in \text{Mat}_{d+1}(\mathbb{K})$ as follows. For $0 \leq i, j \leq d$ the (i, j) -entry is $\delta_{i+j,d}$.

Lemma 2.19. Consider the elements x, y, z of $U_q(\mathfrak{sl}_2)$. Then the matrices that represent these elements with respect to $[x]_{\text{row}}$ are $K_q, SE_{q^{-1}}S, E_q$, respectively, where K_q is from [Definition 2.16](#), E_q is from [Definition 2.17](#) and S is from [Definition 2.18](#).

Proof. Immediate from [Lemma 2.14](#). \square

3. Main result

In this section we give the main result in this paper. The main result is [Theorem 3.10](#). We start with the following assumption.

Assumption 3.1. Let $V_{d,1}$ be an irreducible module of $U_q(\mathfrak{sl}_2)$ of dimension $d+1$. Let $[x]_{\text{row}}$ be the basis from [Definition 2.15](#). Recall that with respect to $[x]_{\text{row}}$, the matrices respecting the actions of x, y, z on $V_{d,1}$ are $K_q, SE_{q^{-1}}S, E_q$, respectively. Let A be a linear transformation on $V_{d,1}$, and let B denote the matrix representing A with respect to $[x]_{\text{row}}$. Assume that on $V_{d,1}$ the pair A, x^{-1} , the pair A, y and the pair A, z are all Leonard pairs.

Note 3.2. With reference to [Assumption 3.1](#), for any scalars ζ, ξ in \mathbb{K} with $\zeta \neq 0$, on $V_{d,1}$ the pair $\zeta A + \xi I, x^{-1}$, the pair $\zeta A + \xi I, y$ and the pair $\zeta A + \xi I, z$ are all Leonard pairs. Here I denotes the identity.

Lemma 3.3. With reference to [Assumption 3.1](#), the matrix B is of irreducible tridiagonal. Moreover, the eigenvalues of A are of the form

$$\alpha^* + bq^{2i-d} + b'q^{d-2i} \quad (0 \leq i \leq d),$$

where α^*, b, b' are some scalars in \mathbb{K} .

Proof. Consider the Leonard pair x^{-1}, A on $V_{d,1}$. Recall that K_q^{-1} is the matrix representing the action of x^{-1} on $V_{d,1}$ with respect to the basis $[x]_{row}$. Denote the diagonal entries $(K_q^{-1})_{i,i}$ of K_q^{-1} by θ_i for $0 \leq i \leq d$. Note that K_q^{-1} is a diagonal matrix and its diagonal entries θ_i are of the form

$$\theta_i = q^{2i-d} \quad (0 \leq i \leq d). \quad (14)$$

Comparing (2) and (14), we find $\{\theta_i\}_{i=0}^d$ is a standard ordering of the eigenvalues of the action of x^{-1} on $V_{d,1}$. By this and the comments below Definition 2.1, the basis $[x]_{row}$ satisfies Definition 2.1(ii). So, the matrix B is of irreducible tridiagonal. By Lemma 2.2, there exist some scalars α^* , b , b' such that the ordering of the eigenvalues of A is the same as in (3), and hence the results hold. \square

Definition 3.4. With reference to Assumption 3.1 and Lemma 3.3, write

$$B = \begin{pmatrix} x_0 & y_1 & & & \\ z_1 & x_1 & y_2 & & \\ & z_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot & y_d \\ & & & & z_d & x_d \end{pmatrix},$$

where x_i , y_i , z_i are some scalars in \mathbb{K} with $y_i z_i \neq 0$ for $1 \leq i \leq d$.

Lemma 3.5. With reference to Assumption 3.1 and Definition 3.4, the Leonard pairs y, A and z, A on $V_{d,1}$ have LB-TD form. Moreover, there exist some scalars b , b' , c , α^* in \mathbb{K} such that the following hold.

$$x_i = \alpha^* + (b + b')q^{d-2i} \quad (0 \leq i \leq d), \quad (15)$$

$$y_i = (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})(q^{-d} - q^{2i-d})^{-1}bb'c^{-1} \quad (1 \leq i \leq d), \quad (16)$$

$$z_i = -c(q^{-2i+1} - q) \quad (1 \leq i \leq d), \quad (17)$$

where, b , b' , c satisfy the following conditions.

$$b \notin \{b'q^{2d-2}, b'q^{2d-4}, \dots, b'q^{2-2d}\}, \quad (18)$$

$$bc^{-1}, b'c^{-1} \notin \{q^{d-1}, q^{d-3}, \dots, q^{1-d}\}. \quad (19)$$

Proof. We first prove the Leonard pair y, A on $V_{d,1}$ has LB-TD form. Recall that v_0, v_1, \dots, v_d is the basis $[x]_{row}$ for $V_{d,1}$ from Definition 2.15, with respect to which the matrix representing the action of y is $SE_{q^{-1}}S$ and the matrix representing A is B . Set $\alpha_i = k_0 k_1 \cdots k_i$ for $0 \leq i \leq d$, where $k_0 = 1$, $k_i = q^{-d} - q^{2i-d}$ ($1 \leq i \leq d$). Then with

respect to the basis $\alpha_0 v_0, \alpha_1 v_1, \dots, \alpha_d v_d$, the matrices representing the actions of y and A are

$$\begin{pmatrix} \theta_0 & & & & 0 \\ 1 & \theta_1 & & & \\ & 1 & \cdot & & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot \\ 0 & & & 1 & \theta_d \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x_0 & k_1 y_1 & & & \\ z_1/k_1 & x_1 & k_2 y_2 & & \\ & z_2/k_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & k_d y_d \\ & & & & z_d/k_d & x_d \end{pmatrix}, \quad (20)$$

respectively, where $\theta_i = q^{2i-d}$ ($0 \leq i \leq d$). By the above arguments and [Definition 2.4](#), we find the matrices in (20) form a LB-TD Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$, and hence the Leonard pair y, A on $V_{d,1}$ has LB-TD form.

Next, we show the entries in the matrix B are as in (15)–(17). By [Lemma 2.6](#) and since $\theta_i = q^{2i-d}$ ($0 \leq i \leq d$) and the matrices in (20) form a LB-TD Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$, there exist some scalars α^*, b, b', c with $c \neq 0$ such that

$$x_i = \alpha^* + (b + b')q^{d-2i} \quad (0 \leq i \leq d), \quad (21)$$

$$k_i y_i = (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})bb'c^{-1} \quad (1 \leq i \leq d), \quad (22)$$

$$z_i/k_i = -cq^{d-2i+1} \quad (1 \leq i \leq d), \quad (23)$$

where the scalars b, b', c satisfy the conditions (18), (19). Obviously, (15)–(17) follow from (21)–(23).

Finally, we prove the Leonard pair z, A on $V_{d,1}$ has LB-TD form. Recall that E_q is the matrix representing the action of z with respect to the basis v_0, v_1, \dots, v_d . Set $\beta_i = l_0 l_1 \cdots l_i$ for $0 \leq i \leq d$, where $l_0 = 1, l_i = q^d - q^{d-2i}$ ($1 \leq i \leq d$). Then with respect to the basis $\beta_0 v_d, \beta_1 v_{d-1}, \dots, \beta_d v_0$, the matrices representing the actions of z and A are

$$\begin{pmatrix} \theta_d & & & & 0 \\ 1 & \theta_{d-1} & & & \\ & 1 & \cdot & & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot \\ 0 & & & 1 & \theta_0 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} x_d & l_1 z_d & & & \\ y_d/l_1 & x_{d-1} & l_2 z_{d-1} & & \\ & y_{d-1}/l_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & l_d z_1 \\ & & & & y_1/l_d & x_0 \end{pmatrix}, \quad (24)$$

respectively, where $\theta_{d-i} = q^{d-2i}$ ($0 \leq i \leq d$). By the above arguments and [Definition 2.4](#), we find the matrices in (24) form a LB-TD Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$, and hence the Leonard pair z, A on $V_{d,1}$ has LB-TD form. \square

Definition 3.6. Define the matrix $C \in \text{Mat}_{d+1}(\mathbb{K})$ to be

$$C = \begin{pmatrix} x_0 & y_1 & & & \\ z_1 & x_1 & y_2 & & \\ & z_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & y_d \\ & & & & z_d & x_d \end{pmatrix},$$

where

$$x_i = (b + b')q^{d-2i} \quad (0 \leq i \leq d), \quad (25)$$

$$y_i = (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})(q^{-d} - q^{2i-d})^{-1}bb'c^{-1} \quad (1 \leq i \leq d), \quad (26)$$

$$z_i = -c(q^{-2i+1} - q) \quad (1 \leq i \leq d), \quad (27)$$

and b, b', c are some scalars in \mathbb{K} satisfying the following conditions.

$$b \notin \{b'q^{2d-2}, b'q^{2d-4}, \dots, b'q^{2-2d}\}, \quad (28)$$

$$bc^{-1}, b'c^{-1} \notin \{q^{d-1}, q^{d-3}, \dots, q^{1-d}\}. \quad (29)$$

Proposition 3.7. With reference to [Definitions 2.17, 2.18 and 3.6](#), the pair of matrix $SE_{q^{-1}}S, C$ forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$.

Proof. Define the matrix $P_1 \in \text{Mat}_{d+1}(\mathbb{K})$ to be

$$P_1 = \begin{pmatrix} k_0 & & & & 0 \\ & k_0k_1 & & & \\ & & \cdot & & \\ & & & \cdot & \\ & & & & \cdot \\ 0 & & & & & k_0k_1 \cdots k_d \end{pmatrix},$$

where $k_0 = 1$, $k_i = q^{-d} - q^{2i-d}$ ($1 \leq i \leq d$). Then the matrices $P_1^{-1}SE_{q^{-1}}SP_1$ and $P_1^{-1}CP_1$ are of the form

$$\begin{pmatrix} \theta_0 & & & & 0 \\ 1 & \theta_1 & & & \\ & 1 & \cdot & & \\ & & \cdot & \cdot & \\ & & & \cdot & \cdot \\ 0 & & & & 1 & \theta_d \end{pmatrix}, \quad \begin{pmatrix} x_0 & k_1y_1 & & & \\ z_1/k_1 & x_1 & k_2y_2 & & \\ & z_2/k_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & k_dy_d \\ & & & & z_d/k_d & x_d \end{pmatrix},$$

respectively, where $\theta_i = q^{2i-d}$ ($0 \leq i \leq d$). Moreover, using (25)–(27), we find the entries of matrix $P_1^{-1}CP_1$ are as follows.

$$\begin{aligned} x_i &= (b + b')q^{d-2i} \quad (0 \leq i \leq d), \\ k_i y_i &= (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})bb'c^{-1} \quad (1 \leq i \leq d), \\ z_i/k_i &= -cq^{d-2i+1} \quad (1 \leq i \leq d). \end{aligned}$$

Then by (28), (29) and Lemma 2.6, the LB-TD pair $P_1^{-1}SE_{q^{-1}}SP_1, P_1^{-1}CP_1$ forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$, and hence $SE_{q^{-1}}S, C$ is a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$. \square

Proposition 3.8. *With reference to Definitions 2.17 and 3.6, the pair of matrix E_q, C forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$.*

Proof. Define the matrix P_2 in $\text{Mat}_{d+1}(\mathbb{K})$ to be

$$P_2 = \begin{pmatrix} l_0 & & & & 0 \\ & l_0 l_1 & & & \\ & & l_0 l_1 l_2 & & \\ & & & \ddots & \\ 0 & & & & l_0 l_1 \cdots l_d \end{pmatrix},$$

where $l_0 = 1, l_i = q^d - q^{d-2i}$ ($1 \leq i \leq d$). Then the matrices $P_2^{-1}SE_qSP_2$ and $P_2^{-1}SCSP_2$ are, respectively, of the form

$$\begin{pmatrix} \theta_d & & & & 0 \\ 1 & \theta_{d-1} & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ 0 & & & 1 & \theta_0 \end{pmatrix}, \quad \begin{pmatrix} x_d & l_1 z_d & & & \\ y_d/l_1 & x_{d-1} & l_2 z_{d-1} & & \\ & y_{d-1}/l_2 & \ddots & \ddots & \\ & & \ddots & \ddots & l_d z_1 \\ & & & y_1/l_d & x_0 \end{pmatrix},$$

where $\theta_{d-i} = q^{d-2i}$ ($0 \leq i \leq d$). Moreover, using (25)–(27), we find the entries of matrix $P_2^{-1}SCSP_2$ are as follows.

$$x_{d-i} = (b + b')q^{2i-d} \quad (0 \leq i \leq d), \quad (30)$$

$$z_{d-i+1}l_i = -c(q^{-d+i-1} - q^{d-i+1})(q^i - q^{-i}) \quad (1 \leq i \leq d), \quad (31)$$

$$y_{d-i+1}/l_i = -q^{-d+2i-1}bb'c^{-1} \quad (1 \leq i \leq d). \quad (32)$$

Replace q with q^{-1} and $bb'c^{-1}$ with c_1 in (30)–(32). Then $\theta_{d-i} = q^{2i-d}$ ($0 \leq i \leq d$) and (30)–(32) become

$$\begin{aligned}x_{d-i} &= (b + b')q^{d-2i} \quad (0 \leq i \leq d), \\z_{d-i+1}l_i &= -(q^{-d+i-1} - q^{d-i+1})(q^i - q^{-i})bb'c_1^{-1} \quad (1 \leq i \leq d), \\y_{d-i+1}/l_i &= -c_1q^{d-2i+1} \quad (1 \leq i \leq d),\end{aligned}$$

respectively. Note that $bc_1^{-1} = (b'c^{-1})^{-1}$ and $b'c_1^{-1} = (bc^{-1})^{-1}$. Then by (28), (29) and Lemma 2.6, the LB-TD pair $P_2^{-1}SE_qSP_2, P_2^{-1}SCSP_2$ forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$, and hence E_q, C is a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$. \square

Proposition 3.9. *With reference to Definitions 2.16 and 3.6, the pair of matrix K_q^{-1}, C forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$.*

Proof. Define a lower triangular matrix P in $\text{Mat}_{d+1}(\mathbb{K})$ as follows:

$$P_{i,0} = q^{(2-i)(d-1)}b'^{2-i}c^{i-2}$$

for $0 \leq i \leq d$ and

$$P_{i,j} = q^{(2-i)(d-1)}b'^{2-i}c^{i-2} \prod_{k=1}^j (q^{2i-d} - q^{2k-2-d})$$

for $1 \leq j \leq i \leq d$, where b', c are from Definition 3.6. Then the matrices $P^{-1}K_q^{-1}P$ and $P^{-1}CP$ are of the form

$$\begin{pmatrix} \theta_0 & & & & 0 \\ 1 & \theta_1 & & & \\ & 1 & \ddots & & \\ & & \ddots & \ddots & \\ 0 & & & 1 & \theta_d \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} \theta_0^* & \varphi_1 & & & 0 \\ & \theta_1^* & \varphi_2 & & \\ & & \ddots & \ddots & \\ & & & \ddots & \varphi_d \\ 0 & & & & \theta_d^* \end{pmatrix},$$

respectively, where $\theta_i = q^{2i-d}$, $\theta_i^* = bq^{2i-d} + b'q^{d-2i}$ for $0 \leq i \leq d$ and $\varphi_i = (q^i - q^{-i})(q^{i-d-1} - q^{d-i+1})bq^{2i-d-1}$ for $1 \leq i \leq d$. Let ϕ_i be the scalar $(q^i - q^{-i})(q^{i-d-1} - q^{d-i+1})b'q^{d-2i+1}$ for $1 \leq i \leq d$. By calculation, we find the sequence $(\{\theta_i\}_{i=0}^d, \{\theta_i^*\}_{i=0}^d, \{\varphi_i\}_{i=1}^d, \{\phi_i\}_{i=1}^d)$ satisfies Conditions (PA1)–(PA5) in Definition 2.8, and hence it is a parameter array. So the pair of matrix $P^{-1}K_q^{-1}P, P^{-1}CP$ forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$ by Lemma 2.9, which implies that the pair of matrix K_q^{-1}, C also forms a Leonard pair in $\text{Mat}_{d+1}(\mathbb{K})$. \square

Theorem 3.10. *Recall $V_{d,1}$ is an irreducible $U_q(\mathfrak{sl}_2)$ -module of dimension $d+1$ and $[x]_{\text{row}}$ is the basis for $V_{d,1}$ from Definition 2.15. Let $A \in \text{End}(V_{d,1})$. Then on $V_{d,1}$ the pair A, x^{-1} , the pair A, y and the pair A, z are all Leonard pairs if and only if with respect to $[x]_{\text{row}}$ the matrix representing A is of the form*

$$\begin{pmatrix} x_0 & y_1 & & & \\ z_1 & x_1 & y_2 & & \\ & z_2 & \cdot & \cdot & \\ & & \cdot & \cdot & \cdot \\ & & & \cdot & \cdot & y_d \\ & & & & z_d & x_d \end{pmatrix},$$

where

$$\begin{aligned} x_i &= \alpha^* + (b + b')q^{d-2i} \quad (0 \leq i \leq d), \\ y_i &= (q^i - q^{-i})(q^{d-i+1} - q^{i-d-1})(q^{-d} - q^{2i-d})^{-1}bb'c^{-1} \quad (1 \leq i \leq d), \\ z_i &= -c(q^{-2i+1} - q) \quad (1 \leq i \leq d), \end{aligned}$$

for some scalars α^* , b , b' and nonzero scalar c in \mathbb{K} , which satisfy the following conditions

$$\begin{aligned} b &\notin \{b'q^{2d-2}, b'q^{2d-4}, \dots, b'q^{2-2d}\}, \\ bc^{-1}, b'c^{-1} &\notin \{q^{d-1}, q^{d-3}, \dots, q^{1-d}\}. \end{aligned}$$

Proof. The only if part is from [Lemma 3.5](#) and the if part is from [Propositions 3.7, 3.8, 3.9](#) and [Note 3.2](#). \square

We finish this paper with some comments.

Lemma 3.11. *With reference to [Assumption 3.1](#), on $V_{d,1}$, the pair A, x is not a Leonard pair.*

Proof. Assume that on $V_{d,1}$, the pair A, x is a Leonard pair for a contradiction. Then by [Theorem 3.10](#), the matrix representing A with respect to the basis $[x]_{row}$ is of the form in [Theorem 3.10](#). By [Note 3.2](#), we assume that $\alpha^* = 0$. Namely, C is the matrix representing A with respect to the basis $[x]_{row}$. Recall that K_q is the matrix representing the action of x on $V_{d,1}$ with respect to the basis $[x]_{row}$. Note that $\{bq^{2i-d} + b'q^{d-2i}\}_{i=0}^d$ (resp. $\{q^{d-2i}\}_{i=0}^d$) is a standard ordering of the eigenvalues of A (resp. the action of x). Then by [Lemmas 2.10, 2.11](#), we obtain

$$K_q^2 C - (q^2 + q^{-2})K_q C K_q + C K_q^2 = \omega K_q + \eta I, \quad (33)$$

$$C^2 K_q - (q^2 + q^{-2})C K_q C + K_q C^2 - \varrho^* K_q = \omega C + \eta^* I, \quad (34)$$

where

$$\varrho^* = -bb'(q^2 - q^{-2})^2, \quad (35)$$

$$\omega = (q - q^{-1})^2((q^{d+1} + q^{-d-1})\xi - (b + b')), \quad (36)$$

$$\eta = -(q - q^{-1})(q^2 - q^{-2})\xi, \quad (37)$$

$$\eta^* = -(q - q^{-1})(q^2 - q^{-2})((b + b')\xi - bb'(q^{d+1} + q^{-d-1})) \quad (38)$$

and ξ is some scalar in \mathbb{K} .

For $1 \leq i \leq d$, comparing the $(i-1, i)$ -entry of both sides of (34) and simplifying the result by using (36), we obtain

$$\begin{aligned} & (q - q^{-1})^2(q^{d+1} + q^{-d-1})\xi \\ &= (b + b')(q^{2d-4i+4} - q^{2d-4i+6} + q^{2d-4i} - q^{2d-4i-2} + q^2 + q^{-2} - 2), \end{aligned}$$

which forces $b + b' = 0$ and $\xi = 0$. So $\omega = 0$ by (36) and $x_i = 0$ for $0 \leq i \leq d$ by (25).

For $1 \leq i \leq d$, comparing the (i, i) -entry of both sides of (34) by using the results $\omega = 0$ and $x_i = 0$ for $0 \leq i \leq d$, we obtain

$$z_i y_i (2q^{d-2i} - (q^2 + q^{-2})q^{d-2i+2}) + y_{i+1} z_{i+1} (2q^{d-2i} - (q^2 + q^{-2})q^{d-2i-2}) = \varrho^* q^{d-2i} + \eta^*. \quad (39)$$

Substituting $y_i, z_i, \varrho^*, \eta^*$ in (39) using (26), (27), (35) and (38), respectively, and simplifying the result, we obtain

$$\begin{aligned} & bb'(q - q^{-1})(q^2 - q^{-2})(q^{d+1} + q^{-d-1}) \\ &= -bb'q^{d-2i}(q^{2d-4i+6} - q^{2d-4i+2} - q^{2d-4i-2} + q^{2d-4i-6} - q^{2d-2i+6} + q^{2d-2i+2} \\ & \quad + q^{2d-2i} - q^{2d-2i-4} - q^{-2i+4} + q^{-2i} + q^{-2i-2} - q^{-2i-6}), \end{aligned}$$

which forces $bb' = 0$. By the above arguments, we find that $b = b' = 0$, which contradicts to the assumption that A, x is a Leonard pair on $V_{d,1}$. \square

Lemma 3.12. With reference to Assumption 3.1 and Theorem 3.10, on $V_{d,1}$,

$$A = \alpha^* 1 + (b + b')x + \frac{c}{q - q^{-1}}[x, y] + \frac{bb'c^{-1}}{q - q^{-1}}[z, x]. \quad (40)$$

Proof. Let $[x]_{row}$ be the basis for $V_{d,1}$ from Definition 2.15. Compare the matrix representing the action of each side of (40) with respect to the basis $[x]_{row}$. We find both sides are equal to the matrix $\alpha^* I + C$, where I denotes the identity and C is from Definition 3.6. \square

Acknowledgements

The authors are grateful to professor P. Terwilliger and professor T. Ito for the advice they offered during their study of Leonard pairs and tridiagonal pairs. This work was supported by the NSF of China (No. 11271257 and No. 11471097), the Specialized Research Fund for the Doctoral Program of Higher Education of China (No. 20121303110005).

References

- [1] H. Alnajjar, B. Curtin, Leonard pairs from the equitable basis of sl_2 , *Electron. J. Linear Algebra* 20 (2010) 490–505.
- [2] H. Alnajjar, Leonard pairs from the equitable generators of $U_q(sl_2)$, *Dirasat Pure Sci.* 37 (2010) 31–35.
- [3] H. Alnajjar, Leonard pairs associated with the equitable generators of the quantum algebra $U_q(sl_2)$, *Linear Multilinear Algebra* 59 (10) (2011) 1127–1142.
- [4] E. Bannai, T. Ito, *Algebraic Combinatorics I: Association Schemes*, Benjamin/Cummings, London, 1984.
- [5] G. Benkart, P. Terwilliger, The equitable basis for sl_2 , *Math. Z.* 268 (2008) 535–557.
- [6] C. Kassel, *Quantum Groups*, Springer-Verlag, New York, 1995.
- [7] T. Ito, P. Terwilliger, C.W. Weng, The quantum algebra $U_q(sl_2)$ and its equitable presentation, *J. Algebra* 298 (2006) 284–301.
- [8] K. Nomura, P. Terwilliger, Krawtchouk polynomials, the Lie algebra sl_2 , and Leonard pairs, *Linear Algebra Appl.* 437 (1) (2012) 345–375.
- [9] K. Nomura, Leonard pair having LB-TD form, *Linear Algebra Appl.* 455 (2014) 1–21.
- [10] P. Terwilliger, Two linear transformations each tridiagonal with respect to an eigenbasis of the other, *Linear Algebra Appl.* 330 (2001) 149–203.
- [11] P. Terwilliger, Introduction to Leonard pairs, OPSFA Rome 2001, *J. Comput. Appl. Math.* 153 (2003) 463–475.
- [12] P. Terwilliger, R. Vidunas, Leonard pairs and the Askey–Wilson relations, *J. Algebra Appl.* 3 (2004) 411–426.
- [13] P. Terwilliger, An algebraic approach to the Askey scheme of orthogonal polynomials, in: *Orthogonal Polynomials and Special Functions: Computation and Applications*, in: *Lecture Notes in Math.*, vol. 1883, Springer, Berlin, 2006, pp. 255–330.
- [14] P. Terwilliger, Finite-dimensional irreducible $U_q(sl_2)$ -modules from the equitable point of view, *Linear Algebra Appl.* 439 (2013) 358–400.