

Expert Q&A;: Are Steroid Tapers Always Necessary?

Stopping steroids abruptly can backfire. Our clinical board explains safe tapering.

By **Living With Arthritis Editorial Team** · Reviewed by **Clinical Review Board** (Multidisciplinary UK clinicians) · 2026-06-14 · livingwitharthritis.org.uk

Overview



Openly licensed image via Openverse

Stopping steroids abruptly can backfire. Our clinical board explains safe tapering.

What the evidence shows

UK guidance from NICE, the British Society for Rheumatology and Versus Arthritis consistently emphasises a combined approach: appropriate medication, regular movement, and supportive lifestyle changes. Recent reviews continue to back this consensus.

What this means for you



Openly licensed image via Openverse

- ****Talk to your clinician.**** Personalised advice always beats blanket rules.
- ****Track your symptoms.**** Use a simple diary so you can spot patterns over weeks, not days.

- ****Move within comfort.**** Gentle, regular activity protects joints and mood.
 - ****Watch the warning signs.**** New swelling, fever, or rapid loss of function deserve same-week review.
-

Practical next steps

1. Read our companion piece for a deeper dive: [\[related guidance\]\(/conditions/rheumatoid-arthritis\)](#). 2. Bring three questions to your next GP or rheumatology appointment. 3. Share this article with a family member who supports you day to day.

Editorial standards



Openly licensed image via Openverse

This article was prepared by the Living With Arthritis Editorial Team and reviewed by our Clinical Review Board against UK NICE guidance and peer-reviewed sources. We update content at least annually.

What the Evidence Says

Current evidence from systematic reviews consistently highlights the risks associated with suddenly stopping long-term steroid treatment. These reviews, which pull together findings from many individual studies, show that people can experience a flare-up of their underlying condition or withdrawal symptoms if they reduce their dose too quickly. Clinical guidelines across the UK strongly recommend a gradual reduction, tailored to the individual's condition and previous dose.

Randomised trials, considered a strong form of evidence, have explored different tapering schedules for various conditions. While there's no single "best" way to taper for everyone, these studies generally support a slow, step-by-step reduction rather than abrupt cessation. This approach helps the body adjust and reduces the chances of negative side effects.

The consensus from expert opinion and published research is clear: managing steroid reduction safely requires careful planning and oversight from a healthcare professional. Self-managing a steroid taper without medical guidance significantly increases the risk of complications, including adrenal suppression, where your body temporarily stops producing enough of its own natural steroids.

Common Mistakes to Avoid

* **Stopping steroids purely because you feel better:** Even if symptoms improve, your body still needs to adjust to a lower dose. * **Cutting your dose in half all at once:** This can be too big a step and trigger withdrawal or a flare. * **Listening to advice from friends or online forums over your doctor:** Everyone's situation is unique; what worked for someone else might not be right for you. * **Forgetting to pick up your next prescription for a lower dose:** This can lead to an unplanned abrupt stop. * **Ignoring withdrawal symptoms like fatigue or joint pain, hoping they'll pass:** These are signals your body might be reducing too quickly.

When to Speak to a Clinician

It's always best to discuss any concerns about your medication with a healthcare professional. If you're on steroids and experience new or worsening symptoms, or if you're struggling with side effects, don't wait. Contact your doctor, rheumatology nurse, or pharmacist for advice.

* Sudden increase in pain or stiffness * Extreme fatigue or weakness * Feeling generally unwell or feverish * Significant mood changes, like anxiety or feeling low * Developing new, unexplained symptoms

Frequently Asked Questions

What happens if I stop steroids suddenly? Stopping steroids abruptly, especially after taking them for a long time, can lead to withdrawal symptoms or a flare-up of your arthritis. Your body might not be able to produce enough of its own natural steroids, leading to issues like extreme tiredness, muscle pain, nausea, and even low blood pressure.

How long does a steroid taper usually take? The length of a steroid taper varies a lot depending on how long you've been taking them, the dose, and your particular condition. It could range from a few weeks to several months, or even longer for some people. Your doctor will create a personalised plan for you.

Can I drink alcohol while tapering steroids? Moderate alcohol consumption is generally acceptable for most people while tapering steroids, but it's wise to discuss this with your doctor. Some people find alcohol can worsen certain side effects, like stomach upset, or interfere with their sleep, which is important for recovery.

Q&A

Question: What is the key thing to know about expert q&a;: are steroid tapers always necessary??

Short answer: Stopping steroids abruptly, especially after long-term use, can cause withdrawal symptoms or a flare-up. Your body may not produce enough natural steroids, causing extreme tiredness, muscle pain, nausea, or low blood pressure. Guidelines

recommend a gradual reduction, tailored by a healthcare professional to your condition and previous dose, to help your body adjust.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Versus Arthritis: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.

General information only — not medical advice. Consult your GP or specialist before changing medication or starting a new programme. Emergency: 999. NHS advice: 111. Living With Arthritis UK · livingwitharthritis.org.uk