

Building Strength and Resilience: Exercise for Frailty Prevention

Exercise is one of the most powerful tools for preventing and reversing frailty. Here's how to build a safe, effective routine at any fitness level.

By **Maxwell** · Reviewed by **Maxwell** (First Contact Practitioner, HCPC PH128483, CSP Member) · 2026-06-21 · livingwitharthritis.org.uk

Regular physical activity is consistently shown to be one of the most effective tools for preventing and even reversing frailty. Exercise helps maintain muscle mass and strength, improves balance and coordination, supports bone density, and contributes to better cardiovascular health — all of which directly counter the physiological decline that defines frailty. The good news is that effective exercise doesn't require a gym, special equipment, or even high intensity. Consistent, appropriate movement at any level makes a meaningful difference.



Openly licensed image via Openverse

Why Exercise Is Central to Frailty Prevention

Muscles are responsive to demand. When challenged regularly, they grow stronger and maintain mass. When underused, they weaken and shrink. This means exercise is not optional for maintaining strength and independence — it's fundamental. Research consistently shows that regular exercise can slow, halt, or even partially reverse frailty, even in people who are already significantly frail.

Beyond the direct physical benefits, exercise improves mood, supports sleep, and contributes to better cognitive function. It also builds confidence and a sense of control, which can be psychologically important for someone managing a chronic condition.

The Four Pillars of Frailty-Prevention Exercise



Openly licensed image via Openverse

An effective exercise routine for frailty prevention typically includes four components, and ideally a programme combines all four rather than focusing on just one.

****Strength training**** maintains muscle mass and strength. This can involve resistance bands, light weights, or bodyweight exercises. The key is working against resistance that challenges the muscles.

****Balance exercises**** reduce fall risk and improve confidence in movement. These might include standing on one leg, heel-to-toe walking, or tai chi.

****Flexibility work**** maintains range of motion, which is essential for function and quality of life. Gentle stretching or activities like yoga can help.



Openly licensed image via Openverse

****Light aerobic activity**** supports cardiovascular health and overall endurance. Walking, swimming, or stationary cycling are accessible options for most people.

Getting Started Safely

For someone who is frail or has been inactive for a while, starting slowly and building gradually is essential. This might mean beginning with seated exercises, very light resistance, and short walking sessions, then progressively increasing duration and intensity as strength and confidence improve.

Working with a physical therapist initially can help ensure exercises are appropriate and performed safely, particularly for anyone with significant frailty, balance concerns, or multiple health conditions. An initial assessment can identify specific limitations and areas of focus.

If you have a medical condition, particularly heart disease, diabetes, or a history of stroke, it's important to get medical clearance before starting a new exercise programme.

Sample Approach for Beginners

A simple starting routine might include two to three short sessions per week, each lasting 20-30 minutes. A session might include:

A few minutes of seated exercises, such as seated marching (lifting knees while sitting), arm circles, and arm lifts with light weights or resistance bands.

A few balance exercises performed near a sturdy support, such as standing on one leg for 10-20 seconds, or heel-to-toe walking along a line.

Gentle stretching, holding each stretch for 15-30 seconds without bouncing.

A short walk, perhaps 5-10 minutes, indoors or outdoors as tolerated.

As strength and confidence build, the duration, resistance, and complexity of exercises can gradually increase. Someone who starts with 5-minute walks might build to 15-minute walks within a few weeks. Resistance can be increased by using slightly heavier weights or stronger resistance bands.

Specific Strength Exercises

Exercises that target major muscle groups are particularly important:

****Sit-to-stands****: Standing up from a chair and sitting back down (with support available if balance is poor). This builds leg strength essential for mobility.

****Leg lifts****: Lying on your back or side and lifting one leg straight up. Strengthens leg muscles.

****Wall push-ups****: Standing facing a wall and pushing against it with your hands. A gentler option than floor push-ups.

****Rowing movements****: Using a resistance band or light weights, pulling toward your body. Strengthens back and arms.

These exercises can be done in the home, require minimal or no equipment, and can be modified to match current fitness level.

Staying Consistent and Motivated

Consistency matters more than intensity, particularly in the early stages. Exercising with a friend, joining a group class designed for older adults, or working with a physical therapist or trainer can improve both safety and motivation. Some gyms or community centres offer exercise classes specifically for older or frail adults.

Tracking small improvements — an extra minute of walking, an easier time standing up from a chair, improved balance — can provide encouraging evidence of progress, even when changes feel gradual.

Setting realistic goals helps maintain motivation. Rather than "get fit," a more achievable goal might be "walk for 10 minutes without stopping" or "rise from a chair without using my hands."

When to Seek Professional Guidance

A doctor should be consulted before starting if you have uncontrolled high blood pressure, significant heart disease, have had a recent heart attack or stroke, or have other significant medical concerns.

A physical therapist can be invaluable for someone with significant frailty, multiple health conditions, or previous falls. They can design an individualised programme, ensure exercises are done safely, and progress the programme appropriately.

Bringing It All Together

The research is clear: it's rarely too late to benefit from exercise, even for adults who are already frail. Starting where you are, building gradually, and staying consistent offers one of the most powerful, accessible paths toward greater strength, independence, and resilience.

Some studies have shown that older adults can regain significant strength and function within 8-12 weeks of starting an appropriate exercise programme. Others have shown that people in their 80s and 90s can still build meaningful muscle with training. The key is to begin, progress gradually, and stay consistent.

****This article is for general information only and is not a substitute for professional medical advice. Always consult with your healthcare provider before starting a new exercise programme, particularly if you have medical conditions, take multiple medications, or have concerns about your balance or cardiovascular health.****

Q&A

Question: What is the key thing to know about building strength and resilience: exercise for frailty prevention?

Short answer: Regular physical activity helps prevent and reverse frailty by maintaining muscle, improving balance, and supporting bone density. An effective programme combines strength, balance, flexibility, and light aerobic exercises. Start slowly and build gradually — a physical therapist can guide you. Consistency matters; even small improvements make a difference. Consult a doctor before starting if you have existing medical conditions.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Arthritis UK: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.

General information only — not medical advice. Consult your GP or specialist before changing medication or starting a new programme. Emergency: 999. NHS advice: 111. Living With Arthritis · livingwitharthritis.org.uk