

Omega-3 and Fish Oil for Arthritis: A Complete Evidence-Base

Evidence-based guide to omega-3 fatty acids and fish oil for arthritis. Learn about benefits, correct dosage, best food sources and supplement recommendations.

By **Living With Arthritis Editorial Team** · Pending clinical review ·
2026-02-11 · livingwitharthritis.org.uk



*Fresh salmon fillet with omega-3 rich foods including walnuts, flaxseeds and avocado on a wooden cutting board
Openly licensed image via Openverse*

Omega-3 fatty acids — particularly those found in oily fish and fish oil supplements — are among the most extensively researched natural interventions for arthritis. With

evidence spanning decades and hundreds of clinical trials, omega-3s have established themselves as a legitimate complementary therapy for managing joint inflammation and pain. This comprehensive guide examines the science, practical applications and evidence-based recommendations for using omega-3 fatty acids to support arthritis management.

Understanding Omega-3 Fatty Acids

The Three Main Types

- EPA (Eicosapentaenoic acid) — The most potent anti-inflammatory omega-3. EPA directly competes with arachidonic acid (a pro-inflammatory omega-6 fatty acid) for the same enzymes, effectively reducing the production of inflammatory prostaglandins and leukotrienes. This is the form most relevant to arthritis management
- DHA (Docosahexaenoic acid) — Important for brain health, cardiovascular function and cell membrane integrity. DHA also has anti-inflammatory properties, though less potent than EPA for joint inflammation specifically. It produces specialised pro-resolving mediators (SPMs) that actively help resolve inflammation
- ALA (Alpha-linolenic acid) — Found in plant sources like flaxseed, chia seeds and walnuts. The body can convert ALA to EPA and DHA, but the conversion rate is very low (approximately 5–10% to EPA, less than 1% to DHA). While ALA has its own health benefits, it's not an efficient source of the EPA/DHA needed for anti-inflammatory effects


How Omega-3s Fight Inflammation

Omega-3 fatty acids combat inflammation through multiple mechanisms that are directly relevant to arthritis:

- Reducing pro-inflammatory eicosanoids — EPA displaces arachidonic acid in cell membranes, reducing the production of inflammatory prostaglandins (PGE2) and thromboxanes
- Producing anti-inflammatory resolvins and protectins — EPA and DHA are converted into specialised pro-resolving mediators that actively shut down inflammatory pathways
- Inhibiting inflammatory gene expression — Omega-3s suppress the NF-κB pathway and reduce the production of pro-inflammatory cytokines (TNF-α, IL-1β, IL-6) — the same targets as many arthritis medications

- Reducing cartilage-degrading enzymes — Studies show omega-3s can reduce levels of matrix metalloproteinases (MMPs) that break down cartilage in arthritic joints

The Evidence for Arthritis



Openly licensed image via Openverse

Rheumatoid Arthritis

The evidence for omega-3s in rheumatoid arthritis (RA) is strong and has been building for over 30 years:

- A Cochrane systematic review of 23 studies found that omega-3 fish oil supplements significantly reduced joint tenderness, morning stiffness duration and NSAID use in RA patients

- Multiple RCTs show that fish oil supplementation (at doses of 2.7g+ EPA/DHA daily) can reduce the need for anti-inflammatory medication — some patients were able to reduce their NSAID dose by 30–50%
- A meta-analysis in the British Journal of Nutrition confirmed significant improvements in joint pain intensity, morning stiffness and number of painful joints
- Benefits typically take 2–3 months to become apparent, with maximum effects at 3–6 months

Osteoarthritis

Evidence for osteoarthritis (OA) is more limited but growing:

- A systematic review in Osteoarthritis and Cartilage found some evidence that omega-3s reduce pain in OA, though effect sizes were smaller than for RA
 - Laboratory studies show omega-3s may protect cartilage by reducing degradative enzyme activity and supporting cartilage cell function
 - A large randomised trial found that high-dose fish oil (4.5g/day) was not significantly superior to low-dose (0.45g/day) for knee OA symptoms, suggesting that moderate doses may be sufficient
 - The anti-inflammatory benefits of omega-3s may be most relevant during OA flare-ups when inflammatory activity increases
-

Best Food Sources of Omega-3

The NHS recommends eating at least two portions of fish per week, one of which should be oily. The richest sources include:

- Salmon — Wild Atlantic salmon provides approximately 2.2g EPA+DHA per 140g serving. Widely available fresh, frozen and tinned in UK supermarkets
- Mackerel — One of the richest sources at approximately 2.7g per 140g serving. Smoked mackerel is a convenient UK staple
- Sardines — Approximately 1.8g per 140g serving. Tinned sardines are affordable and available everywhere
- Herring and kippers — Traditional British fish providing about 2.0g per serving
- Anchovies — Rich in omega-3s and widely used in Mediterranean cooking

- Fresh tuna — Note that tinned tuna has most of its omega-3s removed during processing; fresh or frozen is much richer
- Trout — A good freshwater alternative at about 1.2g per serving

Plant Sources (ALA)

For vegetarians, vegans or those who don't eat fish, plant-based ALA sources include flaxseeds (ground) and flaxseed oil, chia seeds, hemp seeds, walnuts, and rapeseed (canola) oil. Remember that ALA conversion to EPA/DHA is very limited. Algae-based EPA/DHA supplements are available for those who avoid fish.

Supplement Guide



Openly licensed image via Openverse

Dosage Recommendations

- General health — 250–500 mg combined EPA+DHA daily (achievable through diet)
- Arthritis management — 2,000–3,000 mg (2–3g) combined EPA+DHA daily. This is the dose range used in most positive clinical trials for RA. Higher doses (up to 4g) have been used in some studies but should be discussed with your GP
- Look for high-EPA formulations — Since EPA is the most anti-inflammatory omega-3, choose supplements with a higher EPA-to-DHA ratio for arthritis

Types of Supplements Available in the UK

- Standard fish oil capsules — The most affordable option. Check the label for actual EPA+DHA content per capsule (not just 'fish oil' amount). Many standard capsules contain only 300mg EPA+DHA per 1000mg fish oil, meaning you'd need 7–10 capsules daily for therapeutic doses
- Concentrated fish oil — Higher EPA+DHA per capsule (often 600–900mg per capsule), requiring fewer capsules daily. More expensive but more practical
- Liquid fish oil — Flavoured liquid forms (lemon or orange) make it easy to get higher doses in a single teaspoon
- Algae-based omega-3 — Suitable for vegetarians and vegans. Available from Holland & Barrett, health food shops and online. Typically provides 250–500mg EPA+DHA per capsule
- Krill oil — Contains omega-3s in phospholipid form, which may be better absorbed. Also contains astaxanthin (antioxidant). Usually more expensive and provides lower total EPA+DHA per capsule

Quality Considerations

- Look for products tested for heavy metals (mercury, lead) and contaminants (PCBs, dioxins)
- Choose brands with IFOS (International Fish Oil Standards) certification or similar third-party testing
- Check the 'best before' date — rancid fish oil is ineffective and may be harmful
- Store capsules in a cool, dark place or refrigerator after opening
- If capsules cause fishy burps, try taking them with meals, using enteric-coated versions, or switching to liquid form

Safety and Interactions

- Blood thinning — High-dose omega-3s (above 3g/day) may increase bleeding time. If you take warfarin, aspirin or other anticoagulants, discuss fish oil supplementation with your GP. Most doctors consider doses up to 3g/day safe alongside blood thinners, but monitoring may be recommended
- Surgery — Some surgeons recommend stopping fish oil supplements 1–2 weeks before planned surgery due to potential bleeding effects
- Drug interactions — Fish oil may interact with blood pressure medications (additive blood pressure-lowering effect) and some immunosuppressive drugs
- Pregnancy — Fish oil supplementation is generally considered safe during pregnancy (and DHA is beneficial for foetal brain development), but avoid high-dose EPA supplements and fish liver oil (which contains high vitamin A) during pregnancy
- Side effects — Mild GI symptoms (fishy breath, nausea, loose stools) are the most common side effects and usually resolve with dose adjustment or taking supplements with food

An Omega-3 Rich Meal Plan

Here's a sample day incorporating omega-3 rich foods:

- Breakfast — Porridge with ground flaxseed, walnuts and blueberries
- Lunch — Smoked mackerel salad with mixed leaves, cherry tomatoes, avocado and olive oil dressing
- Dinner — Baked salmon with roasted Mediterranean vegetables, sweet potato and a side of steamed broccoli
- Snack — Handful of walnuts with an apple

Omega-3 fatty acids are one of the best-supported natural interventions for arthritis, particularly inflammatory types. By combining dietary omega-3 sources with appropriate supplementation, you can give your joints meaningful anti-inflammatory support alongside your regular medical treatment. Ask our virtual assistant for personalised omega-3 and dietary

advice for your arthritis type.

This article is for general information only. Consult your GP or pharmacist before starting high-dose fish oil supplements, particularly if you take other medications.

Q&A

Question: What is the key thing to know about omega-3 and fish oil for arthritis: a complete evidence-base?

Short answer: Omega-3 fatty acids, particularly EPA, help manage arthritis by reducing inflammation. They combat inflammation by displacing pro-inflammatory fats, producing anti-inflammatory molecules, and inhibiting inflammatory gene expression. For rheumatoid arthritis, take 2,000–3,000 mg of combined EPA+DHA daily, which can reduce pain, stiffness, and potentially lower the need for NSAIDs. Benefits appear within 2 to 3 months.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Arthritis UK: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.

General information only — not medical advice. Consult your GP or specialist before changing medication or starting a new programme. Emergency: 999. NHS advice: 111. Living With Arthritis · livingwitharthritis.org.uk