

Expert Brief: Exercise Order for Maximum Pain Relief

Mobility before strength before cardio: the order that produces the largest reduction in arthritis pain scores. A physiotherapist explains why.

By **Living With Arthritis Editorial Team** · Reviewed by **Clinical Review Board** (HCPC-registered physiotherapists, BDA-registered dietitians) · 2026-06-04 · livingwitharthritis.org.uk

What this guide covers



Openly licensed image via Openverse

This piece summarises what the current evidence says about **expert brief: exercise order for maximum pain relief**, the practical steps that help most, and the red flags that mean you should speak to your GP or physiotherapist. It is written for people living in the UK and reviewed by our clinical team.

The underlying mechanism

Arthritis pain rarely has a single cause. Joint cartilage, synovial fluid, ligaments, surrounding muscle and the nervous system all contribute. Understanding which of these is driving your specific symptom is the first step toward targeted relief. For an in-depth condition overview, see our [\[related guide\]\(/exercises\)](#).

Inflammatory and degenerative forms behave differently. Inflammatory arthritis (rheumatoid, psoriatic, ankylosing spondylitis) typically worsens with rest and improves with gentle movement. Degenerative arthritis (osteoarthritis) typically worsens with prolonged use. Knowing which pattern matches your day-to-day experience helps you choose the right management approach.



Openly licensed image via Openverse

What helps — by strength of evidence

1. ****Movement and graded exercise.**** NICE NG226 (2022) and Cochrane reviews place structured exercise as a first-line treatment. Aim for 150 minutes of moderate activity per week, split into 20–30 minute sessions. 2. ****Anti-inflammatory eating.**** A Mediterranean-style pattern — oily fish twice a week, plenty of vegetables, olive oil, whole grains — is linked to reduced pain scores in OA and RA trials. 3. ****Weight management.**** A 5 percent reduction in body weight reduces knee-OA symptoms measurably; a 10 percent reduction can match the effect of some medications. 4. ****Heat and cold.**** Heat eases stiffness; cold eases acute swelling. Twenty-minute applications, with a towel barrier. 5. ****Sleep hygiene.**** Pain perception rises sharply on fewer than six hours of sleep. Prioritise a consistent bedtime.

A practical seven-day starting plan

Begin with the smallest dose that feels manageable — five minutes of movement, one anti-inflammatory meal a day, a 10 pm screen-off rule. Track only three things weekly: pain on waking (0–10), longest pain-free interval, and one functional win (e.g., climbed stairs without grabbing the rail). Small, consistent inputs outperform big sporadic efforts.



Openly licensed image via Openverse

When to seek clinical advice

Book an appointment with your GP or a first-contact physiotherapist if you notice: sudden hot, red, swollen joints; unexplained weight loss alongside joint pain; morning stiffness lasting longer than an hour for more than two weeks; or new numbness or weakness. These can signal inflammatory disease that benefits from early treatment.

Frequently asked questions

****Will this approach work alongside my medication?**** Yes. Exercise, diet and pacing complement prescribed treatments and often improve their effectiveness. Always check with your prescriber before stopping any medication.

****How long until I notice a difference?**** Most people report meaningful improvement at 6–12 weeks of consistent practice. Earlier wins (less morning stiffness, better sleep) often appear within 2–3 weeks.

****Is this safe if I have other conditions?**** The principles above are broadly safe for adults with stable cardiovascular, metabolic and respiratory conditions. If you are unsure, ask your GP for a referral to a first-contact physiotherapist for a tailored plan.

This article is general health information, not medical advice. Always consult a qualified healthcare professional for decisions about your individual care.

Q&A

Question: What is the key thing to know about expert brief: exercise order for maximum pain relief?

Short answer: The best exercise order for pain relief depends on the type of arthritis. Inflammatory arthritis improves with gentle movement, while degenerative arthritis may worsen with prolonged use. Structured exercise, specifically 150 minutes of moderate activity per week, is recommended. Start slowly, aiming for 20-30 minute sessions.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Arthritis UK: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.