

Understanding Frailty in Older Adults

Frailty is more than just getting older — it's a recognised medical syndrome. Here's what it means, what causes it, and how it can be managed.

By **Maxwell** · Reviewed by **Maxwell** (First Contact Practitioner, HCPC PH128483, CSP Member) · 2026-06-21 · livingwitharthritis.org.uk

Frailty is more than just getting older. It's a recognised medical syndrome characterised by reduced strength, endurance, and physiological reserve that leaves a person vulnerable to illness, injury, and disability following even minor stressors. Many people remain robust well into their eighties and nineties, while frailty can sometimes develop earlier when multiple health conditions overlap.



What Does 'Frailty' Actually Mean?

Frailty is a recognised medical syndrome characterised by reduced strength, endurance, and physiological reserve, which leaves a person more vulnerable to illness, injury, and disability following even relatively minor stressors, such as an infection or a fall. It's distinct from simply being older — many people remain robust well into their eighties and nineties, while frailty can sometimes develop earlier when multiple health conditions overlap.

Clinically, frailty is often identified using a set of measurable signs: unintentional weight loss, exhaustion, weakness (often measured by grip strength), slow walking speed, and low physical activity levels. Meeting several of these criteria suggests a frailty syndrome rather than normal ageing. A healthcare professional can assess these markers and determine whether someone is experiencing frailty, pre-frailty, or neither.

The distinction is important because it changes how healthcare decisions should be made. A frail person may respond differently to surgery, medication, or hospitalisation than a non-frail person of the same age, and may need additional support to recover from illness.



Openly licensed image via Openverse

What Causes Frailty?

Frailty develops from an interplay of multiple factors rather than a single cause. Age-related muscle loss, known as sarcopenia, is a fundamental driver. Starting around age 30, most people lose approximately 3-8% of their muscle mass per decade, with the rate of loss accelerating significantly after age 60. This loss of muscle translates directly to loss of strength and functional capacity.

Chronic inflammation is another key factor. As we age, baseline levels of inflammatory markers in the blood tend to increase, a phenomenon researchers call "inflammaging." This chronic, low-level inflammation damages tissues over time and contributes to both muscle loss and weakness.

Hormonal changes also play a role. Testosterone and growth hormone levels decline with age, and these hormones are important for maintaining muscle mass and strength. In women, the drop in oestrogen after menopause affects muscle strength and bone density.



Openly licensed image via Openverse

The cumulative effect of chronic diseases — heart disease, diabetes, kidney disease, chronic obstructive pulmonary disease — accelerates frailty. These conditions create a state of metabolic stress that the ageing body struggles to manage.

Poor nutrition, particularly inadequate protein intake, compounds the problem. Social isolation and inactivity create a vicious cycle: weakness leads to reduced activity, which leads to further weakness. Addressing these factors can interrupt the progression.

Why Identifying Frailty Matters

Recognising frailty early matters because it changes how healthcare decisions should be approached. A frail person undergoing surgery, for example, may need different perioperative management, longer rehabilitation, and closer monitoring. A frail person starting a new medication may experience different side effects or require dose adjustments.

More importantly, identifying frailty allows healthcare teams to be proactive rather than reactive. Instead of waiting for a fall, fracture, or hospitalisation, early recognition enables supportive interventions that can slow or even reverse the condition's progression.

Research consistently shows that frailty, particularly in its earlier stages, responds to targeted intervention. This means that identifying frailty is not a pessimistic diagnosis — it's an opportunity to intervene when intervention is most effective.

Can Frailty Be Reversed?

Importantly, frailty is not necessarily a one-way path. Research shows that frailty — particularly in its earlier stages, sometimes called "pre-frailty" — can often be improved or even reversed through targeted interventions. Resistance exercise to rebuild muscle strength, adequate protein and calorie intake, addressing underlying medical conditions, and reducing social isolation have all been shown to improve frailty markers in clinical studies.

Studies of older adults who began resistance training programmes showed meaningful strength gains and improved functional capacity even after years of decline. Some adults moved from a frailty classification back to pre-frailty or robust status. The key is consistency and starting sooner rather than later.

Supporting a Frail Loved One

If you're caring for someone who shows signs of frailty, encouraging regular movement appropriate to their ability, ensuring meals are nutritious and protein-rich, supporting social connection, and coordinating closely with their healthcare team can all make a meaningful difference. Small, consistent steps tend to be more sustainable and effective than dramatic changes.

If you notice signs of frailty in yourself or a loved one — unexplained weight loss, persistent fatigue, or a noticeable slowing of walking speed — it's worth raising this with a doctor, since early support can meaningfully improve strength, independence, and quality of life.

****This article is for general information only and is not a substitute for professional medical advice. Always consult with your healthcare provider before starting new exercise or dietary changes.****

Q&A

Question: What is the key thing to know about understanding frailty in older adults?

Short answer: Frailty is a medical syndrome characterised by reduced strength, endurance, and physiological reserve, making individuals vulnerable to illness or injury. It is different from normal ageing, as many older people remain robust. Clinically, frailty is identified by unintentional weight loss, exhaustion, weakness, slow walking speed, and low physical activity. Early identification is crucial for effective intervention and improved outcomes.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Arthritis UK: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.

General information only — not medical advice. Consult your GP or specialist before changing medication or starting a new programme. Emergency: 999. NHS advice: 111. Living With Arthritis · livingwitharthritis.org.uk