

Steroid Injections vs Oral Steroids: When Each Makes Sense

Targeted joint injections and short oral courses each have a role. We unpack the trade-offs.

By **Living With Arthritis Editorial Team** · Reviewed by **Clinical Review Board** (Multidisciplinary UK clinicians) · 2026-06-13 · livingwitharthritis.org.uk

Overview



Openly licensed image via Openverse

Targeted joint injections and short oral courses each have a role. We unpack the trade-offs.

What the evidence shows

UK guidance from NICE, the British Society for Rheumatology and Arthritis UK consistently emphasises a combined approach: appropriate medication, regular movement, and supportive lifestyle changes. Recent reviews continue to back this consensus.

What this means for you




Openly licensed image via Openverse

- **Talk to your clinician.** Personalised advice always beats blanket rules.
 - **Track your symptoms.** Use a simple diary so you can spot patterns over weeks, not days.
 - **Move within comfort.** Gentle, regular activity protects joints and mood.
 - **Watch the warning signs.** New swelling, fever, or rapid loss of function deserve same-week review.
-

Practical next steps

1. Read our companion piece for a deeper dive: [\[related guidance\]\(/conditions/osteoarthritis\)](#).
 2. Bring three questions to your next GP or rheumatology appointment.
 3. Share this article with a family member who supports you day to day.
-

Editorial standards



This article was prepared by the Living With Arthritis Editorial Team and reviewed by our Clinical Review Board against UK NICE guidance and peer-reviewed sources. We update content at least annually.

What the Evidence Says

Systematic reviews of treatment options for arthritis consistently highlight that neither steroid injections nor oral steroids are long-term solutions. While both can be very effective in managing flare-ups and reducing pain, they don't tackle the underlying disease progression. Randomised controlled trials show that steroid injections can provide targeted pain relief for weeks to months, particularly in single inflamed joints or bursae, whereas short courses of oral steroids are generally considered for more widespread inflammation.

Clinical guidelines from professional bodies emphasise that these treatments are part of a broader management strategy. They are most beneficial when used alongside other interventions like regular exercise, physical therapy, weight management where appropriate, and other pain relief methods. The goal is to improve function and quality of life, allowing individuals to engage in activities that support long-term joint health.

The body of evidence also points to potential risks with overuse or high doses of steroids. Repeated steroid injections into the same joint, for instance, can sometimes weaken tissues over time, while prolonged oral steroid use carries a wider range of potential side effects impacting bone density, blood sugar, and susceptibility to infection. Therefore, careful consideration of benefits versus risks is always advised.

Common Mistakes to Avoid

* **Expecting a permanent cure:** Neither injections nor oral steroids stop arthritis progression. *Remember they are tools for managing symptoms and flare-ups.* * **Insisting on steroids for every ache:** Not every pain is due to inflammation that steroids will help. *Discuss other pain relief options with your clinician.* * **Stopping oral steroids suddenly:** This can cause withdrawal symptoms or a flare. *Always follow your clinician's tapering instructions carefully.* * **Ignoring lifestyle changes because of steroid relief:** Steroids offer temporary relief, not a pass to neglect movement or diet. *Continue with gentle exercise and healthy eating habits.* * **Having injections too frequently:** Frequent injections into the same

joint carry risks. *Follow your clinician's advice on injection frequency.*

When to Speak to a Clinician

It's really important to know when to seek professional advice, especially when dealing with new or worsening symptoms. While a little discomfort is part of managing arthritis, certain signals suggest it's time to get things checked out by your GP, physiotherapist, or rheumatologist.

* Sudden, severe joint pain that doesn't ease. * New or significantly increased joint swelling and redness. * Fever alongside joint symptoms, which could indicate infection. * Rapid loss of function or inability to move a joint you could previously move. * Symptoms that are getting worse despite following your current treatment plan.

Frequently Asked Questions

How long do steroid injections take to work? Typically, you might start to feel the effects of a steroid injection within a few days, though for some, it can take up to a week. The full pain relief can then last from several weeks to a few months, depending on the individual and the joint injected.

Can I have multiple steroid injections? Yes, it's possible to have multiple steroid injections, but your clinician will carefully consider the frequency and number. Too many injections into the same joint can, over time, weaken surrounding tissues, so there are usually limits on how often and how many you can receive.

Are there alternatives to steroids for arthritis flare-ups? Absolutely. Depending on the type and severity of your arthritis, alternatives might include non-steroidal anti-inflammatory drugs (NSAIDs), other pain relief medications, physical therapy, heat or cold packs, and various topical treatments. Your clinician can discuss the best options for your specific situation.

Q&A

Question: What is the key thing to know about steroid injections vs oral steroids: when each makes sense?

Short answer: Steroid injections typically alleviate pain and inflammation in specific joints for several weeks to months, often taking a few days to a week to become effective. Oral steroids are generally used for widespread inflammation. Neither is a long-term solution, and both carry risks with overuse. They should be part of a broader management strategy including exercise, therapy, and weight management.

Question: When should I seek professional advice?

Short answer: If symptoms persist or worsen, pain is severe and unresponsive to usual relief, or you develop new symptoms, contact your GP, physiotherapist or rheumatology team promptly.

Question: Where can I get free arthritis support in the UK?

Short answer: Arthritis UK: 0800 5200 520. NRAS: 0800 298 7650. Free tools and guides at livingwitharthritis.org.uk.

General information only — not medical advice. Consult your GP or specialist before changing medication or starting a new programme. Emergency: 999. NHS advice: 111. Living With Arthritis · livingwitharthritis.org.uk