

Formulae sheet for Electromagnetism

A. Operator in different coordinates

→ Cartesian coordinates:

$$\nabla = \left(\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z} \right), \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

$$d\vec{\ell} = dx\hat{x} + dy\hat{y} + dz\hat{z}, dV = dxdydz$$

→→ Gradient:

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right)$$

→→ Divergence:

$$\nabla \cdot \vec{F} = \frac{\partial F_x}{\partial x} + \frac{\partial F_y}{\partial y} + \frac{\partial F_z}{\partial z}$$

→→ Curl:

$$\nabla \times \vec{F} = \left(\frac{\partial F_z}{\partial y} - \frac{\partial F_y}{\partial z}, \frac{\partial F_x}{\partial z} - \frac{\partial F_z}{\partial x}, \frac{\partial F_y}{\partial x} - \frac{\partial F_x}{\partial y} \right)$$

→ Cylindrical coordinates:

$$\nabla = \left(\frac{\partial}{\partial r}, \frac{1}{r} \frac{\partial}{\partial \theta}, \frac{\partial}{\partial z} \right), \nabla^2 = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} + \frac{\partial^2}{\partial z^2}$$

$$d\vec{\ell} = dr\hat{r} + r d\theta\hat{\theta} + dz\hat{z}$$

$$dV = r dr d\theta dz$$

→→ Gradient:

$$\nabla f = \left(\frac{\partial f}{\partial r}, \frac{1}{r} \frac{\partial f}{\partial \theta}, \frac{\partial f}{\partial z} \right)$$

→→ Divergence:

$$\nabla \cdot \vec{F} = \frac{1}{r} \frac{\partial}{\partial r} (r F_r) + \frac{1}{r} \frac{\partial F_\theta}{\partial \theta} + \frac{\partial F_z}{\partial z}$$

→→ Curl:

$$\nabla \times \vec{F} = \left(\frac{1}{r} \frac{\partial F_z}{\partial \theta} - \frac{\partial F_\theta}{\partial z}, \frac{\partial F_r}{\partial z} - \frac{\partial F_z}{\partial r}, \right.$$

$$\left. \frac{1}{r} \frac{\partial}{\partial r} (r F_\theta) - \frac{\partial F_r}{\partial \theta} \right)$$

→ Spherical coordinates:

$$\nabla = \left(\frac{\partial}{\partial r}, \frac{1}{r} \frac{\partial}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \right)$$

$$\nabla^2 = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2 \theta} \frac{\partial^2}{\partial \phi^2}$$

$$d\vec{\ell} = dr\hat{r} + r d\theta\hat{\theta} + r \sin \theta d\phi\hat{\phi}$$

$$dV = r^2 \sin \theta dr d\theta d\phi$$

→→ Gradient:

$$\nabla f = \left(\frac{\partial f}{\partial r}, \frac{1}{r} \frac{\partial f}{\partial \theta}, \frac{1}{r \sin \theta} \frac{\partial f}{\partial \phi} \right)$$

→→ Divergence:

$$\nabla \cdot \vec{F} = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 F_r \right) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta F_\theta) + \frac{1}{r \sin \theta} \frac{\partial F_\phi}{\partial \phi}$$

→→ Curl:

$$\nabla \times \vec{F} = \frac{1}{r \sin \theta} \left(\frac{\partial}{\partial \theta} (F_\phi \sin \theta) - \frac{\partial F_\theta}{\partial \phi}, \right.$$

$$\left. \frac{1}{r} \left(\frac{\partial F_r}{\partial \phi} - \frac{\partial}{\partial r} (r F_\phi) \right), \right.$$

$$\left. \frac{1}{r} \left(\frac{\partial F_\theta}{\partial r} - \frac{\partial F_r}{\partial \theta} \right) \right)$$

B. Vector formulae

$$\begin{aligned} \rightarrow \vec{a} \cdot (\vec{b} \times \vec{c}) &= \vec{b} \cdot (\vec{c} \times \vec{a}) = \vec{c} \cdot (\vec{a} \times \vec{b}) \\ \rightarrow \vec{a} \times (\vec{b} \times \vec{d}) &= (\vec{a} \cdot \vec{d})\vec{b} - (\vec{a} \cdot \vec{b})\vec{d} \\ \rightarrow (\vec{a} \times \vec{b}) \cdot (\vec{c} \times \vec{d}) &= (\vec{a} \cdot \vec{c})(\vec{b} \cdot \vec{d}) - (\vec{a} \cdot \vec{d})(\vec{b} \cdot \vec{c}) \\ \rightarrow \nabla \times \nabla \psi &= 0 \\ \rightarrow \nabla \cdot (\nabla \times \vec{a}) &= 0 \\ \rightarrow \nabla \times (\nabla \times \vec{a}) &= \nabla(\nabla \cdot \vec{a}) - \nabla^2 \vec{a} \\ \rightarrow \nabla \cdot (\psi \vec{a}) &= \nabla \psi \cdot \vec{a} + \psi \nabla \cdot \vec{a} \\ \rightarrow \nabla \times (\psi \vec{a}) &= \nabla \psi \times \vec{a} + \psi \nabla \times \vec{a} \\ \rightarrow \nabla(\vec{a} \cdot \vec{b}) &= (\vec{a} \cdot \nabla)\vec{b} + (\vec{b} \cdot \nabla)\vec{a} + \vec{a} \times (\nabla \times \vec{b}) + \vec{b} \times (\nabla \times \vec{a}) \\ \rightarrow \nabla \cdot (\vec{a} \times \vec{b}) &= \vec{b} \cdot (\nabla \times \vec{a}) - \vec{a} \cdot (\nabla \times \vec{b}) + (\vec{b} \cdot \nabla)\vec{a} - (\vec{a} \cdot \nabla)\vec{b} \\ \rightarrow \nabla \times (\vec{a} \times \vec{b}) &= (\vec{b} \cdot \nabla)\vec{a} - (\vec{a} \cdot \nabla)\vec{b} + \vec{a}(\nabla \cdot \vec{b}) - \vec{b}(\nabla \cdot \vec{a}) \\ \rightarrow \nabla \cdot \vec{r} &= 3 \\ \rightarrow \nabla \times \vec{r} &= 0 \\ \rightarrow \nabla \cdot [\vec{r} \cdot f(r)] &= \frac{2}{r} f + \frac{\partial f}{\partial r} \\ \rightarrow \nabla \times [\vec{r} \cdot f(r)] &= 0 \\ \rightarrow (\vec{a} \cdot \nabla) \hat{r} f(r) &= \frac{f(r)}{r} [\vec{a} - \hat{r}(\vec{a} \cdot \hat{r})] + f(\vec{a} \cdot \hat{r}) \frac{\partial \hat{r}}{\partial r} \end{aligned}$$

C. Theorems from vector calculus

→ Divergence Theorem:

$$\int_V \nabla \cdot \vec{A} d^3x = \int_S \vec{A} \cdot \hat{n} dS$$

$$\int_V \nabla \psi d^3x = \int_S \psi \hat{n} dS$$

$$\int_V \nabla \times \vec{A} d^3x = \int_S \hat{n} \times \vec{A} dS$$

→ Green's First Identity:

$$\int_V (\phi \nabla^2 \psi + \nabla \phi \cdot \nabla \psi) d^3x = \int_S \phi \hat{n} \cdot \nabla \psi dS$$

→ Green's Theorem:

$$\int_V (\phi \nabla^2 \psi - \psi \nabla^2 \phi) d^3x = \int_S (\phi \nabla \psi - \psi \nabla \phi) \cdot \hat{n} dS$$

→ Stokes's Theorem:

$$\int_S (\nabla \times \vec{A}) \cdot \hat{n} dS = \oint_C \vec{A} \cdot d\vec{l}$$

$$\int_S \hat{n} \times \nabla \psi dS = \oint_C \psi d\vec{l}$$

D. Vector calculus with delta-function

→ The Dirac delta function $\delta(\vec{r})$ is a distribution that satisfies:

$$\int_{\mathbb{R}^3} \delta(\vec{r}) d^3r = 1$$

and is zero everywhere except at $\vec{r} = \vec{0}$.

→ The sifting property of the Dirac delta function allows us to extract function values:

$$\int_{\mathbb{R}^3} f(\vec{r}) \delta(\vec{r} - \vec{r}_0) d^3r = f(\vec{r}_0)$$

→ The Laplacian of the Dirac delta function in three dimensions is:

$$\nabla^2 \delta(\vec{r}) = -4\pi \delta(\vec{r})$$

→ The gradient and Laplacian of the function $\frac{1}{r}$ is:

$$\nabla \left(\frac{1}{r} \right) = -\frac{\vec{r}}{r^3}, \quad \nabla^2 \left(\frac{1}{r} \right) = -4\pi \delta(\vec{r})$$

→ The divergence of a vector field multiplied by a delta function can be expressed as:

$$\nabla \cdot (f(\vec{r}) \delta(\vec{r} - \vec{r}_0)) = \delta(\vec{r} - \vec{r}_0) \nabla f(\vec{r}_0) + f(\vec{r}_0) \nabla \delta(\vec{r} - \vec{r}_0)$$

E. Constants

→ Permittivity of free space:

$$\epsilon_0 = 8.854 \times 10^{-12} \text{ F/m}$$

→ Permeability of free space:

$$\mu_0 = 4\pi \times 10^{-7} \text{ H/m}$$

→ Speed of light in vacuum:

$$c = 3.00 \times 10^8 \text{ m/s}$$

→ Charge of an electron:

$$e = 1.602 \times 10^{-19} \text{ C}$$

→ Avogadro's number:

$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

F. Electrostatics

→ Coulomb's Law:

$$\vec{F} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \hat{r}$$

→ Electric Field:

$$\vec{E} = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\vec{r}')(\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3} dV'$$

or

$$\vec{E} = -\nabla V, \quad V(\vec{r}) = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} dV'$$

The electric field is the gradient of the electric potential.

→ Gauss's Law:

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0}$$

$$\oint_{\partial V} \vec{E} \cdot d\vec{A} = \frac{Q_{\text{enc}}}{\epsilon_0}$$

The divergence of the electric field is proportional to the charge density.

→ Poisson's Equation:

$$\nabla^2 V = -\frac{\rho}{\epsilon_0}$$

Relates the electric potential to the charge density.

→ Ohm's Law for Current Density:

$$\vec{J}_f = \sigma \vec{E} = \frac{1}{\rho} \vec{E}$$

→ Polarization:

$$\vec{P} = \epsilon_0 \chi_f \vec{E} = \epsilon_0 (\epsilon_r - 1) \vec{E}$$

Describes the polarization density in a medium.

→ Electric Displacement Field:

$$\vec{D} = \epsilon_0 \vec{E} + \vec{P} = \epsilon_r \epsilon_0 \vec{E}$$

The electric displacement field accounts for free and bound charges.

→ Surface Bound Charge Density:

$$\sigma_b = \vec{P} \cdot \hat{n}$$

Surface charge density due to polarization.

→ Volume Bound Charge Density:

$$\rho_b = -\nabla \cdot \vec{P}$$

Volume charge density due to polarization.

→ Capacitance:

$$C = \frac{\Delta Q}{\Delta V}, E = \frac{Q^2}{2C}$$

→ The current density J can be expressed as:

$$J = nqv_D$$

where $v_D = -\frac{eE}{m} \langle \tau \rangle$. $\langle \tau \rangle$ is the average time between collisions (mean free time).

G. Magnetostatics

→ Biot-Savart Law:

$$\vec{B}(\vec{r}) = \frac{\mu_0}{4\pi} \int \frac{Id\vec{\ell} \times (\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3}$$

→ Lorentz Force:

$$\vec{F} = q(\vec{v} \times \vec{B})$$

→ Ampère's Law:

$$\nabla \times \vec{B} = \mu_0 \vec{J}$$

$$\oint_{\partial S} \vec{B} \cdot d\vec{\ell} = \mu_0 I_{\text{enc}}$$

In the absence of time-varying electric fields, the curl of the magnetic field is proportional to the current density.

→ Magnetostatic Gauss's Law:

$$\nabla \cdot \vec{B} = 0$$

→ Relation between Magnetic Field and Magnetization:

$$\vec{B} = \mu_0 \vec{H} + \vec{M}$$

→ Magnetization Current Density:

$$\vec{J}_m = \nabla \times \vec{M}$$

→ Magnetic inductance:

$$L = \frac{\Delta \Phi}{\Delta I}, E = \frac{I^2 L}{2}$$

H. Electrodynamics

→ Faraday's Law of induction states that a time-varying magnetic field induces a curl in the electric field.

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t}$$

$$\mathcal{E} = \oint_{\partial S} \vec{E} \cdot d\vec{\ell} = -\frac{d}{dt} \int_S \vec{B} \cdot d\vec{A}$$

The electromotive force (EMF) around a closed loop ∂S is equal to the negative rate of change of the magnetic flux through the surface S bounded by the loop.

→ Maxwell's Equations:

$$\nabla \cdot \vec{E} = \frac{\rho_f}{\epsilon_0} \quad (\text{Gauss's Law})$$

$$\nabla \cdot \vec{B} = 0 \quad (\text{Magnetostatic Gauss's Law})$$

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \quad (\text{Faraday's Law})$$

$$\nabla \times \vec{B} = \mu_0 \vec{J}_f + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t} \quad (\text{Maxwell-Ampère Law})$$

→ Poynting Vector:

$$\vec{S} = \vec{E} \times \vec{H}$$

Represents the energy flux density of an electromagnetic field.

→ Energy Density:

$$u = \frac{1}{2} \epsilon_0 E^2 + \frac{1}{2\mu_0} B^2$$

The energy density of the electromagnetic field.

→ Electromagnetic Wave Equation:

$$\nabla^2 \vec{E} - \mu_0 \epsilon_0 \frac{\partial^2 \vec{E}}{\partial t^2} = 0$$

$$\nabla^2 \vec{B} - \mu_0 \epsilon_0 \frac{\partial^2 \vec{B}}{\partial t^2} = 0$$

Describes the propagation of electromagnetic waves in a vacuum.

→ Continuity Equation in Electrodynamics:

$$\nabla \cdot \vec{J}_f + \frac{\partial \rho_f}{\partial t} = 0$$

Table of Integrals*

Basic Forms

$$\int x^n dx = \frac{1}{n+1} x^{n+1} \quad (1)$$

$$\int \frac{1}{x} dx = \ln |x| \quad (2)$$

$$\int u dv = uv - \int v du \quad (3)$$

$$\int \frac{1}{ax+b} dx = \frac{1}{a} \ln |ax+b| \quad (4)$$

Integrals of Rational Functions

$$\int \frac{1}{(x+a)^2} dx = -\frac{1}{x+a} \quad (5)$$

$$\int (x+a)^n dx = \frac{(x+a)^{n+1}}{n+1}, n \neq -1 \quad (6)$$

$$\int x(x+a)^n dx = \frac{(x+a)^{n+1}((n+1)x-a)}{(n+1)(n+2)} \quad (7)$$

$$\int \frac{1}{1+x^2} dx = \tan^{-1} x \quad (8)$$

$$\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} \quad (9)$$

$$\int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \ln |a^2-x^2| \quad (10)$$

$$\int \frac{x^2}{a^2+x^2} dx = x - a \tan^{-1} \frac{x}{a} \quad (11)$$

$$\int \frac{x^3}{a^2+x^2} dx = \frac{1}{2} x^2 - \frac{1}{2} a^2 \ln |a^2+x^2| \quad (12)$$

$$\int \frac{1}{ax^2+bx+c} dx = \frac{2}{\sqrt{4ac-b^2}} \tan^{-1} \frac{2ax+b}{\sqrt{4ac-b^2}} \quad (13)$$

$$\int \frac{1}{(x+a)(x+b)} dx = \frac{1}{b-a} \ln \frac{a+x}{b+x}, a \neq b \quad (14)$$

$$\int \frac{x}{(x+a)^2} dx = \frac{a}{a+x} + \ln |a+x| \quad (15)$$

$$\int \frac{x}{ax^2+bx+c} dx = \frac{1}{2a} \ln |ax^2+bx+c| - \frac{b}{a\sqrt{4ac-b^2}} \tan^{-1} \frac{2ax+b}{\sqrt{4ac-b^2}} \quad (16)$$

Integrals with Roots

$$\int \sqrt{x} dx = \frac{2}{3} (x-a)^{3/2} \quad (17)$$

$$\int \frac{1}{\sqrt{x \pm a}} dx = 2\sqrt{x \pm a} \quad (18)$$

$$\int \frac{1}{\sqrt{a-x}} dx = -2\sqrt{a-x} \quad (19)$$

$$\int x\sqrt{x-ax} = \frac{2}{3} a(x-a)^{3/2} + \frac{2}{5} (x-a)^{5/2} \quad (20)$$

$$\int \sqrt{ax+bx^2} = \left(\frac{2b}{3a} + \frac{2x}{3} \right) \sqrt{ax+bx^2} \quad (21)$$

$$\int (ax+b)^{3/2} dx = \frac{2}{5a} (ax+b)^{5/2} \quad (22)$$

$$\int \frac{x}{\sqrt{x \pm a}} dx = \frac{2}{3} (x \mp 2a) \sqrt{x \pm a} \quad (23)$$

$$\int \sqrt{\frac{x}{a-x}} dx = -\sqrt{x(a-x)} - a \tan^{-1} \frac{\sqrt{x(a-x)}}{x-a} \quad (24)$$

$$\int \sqrt{\frac{x}{a+x}} dx = \sqrt{x(a+x)} - a \ln [\sqrt{x} + \sqrt{x+a}] \quad (25)$$

Integrals with Logarithms

$$\int \ln ax dx = x \ln ax - x \quad (42)$$

$$\int \frac{\ln ax}{x} dx = \frac{1}{2} (\ln ax)^2 \quad (43)$$

$$\int \ln(ax+b) dx = \left(x + \frac{b}{a} \right) \ln(ax+b) - x, a \neq 0 \quad (44)$$

$$\int \ln(x^2+a^2) dx = x \ln(x^2+a^2) + 2a \tan^{-1} \frac{x}{a} - 2x \quad (45)$$

$$\int \ln(x^2-a^2) dx = x \ln(x^2-a^2) + a \ln \frac{x+a}{x-a} - 2x \quad (46)$$

$$\int \ln(ax^2+bx+c) dx = \frac{1}{a} \sqrt{4ac-b^2} \tan^{-1} \frac{2ax+b}{\sqrt{4ac-b^2}} - 2x + \left(\frac{b}{2a} + x \right) \ln(ax^2+bx+c) \quad (47)$$

$$\int x \ln(ax+b) dx = \frac{bx}{2a} - \frac{1}{4} x^2 + \frac{1}{2} \left(x^2 - \frac{b^2}{a^2} \right) \ln(ax+b) \quad (48)$$

$$\int x \ln(a^2-b^2x^2) dx = -\frac{1}{2} x^2 + \frac{1}{2} \left(x^2 - \frac{a^2}{b^2} \right) \ln(a^2-b^2x^2) \quad (49)$$

Integrals with Exponentials

$$\int e^{ax} dx = \frac{1}{a} e^{ax} \quad (50)$$

$$\int \sqrt{x} e^{ax} dx = \frac{1}{a} \sqrt{x} e^{ax} + \frac{i\sqrt{\pi}}{2a^{3/2}} \operatorname{erf}(i\sqrt{ax}), \text{ where } \operatorname{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt \quad (51)$$

$$\int x e^x dx = (x-1) e^x \quad (52)$$

$$\int x^a e^x dx = \left(\frac{x}{a} - \frac{1}{a^2} \right) e^{ax} \quad (53)$$

$$\int x^2 e^x dx = (x^2 - 2x + 2) e^x \quad (54)$$

$$\int x^2 e^{ax} dx = \left(\frac{x^2}{a} - \frac{2x}{a^2} + \frac{2}{a^3} \right) e^{ax} \quad (55)$$

$$\int x^3 e^x dx = (x^3 - 3x^2 + 6x - 6) e^x \quad (56)$$

$$\int x^n e^{ax} dx = \frac{x^n e^{ax}}{a} - \frac{n}{a} \int x^{n-1} e^{ax} dx \quad (57)$$

$$\int x^n e^{-ax} dx = \frac{(-1)^n}{a^{n+1}} \Gamma[1+n, -ax], \text{ where } \Gamma(a, x) = \int_x^\infty t^{a-1} e^{-t} dt \quad (58)$$

$$\int e^{ax^2} dx = \frac{i\sqrt{\pi}}{2\sqrt{a}} \operatorname{erf}(i\sqrt{ax}) \quad (59)$$

$$\int e^{-ax^2} dx = \frac{\sqrt{\pi}}{2\sqrt{a}} \operatorname{erf}(x\sqrt{a}) \quad (60)$$

$$\int x e^{-ax^2} dx = -\frac{1}{2a} e^{-ax^2} \quad (61)$$

$$\int x^2 e^{-ax^2} dx = \frac{1}{4} \sqrt{\frac{\pi}{a^3}} \operatorname{erf}(x\sqrt{a}) - \frac{x}{2a} e^{-ax^2} \quad (62)$$

Integrals with Trigonometric Functions

$$\int \sin ax dx = -\frac{1}{a} \cos ax \quad (63)$$

$$\int \sin^2 ax dx = \frac{x}{2} - \frac{\sin 2ax}{4a} \quad (64)$$

$$\int \sin^n ax dx = -\frac{1}{a} \cos ax \cdot {}_2F_1\left[\frac{1}{2}, \frac{1-n}{2}, \frac{3}{2}, \cos^2 ax\right] - \frac{1}{a} \cos ax \quad (65)$$

$$\int \sin^3 ax dx = -\frac{3 \cos ax}{4a} + \frac{\cos 3ax}{12a} \quad (66)$$

$$\int \cos ax dx = \frac{1}{a} \sin ax \quad (67)$$

$$\int \cos^2 ax dx = \frac{x}{2} + \frac{\sin 2ax}{4a} \quad (68)$$

$$\int \cos^n ax dx = -\frac{1}{a(1+p)} \cos^{1+p} ax \times {}_2F_1\left[\frac{1+p}{2}, \frac{1}{2}, \frac{3+p}{2}, \cos^2 ax\right] \quad (69)$$

$$\int \cos^3 ax dx = \frac{3 \sin ax}{4a} + \frac{\sin 3ax}{12a} \quad (70)$$

$$\int \cos ax \sin bxdx = \frac{\cos[(a-b)x]}{2(a-b)} - \frac{\cos[(a+b)x]}{2(a+b)}, a \neq b \quad (71)$$

$$\int \sin^2 ax \cos bxdx = -\frac{\sin[2(a-b)x]}{4(2a-b)} + \frac{\sin bx}{2b} - \frac{\sin[(2a+b)x]}{4(2a+b)} \quad (72)$$

$$\int \sin^2 x \cos x dx = \frac{1}{3} \sin^3 x \quad (73)$$

$$\int \cos^2 ax \sin bxdx = \frac{\cos[(2a-b)x]}{4(2a-b)} - \frac{\cos bx}{2b} - \frac{\cos[(2a+b)x]}{4(2a+b)} \quad (74)$$

$$\int \cos^2 ax \sin ax dx = -\frac{1}{3a} \cos^3 ax \quad (75)$$

$$\int \sin^2 ax \cos^2 bxdx = \frac{x}{4} - \frac{\sin 2ax}{8a} - \frac{\sin[2(a-b)x]}{16(a-b)} + \frac{\sin 2bx}{8b} - \frac{\sin[2(a+b)x]}{16(a+b)} \quad (76)$$

$$\int \sin^2 ax \cos^2 ax dx = \frac{x}{8} - \frac{\sin 4ax}{32a} \quad (77)$$

$$\int \tan ax dx = -\frac{1}{a} \ln \cos ax \quad (78)$$

$$\int \tan^2 ax dx = -x + \frac{1}{a} \tan ax \quad (79)$$

$$\int \tan^n ax dx = \frac{(-1)^n}{a^{n+1}} \Gamma[1+n, -ax], \text{ where } \Gamma(a, x) = \int_x^\infty t^{a-1} e^{-t} dt \quad (80)$$

$$\int \tan^3 ax dx = \frac{1}{a} \ln \cos ax + \frac{1}{2a} \sec^2 ax \quad (81)$$

$$\int \sec x dx = \ln |\sec x + \tan x| = 2 \tanh^{-1} \left(\tan \frac{x}{2} \right) \quad (82)$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax \quad (83)$$

$$\int \sec^3 x dx = \frac{1}{2} \sec x \tan x + \frac{1}{2} \ln |\sec x + \tan x| \quad (84)$$

$$\int \sec x \tan x dx = \sec x \quad (85)$$

$$\int \sec^2 x \tan x dx = \frac{1}{2} \sec^2 x \quad (86)$$

$$\int \sec^n x \tan x dx = \frac{1}{n} \sec^n x, n \neq 0 \quad (87)$$

$$\int \csc x dx = \ln \left| \tan \frac{x}{2} \right| = \ln |\csc x - \cot x| + C \quad (88)$$

$$\int \csc^2 ax dx = -\frac{1}{a} \cot ax \quad (89)$$

$$\int \csc^3 x dx = -\frac{1}{2} \cot x \csc x + \frac{1}{2} \ln |\csc x - \cot x| \quad (90)$$

$$\int \csc^n x \cot x dx = -\frac{1}{n} \csc^n x, n \neq 0 \quad (91)$$

$$\int \sec x \csc x dx = \ln |\tan x| \quad (92)$$

Products of Trigonometric Functions and Monomials

$$\int x \cos x dx = \cos x + x \sin x \quad (93)$$

$$\int x \cos ax dx = \frac{1}{a^2} \cos ax + \frac{x}{a} \sin ax \quad (94)$$

$$\int x^2 \cos x dx = 2x \cos x + (x^2 - 2) \sin x \quad (95)$$

$$\int x^2 \cos ax dx = \frac{2x \cos ax}{a^2} + \frac{a^2 x^2 - 2}{a^3} \sin ax \quad (96)$$

$$\int x^n \cos x dx = -\frac{1}{2} (i)^{n+1} [\Gamma(n+1, -ix) + (-1)^n \Gamma(n+1, ix)] \quad (97)$$

$$\int x^n \cos ax dx = \frac{1}{2} (ia)^{1-n} [(-1)^n \Gamma(n+1, -iax) - \Gamma(n+1, iax)] \quad (98)$$

$$\int x \sin x dx = -x \cos x + \sin x \quad (99)$$

$$\int x \sin ax dx = -\frac{x \cos ax}{a} + \frac{\sin ax}{a^2} \quad (100)$$

$$\int x^2 \sin x dx = (2-x^2) \cos x + 2x \sin x \quad (101)$$

$$\int x^2 \sin ax dx = \frac{2-a^2 x^2}{a^3} \cos ax + \frac{2x \sin ax}{a^2} \quad (102)$$

$$\int x^n \sin x dx = -\frac{1}{2} (i)^n [\Gamma(n+1, -ix) - (-1)^n \Gamma(n+1, -ix)] \quad (103)$$

Products of Trigonometric Functions and Exponentials

$$\int e^x \sin x dx = \frac{1}{2} e^x (\sin x - \cos x) \quad (104)$$

$$\int e^{bx} \sin ax dx = \frac{1}{a^2 + b^2} e^{bx} (b \sin ax - a \cos ax) \quad (105)$$

$$\int e^x \cos x dx = \frac{1}{2} e^x (\sin x + \cos x) \quad (106)$$

$$\int e^{bx} \cos ax dx = \frac{1}{a^2 + b^2} e^{bx} (a \sin ax + b \cos ax) \quad (107)$$

$$\int x e^x \sin x dx = \frac{1}{2} e^x (\cos x - x \cos x + x \sin x) \quad (108)$$

$$\int x e^x \cos x dx = \frac{1}{2} e^x (x \cos x - \sin x + x \sin x) \quad (109)$$

Integrals of Hyperbolic Functions

$$\int \cosh ax dx = \frac{1}{a} \sinh ax \quad (110)$$

$$\int e^{ax} \cosh bxdx = \begin{cases} \frac{e^{ax}}{a^2 - b^2} [a \cosh bx - b \sinh bx] & a \neq b \\ \frac{e^{2ax}}{4a} + \frac{x}{2} & a = b \end{cases} \quad (111)$$

$$\int \sinh ax dx = \frac{1}{a} \cosh ax \quad (112)$$

$$\int e^{ax} \sinh bxdx = \begin{cases} \frac{e^{ax}}{a^2 - b^2} [-b \cosh bx + a \sinh bx] & a \neq b \\ \frac{e^{2ax}}{4a} - \frac{x}{2} & a = b \end{cases} \quad (113)$$

$$\int e^{ax} \tanh bxdx = \begin{cases} \frac{e^{(a+2b)x}}{(a+2b)^2} {}_2F_1\left[1 + \frac{a}{2b}, 1 + \frac{a}{2b}, -e^{2bx}\right] & a \neq b \\ \frac{e^{ax} - 2 \tanh^{-1}[e^{ax}]}{a} & a = b \end{cases} \quad (114)$$

$$\int \tanh ax dx = \frac{1}{a} \ln \cosh ax \quad (115)$$

$$\int \cos ax \cosh bxdx = \frac{1}{a^2 + b^2} [a \sin ax \cosh bx + b \cos ax \sinh bx] \quad (116)$$

$$\int \cos ax \sinh bxdx = \frac{1}{a^2 + b^2} [b \cos ax \cosh bx + a \sin ax \sinh bx] \quad (117)$$

$$\int \sin ax \cosh bxdx = \frac{1}{a^2 + b^2} [-a \cos ax \cosh bx + b \sin ax \sinh bx] \quad (118)$$

$$\int \sin ax \sinh bxdx = \frac{1}{a^2 + b^2} [b \cosh bx \sin ax - a \cos ax \sinh bx] \quad (119)$$

$$\int \sinh ax \cosh bxdx = \frac{1}{4a} [-2ax + \sinh 2ax] \quad (120)$$

$$\int \sinh ax \cosh bxdx = \frac{1}{b^2 - a^2} [b \cosh bx \sinh ax - a \cosh ax \sinh bx] \quad (121)$$

*© 2014. From <http://integral-table.com>, last revised June 14, 2014. This material is provided as is without warranty or representation about the accuracy, correctness or suitability of the material for any purpose, and is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 United States License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/> or send a letter to Creative Commons, 171 Second Street, Suite 300, San Francisco, California, 94105, USA.