


PHYSICS 105 - Principles of Physics

Kinematics 1D, Forces & vectors, kinematics 2D, Newton's law
Conservation momentum & collision, Energy & work
Optional: rotational kinematics

Math review: proportion, Sci. Notations, trig., vectors


LAB lasts 2:15
Don't expect less !

If it bites or scratches,
it's biology.
If it stinks or pops,
it's chemistry.
If it doesn't work,
it's physics







Demo: free-fall
EM /



Michael
Faraday





James Prescott
Joule



Richard Feynman

<https://youtu.be/luv6hY6zsd0?t=281>



Hw: <https://www.youtube.com/watch?v=0NbBjNiw4tk> Michio Kaku -watch. Quiz next week.

UNITS we will cover in this class:

- review Math :Proportions, scientific notation
- Notions of motion – 1D kinematics, graphical analysis of motion free-fall.
- , trigonometry, vectors, components of vectors, adding vectors.
- Forces, equilibrium, torques.
- Kinematics 2D, projectile motion, circular motion.
- Newton's laws, impulse, momentum.
- Conservation of momentum, collision.
- Work and energy, conservation of energy, energy of spin.

If time permits: rotational kinematics

Physics experiments involve the measurement of a variety of physical quantities.

These measurements should be accurate and reproducible.

The first step in ensuring accuracy and reproducible is defining the **units** in which the measurements are made.

Units and derived units

SI Base Units

Base Quantity	Name	Symbol
Length	meter	m
Mass	kilogram	kg
Time	second	s
Electric current	ampere	A
Temperature	kelvin	K
Amount of substance	mole	mol
Luminous intensity	candela	cd

Important:

Speed is in m/s
and m/s is noted m s^{-1}

Acceleration is in m/s/s
And is also noted m s^{-2}

Distance (displacement) is in m

SI Derived Units

Derived Quantity	Name	Symbol	Equivalent SI units
Frequency	hertz	Hz	s^{-1}
Force	newton	N	$\text{m} \cdot \text{kg} \cdot \text{s}^{-2}$
Pressure	pascal	Pa	N/m^2
Energy	joule	J	$\text{N} \cdot \text{m}$
Power	watt	W	J/s
Electric charge	coulomb	C	$\text{s} \cdot \text{A}$
Electric potential	volt	V	W/A
Electric resistance	ohm	Ω	V/A
Celsius temperature	degree Celsius	$^{\circ}\text{C}$	K°

Important:

Units can give away equations.

Force is $\text{kg m/s/s} \rightarrow$

Find the equation for force

find the equation for power

find the equation for pressure

Find the equation for energy

Step 0: MATH REVIEW

- Solving proportion

See assignment_conversion_units partially in class then assignment

- scientific notation


See assignment_sci_notin in class

- Read chapter 1 from your book:

Big numbers – little numbers

Arithmetic in Sci Not

Dimensionality



And do quiz2_chapter1_EZphysics_home as take home quiz

- trigonometry: will be covered in lab 3

Following is a list of prefixes and their meanings that are often used in conjunction with SI units:

Multiple	Prefix	Symbol
10^{12}	tera	T
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n

$$\left\{ \begin{array}{l} \text{kilo (k)} = 10^3 \\ 1 \text{ km} = 10^3 \text{ m} \\ 1 \text{ kg} = 10^3 \text{ g} \end{array} \right.$$

$$\left\{ \begin{array}{l} \text{milli (m)} = 10^{-3} \\ 5 \text{ ml} = 5 \times 10^{-3} \text{ l} \\ 5 \text{ mm} = 5 \times 10^{-3} \text{ m} \\ 5 \text{ mg} = 5 \times 10^{-3} \text{ g} \end{array} \right.$$

$$\left\{ \begin{array}{l} \text{nano (n)} = 10^{-9} \\ 7 \text{ ns} = 7 \times 10^{-9} \text{ sec} \\ 7 \text{ nm} = 7 \times 10^{-9} \text{ m} \end{array} \right.$$

$$\left\{ \begin{array}{l} \text{tera (T)} = 10^{12} \\ 5 \text{ TJ} = 5 \times 10^{12} \text{ joules} \end{array} \right.$$

$$\left\{ \begin{array}{l} \text{mega (M)} = 10^6 \\ 25 \text{ MB} = 25 \times 10^6 \text{ B} \\ \text{B is for byte} \end{array} \right. \left\{ \begin{array}{l} \text{centi (c)} = 10^{-2} \\ 5 \text{ cm} = 5 \times 10^{-2} \text{ m} \\ 5 \text{ cg} = 5 \times 10^{-2} \text{ g} \end{array} \right.$$

$$\left\{ \begin{array}{l} \text{micro (\mu)} = 10^{-6} \\ 3 \text{ }\mu\text{m} = 3 \times 10^{-6} \text{ m} \\ 3 \text{ }\mu\text{g} = 3 \times 10^{-6} \text{ g} \end{array} \right.$$

Remember :

$$1\text{km} = 1,000\text{m} (10^3 \text{ meters})$$

$$1\text{m} = 100\text{cm} (10^2 \text{ centimeters})$$

$$1\text{cm} = 10\text{mm} (10 \text{ millimeters})$$

$$1\text{mm} = 1000 \mu\text{m} (10^3 \text{ micrometers})$$

$$1\text{m} = 1000 \text{ mm} (10^3 \text{ mm})$$

$$1\text{m} = 1,000,000 \mu\text{m} (10^6 \mu\text{m})$$

note that: if $1\text{m} = 10^6 \mu\text{m}$ then $1\mu\text{m} = 10^{-6} \text{ m}$

$$1\text{mm} = 10^{-3} \text{ m} \quad 1\text{cm} = 10^{-2} \text{ m}$$

$$1 \text{ mile} = 1,609\text{m}$$

$$10 \text{ N} = 2.2 \text{ pounds}$$

$$1\text{min} = 60\text{s}$$

$$1\text{hour} = 60\text{min} = 3600\text{s}$$

$$1 \text{ day} = 24 \text{ h} = 24 \times 3600 \text{ s} = 86,400\text{s}$$

$$1\text{kg} = 1000 \text{ g}$$

$$1\text{g} = 1,000\text{mg}$$

Following is a list of prefixes and their meanings that are often used in conjunction with SI units:

Multiple	Prefix	Symbol
10^{12}	tera	T
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n

$$1\text{m/s} = 2.2 \text{ mph}$$

On Earth 1 kg is 2.2 pounds

Note: kg is a mass

Pounds measures a weight.

Unit 1: kinematics 1D

- 1. Average speed - slide 12**
- 2. displacement (change in position) vs distance - slide 15**
- 3. velocity vs speed - slide 19**
- 4. acceleration - slide 20**
- 5. kinematics equations – slide 35**
- 6. graphical analysis of motion - slide 45**
- 7. Free-fall – see unit 1B**

Kinematics deals with the concepts that are needed to describe motion.

Dynamics deals with the effect that forces have on motion.

Together, kinematics and dynamics form the branch of physics known as ***Mechanics***.

Kinematics is the portion of mechanics that describes motion without any reference to ?

- a) forces
- b) accelerations
- c) velocities
- d) displacements
- e) time

Distance - Average speed

Distance (d) measures “how far”. Unit is m.

time (t) is the time elapsed. Unit is second.

Average speed (s) is the distance traveled divided by the time required to cover the distance. Unit is m/s.

$$\text{Average speed} = \frac{\text{total distance}}{\text{elapsed time}}$$

SI units for speed: **meters per second** (m/s)

Example:

A NASCAR car travels at an average speed of 190 miles/hour.

A) How long would a 500 mile race take in hours?

B) Convert 190mph to m/s

(remember $1\text{m/s} = 2.2\text{ mph}$)

C) in 5 hours how many miles were covered ?

Convert to km



$$\text{Average speed} = \frac{\text{Distance}}{\text{Elapsed time}}$$


Examples

- 1) How far does a jogger run in 1.5 hours (convert 1st in s) if his average speed is 2.22 m/s? Convert to km
Convert to miles (1 mile = 1,609m)
- 2) If it takes you 45 seconds to run 340m, what is your average speed?
- 3) How long does it take a skier to travel 650m, going 18.5m/s ?
- 4) at 9:00AM you start on a walk. At 10:30AM you are 4.6 miles from Your starting point. what is your average speed in mph ?
Convert to m/s (Sorry units used in Physics are meters, Seconds, kilograms, joule ..) 1 mile = 1609m
- 5) What is the average speed of a runner who finishes the 1,500 meters race in 3 minutes 30 seconds ? In m/s

Your speedometer does not indicate averages; it tells you how fast you are going now. This is your instantaneous speed. If the instantaneous does not change, you are going at a constant speed. The constant value of the speed will be equal to your average speed for the whole trip.

In Physics displacement and distance are not the same physical quantities.

- 1) If you walk from A to B following the red path,
The distance d is the length of the path.
You can measure it using a surveyor wheel:



- 2) If you connect A (start) to B (end) using an arrow
This represent the displacement. \overrightarrow{AB}

In Physics/Math
displacement AB is called a vector

A displacement has a length (length of arrow)
and a direction (like 30NW) .

BUT do not worry (for now) in 1D displacement is easy to understand.
Displacement tells if the motion is @ left /@ right or @ up/@down

Coordinate System

1. Origin

2. Axis

positive coordinate direction

3. Unit Vector



0


Kinematics:

Position as a function of time $x(t)$

Velocity as a function of time $V(t)$

Acceleration as a function of time a



Position x: location on number line



Finding the displacement
In 1D is easy =
Change in position on number line

displacement Δx : change in position in 1D
Connect initial position to final position

$$\Delta x = x_2 - x_1 = -5 - 3 = -8 \text{ units}$$




If $X_1 = -5$ and displacement = 7 what is the final position X_2 ?

Same principle can apply to the vertical scale.
If $Y_1 = -3$ and the $Y_2 = 2$ what is the displacement

If the Y_1 is -10 and the displacement is 5 what is the final position Y_2 ?

We work in one dimension for now along the number line.



In one dimension the displacement is just the change of position on the number line.


Displacement = final position – initial position

Displacement is noted Δx

$\Delta x = x_2 - x_1 = x_{\text{final}} - x_{\text{initial}}$

Here the displacement \overrightarrow{AB} is $x_2 - x_1 = -6 - 3 = -9$ units.
Motion @ left \Rightarrow the displacement is negative
Motion @ right \Rightarrow the displacement is positive

The distance is always positive. Here the distance is 9 units.
Displacement in physics is a physical quantity called vector.
It has a magnitude, a direction and a unit.




$$AB = \Delta x = ?$$

$$CD = \Delta x = ?$$

If you move from A to B then B to A :

- what is the displacement ? (connect starting point to ending point or $X_{\text{final}} - X_{\text{initial}}$)
- what is the distance ?

You can add displacements
 $+5 + 3 = +8$



**Connect the starting point
To the final point**

What is the sum of the displacements ?
 What is the notation ?

A physical quantity that has a direction and a magnitude like a displacement is called a vector. It is represented by an arrow.

$$\begin{array}{c} \text{5} \\ \longrightarrow \end{array} + \begin{array}{c} \text{5} \\ \longrightarrow \end{array} = \begin{array}{c} \text{10} \\ \longrightarrow \end{array}$$

It is easy to add vectors in 1D. Complete the page.
Draw the vector sum and write its notation.

$$\begin{array}{c} \text{5} \\ \longrightarrow \end{array} + \begin{array}{c} \text{-5} \\ \longleftarrow \end{array}$$

$$\begin{array}{c} \text{5} \\ \longrightarrow \end{array} + \begin{array}{c} \text{10} \\ \longrightarrow \end{array} =$$

You walk 5 yards @ right and another 5 yards @ right .
The net sum is **10 yards @ right**

$$\begin{array}{c} \text{5} \\ \longrightarrow \end{array} + \begin{array}{c} \text{-10} \\ \longleftarrow \end{array} =$$

10 yards @ left is simply -10 yards.
Since there are only 2 directions. Minus is for left.

$$\begin{array}{c} \text{5} \\ \longrightarrow \end{array} + \begin{array}{c} \text{-15} \\ \longleftarrow \end{array} =$$

Up is positive
Down is negative.

$$\begin{array}{c} \text{10} \\ \uparrow \end{array} + \begin{array}{c} \text{-5} \\ \downarrow \end{array} =$$

$$\begin{array}{c} \uparrow \\ 1500 \end{array} + \begin{array}{c} \downarrow \\ 2000 \end{array}$$

$$\begin{array}{c} \uparrow \\ 1500 \end{array} + \begin{array}{c} \downarrow \\ 1500 \end{array}$$

$$\begin{array}{c} \downarrow \\ 4000 \end{array} + \begin{array}{c} \uparrow \\ 500 \end{array}$$

In Physics Velocity and speed are not the same physical quantity:

average velocity = displacement / time : $V = \Delta x / \text{time}$

If the object is moving @right, velocity is positive

If the object is moving @left, velocity is negative

So to find the velocity : find the displacement (see previous slides) and divide by the time !

velocity is called a vector – notation is \vec{V}

It has a magnitude, a direction (which way: left or right if 1D motion), a unit (m/s)


(other examples of vectors: force, acceleration, displacement)

average speed = distance / time = d/t (see previous slides)

speed is not a vector (it is called a scalar) – it has a magnitude, a unit (m/s)

(other examples of scalars: time, energy, distance, temperature, volume ...

any quantity without a direction)



1) A race car covers 1700m @ right in 5s. What is the average velocity?
The same race drive in the opposite direction with the wind opposing
Its motion. It covers 1700m @ left in 5.4 s. What is the average velocity ?

2) In NYC you walk 10 blocks @ south (or -10) in 30minutes. What is your average velocity in blocks/hour

In class

3) you drive 100 miles @ east (so +10) in 2 hours.
What is your average velocity in mph and m/s

4) If you walk 5 blocks @ north and you realize this is the wrong way.
You then walk 10 blocks @ south. The all trip lasted 30 minutes.
What is the average velocity (from start to end) blocks/hour
What is the average speed. (not the same)

In class


5) same question for 10 blocks @ west (-10) and 20 blocks @ east.
The trip lasts 45 minutes

6) same question for 100 miles @ North and 50 miles @ south in 2 hours.
In mph in m/s

7) An insect flies 5 inches @ up 3 inches down then 15 inches @ up in 2 s.
What is its average velocity ? In in/s Average speed?

HW for next class

First step in kinematics: the number line or Coordinate system.



DEFINITION OF AVERAGE ACCELERATION

(how fast an object is getting faster)

a = change of velocity / time elapsed = (Vfinal – Vinitial) / time
acceleration is the change of velocity per unit time

Unit is (m/s)/s or m/s/s or m/s²

Remember: acceleration has the same direction as the force applied to the object!
Because a and force are proportional,

Example: acceleration due to gravity is about 22mph/s

Which means that every second a falling object goes 22mph faster every second.

A falling object (dropped from a tall tower) reaches a speed of 88mph after ___ seconds.

The acceleration is @ down because the _____ of the object is pulling it down.

23

22mph per second is _____ m/s per second.

After 3 seconds of fall, the speed reached is _____ m/s


ACCELERATION is a vector.


It has a magnitude (change in velocity per unit time) and a direction

If there is a thrust (or push/pull) @ right, acceleration is positive

If there is a thrust @ left, acceleration is negative.

The notion of *acceleration* emerges when a change in velocity is combined with the time during which the change occurs.





Example Acceleration and Increasing Velocity

Determine the average acceleration of the plane in km/h/s

$V_{\text{initial}} = 0$

$V_{\text{final}} = 250 \text{ km/h}$

time elapsed = 30s

A) Acceleration in km/h/s (see next slide before going to B)


B) How fast it is going after 5 seconds (find speed in km/h)

C) for the same acceleration what is the speed after 5s if the initial speed is 50km/h ?

D) convert the speed found in B to m/s (1km=1000m and 1 hour=3600s)

$$\vec{a} = \frac{+9.0 \text{ km/h}}{\text{s}}$$

Can you find the rate of Increase?



Acceleration is a vector:
9 is the magnitude ; km/h/s is the unit ; and + is the direction

Acceleration and Decreasing Velocity

Initial velocity = 28 m/s

Final velocity = 13 m/s

Between $t=9\text{ s}$ and $t=12\text{ s}$

Find the acceleration

1. 5 m/s/s


2. 1.7 m/s/s

3. -5 m/s/s

The acceleration has the same direction
than the force (push/pull)

Here the pull is @ left so acceleration is @ ____
but the velocity is @ _____

Velocity is a vector too. It has a _____
And a _____.





$t_0 = 9.0\text{ s}$




$t = 12\text{ s}$

$\vec{a} =$




(b)



If velocity and acceleration have the same direction \Rightarrow speeding up
If velocity and acceleration have opposite direction \Rightarrow slowing down

http://www.walter-fendt.de/html5/phen/acceleration_en.htm

 Use $V_0=5$ and $a = 1$
Then $V_0=10$ and $a = -1$ 28

Try also $V_0=0$ $a = -1$ $X_0=50$
Similar to lab2 part 4

Consider a motion with:
 Initial position $X_0 = 0$
 Initial $V_0 = 5\text{m/s}$
 Acceleration $a = -1\text{m/s/s}$

Take home assignment

B) fill the column 2. Remember $V = V_0 + a t$ (so for example if $t = 6$ then $V = 5 + (-1) \cdot 6 = -1\text{m/s}$)
 Column 1 is the time elapsed Column 2 is the velocity

time	velocity	Position X
0		
1		
2		
3		
4		
5		
6		
7		
8		

B) the position X of an object moving with a constant acceleration a and initial velocity 5m/s is given by:

$$X = V_0 \times \text{time} + 0.5 \times a \times \text{time}^2$$

$$\text{So } X = 5t + 0.5(-1)t^2$$

Use the equation to fill column 3

(example if $t = 6$ $x = 5 \cdot 6 + 0.5 \cdot (-1) \cdot (36) = 12$. note: $6^2 = 36$)

C) Make 2 graphs. One graph for:
 Velocity vs time
 Position vs time

D) trace (insert) the best fit line for graph 1
 Display equation on graph.

E) trace (insert) the best fit polynomial degree 2 (parabola)
 Display equation on graph

Starting from rest, a particle confined to move along a straight line is accelerated at a rate of 2 m/s^2 .

- a) The particle travels 2 m during each second.
- b) The particle travels 2 m only during the first second.
- c) The speed of the particle increases by 2 m/s during each second.
- d) The acceleration of the particle increases by 2 m/s^2 during each second.

The drawing shows the position of a rolling ball at one second intervals. Which one of the following phrases best describes the motion of this ball?



- a) constant position
- b) constant velocity
- c) increasing velocity
- d) constant acceleration
- e) decreasing velocity

In which one of the following situations does the car have an acceleration that is directed due north?

- a) A car travels northward with a constant speed of 24 m/s.
- b) A car is traveling southward as its speed increases from 24 m/s to 33 m/s.
- c) A car is traveling southward as its speed decreases from 24 m/s to 18 m/s.
- d) A car is traveling northward as its speed decreases from 24 m/s to 18 m/s.
- e) A car travels southward with a constant speed of 24 m/s.

A postal truck driver driving due east gently steps on her brake as she approaches an intersection to reduce the speed of the truck. What is the direction of the truck's acceleration, if any?

- a) There is no acceleration in this situation.
- b) due north
- c) due east
- d) due south
- e) due west

A sports car starts from rest. After 10.0 s, the speed of the car is 25.0 m/s. What is the magnitude of the car's acceleration?

- a) 2.50 m/s^2
- b) 5.00 m/s^2
- c) 10.0 m/s^2
- d) 25.0 m/s^2
- e) 250 m/s^2

Which one of the following is not a vector quantity?

- a) acceleration
- b) displacement
- c) instantaneous velocity
- d) average velocity
- e) average speed

Which one of the following equations is the correct expression for average acceleration?

a) $a = \frac{v}{t}$

b) $a = \frac{dv}{dt}$

c) $a = \frac{\Delta x}{\Delta t}$

d) $a = \frac{\Delta v}{\Delta t}$

e) $a = \frac{1}{2}vt^2$

1) What is the acceleration of a rocket ship in outer space that takes 5.0s to increase its speed from 1240m/s to 1300m/s ?

(textbook)

2) In outer space a rocket ship is traveling at the enormous speed of 2800m/s. What is its acceleration if it increases its

Speed uniformly and is going 2840m/s after 25 s ?

Convert to mph (do it in class)

3) A car travels with an average speed of 25m/s (do it in class)

A) What is the speed in km/s (1km =1000m)

B) What is the speed in km/h (1m/s = 3.6 km/h)

4) A car travels with an average speed of 58mph. What is the speed in km/h

(1.6km = 1 mile)

5) Starting from rest and moving in a straight line, a runner achieves a velocity of 7m/s in a time of 2s. What is the

Average acceleration of the runner ?

6) starting from rest, a car accelerates at a rate of 4.2m/s/s for a time of 5 seconds.

What is its velocity at the end of this time ?

7) The velocity of a car decreases from 30m/s to 18m/s in a time of 4seconds. What is the average acceleration if the car

In this process ?

Equations of kinematics: the big 5

Notations:

V (V_{final}) final velocity

V_i (V_{initial}) initial velocity \rightarrow velocity at $t=0$

X_i (X_{initial}) initial position \rightarrow position at $t=0$

X (X_{final}) final position

Δx is the displacement ($x-x_i$) how far you are from "home"

a acceleration

t is the time elapsed

V_i is the initial velocity and is
Sometimes noted V_0

X_i is the initial position (home) and
Is sometimes notes x_0

EQUATIONS OF KINEMATICS

for motion with constant acceleration


1) $x = x_i + V_i t + 0.5 a t^2$ or $\Delta x = V_i t + 0.5 a t^2$ if $x_i=0$ then $x = V_i t + 0.5 a t^2$

2) $V = V_i + at$

3) $a = (V - V_i)/t$

4) $V^2 = V_i^2 + 2a(x - x_i)$ or $V^2 = V_i^2 + 2a(\Delta x)$

5) Average velocity $= \Delta x/t$ or $(V + V_i)/2$



EQUATIONS OF KINEMATICS

for motion with constant velocity and acceleration = 0


1) $x = x_i + V_i t$ or $\Delta x = V_i t$ if $x_i=0$ then $x = V_i t$

2) $V_i = V_f$ stays the same

3) $a = 0$

4) NA

5) Average velocity $= \Delta x/t$



**Let's use the 2 equations : $\Delta x = V_i t + 0.5 (a) t^2$ and $V = V_i + at$
or $x = x_i + V_i t + 0.5 (a) t^2$**

1) A car accelerating with an initial velocity of 12m/s accelerates at a constant rate of 2.5m/s/s for a time of 2s.

A) What is its velocity at the end of this time ?

B) What displacement does the car cover during this process (the initial position $X_0 = 0$ find X)

2) A car moving with an initial velocity of 30m/s slows down at a constant rate of -3m/s/s

A) What is the velocity after 3 seconds of deceleration

B) What displacement does the car cover in this time: (Find X with $X_0 = 0$)

3) A runner moving with an initial velocity of 9m/s slows down at a constant rate of -1.5m/s/s over a period of 2 seconds.

A) What is her velocity at the end of this time

B) What displacement did she cover during this process(X with $X_0 = 0$)

4) Starting from rest ($V_{\text{initial}} = 0$) , a car accelerated at a constant rate of 3m/s/s for a time of 5 seconds. $X_i = 0$

A) Compute the velocity of the car at 1s, 2s, 3s, 4s and 5s and plot the velocity versus time.,

B) What is the equation of this line ? (V as a function of time)

C) compute the displacement (so X) traveled for these time and plot the distance against time

t	0	1	2	3	4	5
V						

t	0	1	2	3	4	5
X						

39

D) What is the equation of the position X vs time

5) if $V_0 = 3\text{m/s}$ and $V_{\text{final}} = 9\text{m/s}$ What is the average velocity if the acceleration is constant.

6) If the initial velocity of a car is 30m/s and its acceleration is 10m/s/s . Initial position is 0.

A) what its equation of motion $x=f(t)$.

B) what is the equation of $V=f(t)$ (velocity as a function of time)

7) if the equation of motion of an object is $x = 5 + 10t + 2t^2$ What are:
Initial position ? initial velocity ? Acceleration ?

8) same question for $x = 10t + t^2$

9) $x = 10t^2$

10) if $x = 5 + 10t + 10t^2$


What is the velocity at $t=5s$ (after 5s)

11) if $V = 10 + 2t$

What is the acceleration ?

What is the final velocity when $t=100s$ (so after 100s)?

What is the average speed during that time ?



FIND THE displacement of the boat :

1. 64m

2. 72m

3. 76m


4. 112m

$$[1] \quad x = v_0 t + \frac{1}{2} a t^2$$

$$[2] \quad v = v_0 + a t$$

$$[3] \quad a = \text{constant}$$

$$[4] \quad v^2 = v_0^2 + 2a(x)$$



FIND THE displacement of the airplane :

1. 62m
2. 1m
3. 124m

x	a	v	v_o	t

$$[1] \quad x = v_0 t + \frac{1}{2} a t^2$$

$$[2] \quad v = v_0 + a t$$

$$[3] \quad a = \text{constant}$$

$$[4] \quad v^2 = v_0^2 + 2a(x)$$

2.4.3. In which one of the following situations is the displacement of the ball directly proportional to the elapsed time?

- a) a ball rolls with constant velocity
- b) a ball at rest is given a constant acceleration
- c) a ball rolling with velocity v_0 is given a constant acceleration
- d) a ball rolling uphill experiences a decreasing acceleration
- e) a ball rolling downhill experiences an increasing acceleration

Example 8 An Accelerating Spacecraft

A spacecraft is traveling with a velocity of +3250 m/s. Suddenly the retrorockets are fired, and the spacecraft begins to slow down with an acceleration whose magnitude is 10.0 m/s². What is the velocity of the spacecraft when the displacement of the craft is +215 km, relative to the point where the retrorockets began firing?

x	a	v	v_o	t

2 possible answers.

1. +/- 6262500 m/s
2. +/- 2502.5 m/s
3. +/- 3855 m/s


$$[1] \quad x = v_o t + \frac{1}{2} a t^2$$

$$[2] \quad v = v_o + a t$$


$$[3] \quad a = \text{constant}$$

$$[4] \quad v^2 = v_o^2 + 2a(x)$$

Why 2 possible answers ?



Similar situation:



Throwing a ball straight up.

The weight (force) is always pulling @ down. So acceleration is always @ down (negative)

On the way up the force causes The ball to slow down.

On the way down the force causes The ball to speed up.

Try to sketch the graphs for:
Acceleration / velocity/ position
For the way up and for the way down.

1) How far does a car travel if it starts at rest and accelerated at 3m/s/s for **6.5s** (HW)

2) How long does it take a car, starting from rest, to travel 240m if its acceleration is 1.90m/s/s ? (in class)

3) How far a car travel if it starts at rest and accelerated at 3m/s/s until it reaches a speed of 22m/s ? (in class)

4) A car moving 22m/s has its brakes jammed on and leaves skid marks 45m long. What was its acceleration ?
(negative) (in class)

5) What is the acceleration of a rocket-driven sled that travels 360m in 5s, starting from rest and accelerating uniformly ? (HW)

HW

6) How fast is a car going if it starts at rest and accelerates uniformly at 2.8m/s/s while traveling 220m ? (HW)

8) A car traveling in a straight line with an initial velocity of 14m/s accelerates at 2m/s/s to a velocity of 24m/s .

A) How much time does it take for the car to reach the velocity of 24m/s ?

B) What is the distance covered by the car in this process ?

(HW)

10) A runner moving at an initial velocity of 9m/s slows down at a constant rate of -1.5m/s/s over a period of 2 seconds.

A) What is her velocity at the end of this time ?

B) What distance does she travel during this process ?

(in class)

11) A car moving with an initial velocity of 30m/s slows down at a rate of -3m/s/s .

A) What is the velocity after 3 seconds of deceleration


B) What distance does the car cover during that time?/

(HW)

GRAPHICAL ANALYSIS OF MOTION

EQUATIONS OF KINEMATICS

for motion with velocity and acceleration = 0



1) $x = x_i + V_i t$ (or Δx)




Position vs time $x(t)$ is a straight line. V_i is the slope. X_i the y-intercept


2) $V = V_i$ stays the same

Velocity vs time $V(t) = V_o$ is a horizontal line

3) $a = 0$

Acceleration is 0









$$\text{Slope} = \frac{\Delta x}{\Delta t} = ?$$

What does the slope of the graph position vs time represent?


What is the unit ? What is the slope

What is the the equation of motion $x(t)$

2.7 Graphical Analysis of Velocity and Acceleration



Describe the motion, find the velocities for parts 1, 2 and 3, don't forget the directions.



A) from graph 3 what is the acceleration ?

B) from graph 2 what is the velocity ?

C) from graph 1 what is the initial position X_i ?

What is the slope ? What does the slope represents ?

What is the equation of motion $x(t)$?

Hint: remember the equation of a line is $y = mx + b$ with m is the slope and b the y-intercept

D) fill the table (use the equation from previous question)

E) from the table : the change of position per second: (that is the rise if run = 1s)

Stays the same ? Increases ?


F) copy the table in a spreadsheet. Make a scatter plot. Fit the best fit line.

Display equation. What is the equation ? So ?

(remember to label the axis with the right units. Also write a title).


t	x
0	
1	
2	
3	
4	
5	

<http://www.walter-fendt.de/ph14e/acceleration.htm>



- A) from graph 3 what is the acceleration ?
 B) from graph 2 what is the velocity ?
 Why is the velocity negative (what do you think it means ?)
 C) from graph 1 what is the initial position x_i ?
 What is the slope ? What does the slope represents ?
 What is the equation of motion $x(t)$?
 D) fill the table
 E) from the table : the change of position per second:
 Stays the same ? Increases ?
 F) copy the table in a spreadsheet. Make a scatter plot. Fit the best fit line.
 Display equation. What is the equation ? (with physical quantities)
 don't forget to label the axis with units and the title.

t	x
0	
1	
2	
3	
4	
5	



Complete the following statement: For an object moving at constant velocity, the distance traveled

- a) increases for each second that the object moves.
- b) is the same regardless of the time that the object moves.
- c) is the same for each second that the object moves.
- d) cannot be determined, even if the elapsed time is known.
- e) decreases for each second that the object moves.

Consider the motion of this car and the graph of its position vs time.

What is happening at $t = 16\text{s}$?

Describe the direction of motion for each part.


What is the velocity during the first part of motion? (you can use table)

What is the velocity during the second part of the motion? (Use table)

What is the initial position of the car ?

What is the equation of motion for the first part of motion $x(t)$?

t	x
0	0
1	-5
2	-10
3	-15
4	-20
5	-25
6	-30
7	-35
8	-40
9	-45
10	-50
11	-55
12	-60
13	-65
14	-70
15	-75
16	-80
17	-76
18	-72
19	-68
20	-64
21	-60



Sketch the graph velocity vs time

EQUATIONS OF KINEMATICS for motion with constant acceleration

1) $x = x_0 + v_0 t + 0.5 a t^2$





The graph $x(t)$ or position vs time is a parabola. The y-intercept is v_0 .
The slope increases = acceleration. The displacement increases every second.



2) $v = v_0 + a t$

The graph is a line. The slope is the acceleration a . The y-intercept is v_0

3) $a = \text{constant}$

The graph is a horizontal line. A is constant.





A) from graph 3) what is the acceleration $a =$ _____.

B) from graph 2) what is the initial velocity $V_0 =$ ____

What is the equation $V(t)$? (use A) and V_0)

C) From graph 1) what is the initial position X_0

What is the equation of motion $x(t)$?

D) fill the tables

E) copy and paste the table time (t) / position (x) in a spreadsheet.

Make a scatter plot. Fit a polynomial degree 2 (quadratic equation) and display equation

F) Copy and paste the table time(t)/ velocity(t) in a spreadsheet.


Make a scatter plot. Fit a line and display equations.

t	x
0	
1	
2	
3	
4	
5	
6	

t	V
0	
1	
2	
3	
4	
5	
6	

Exercise: consider the following object accelerating at a constant rate.


- 1) fill the tables
- 2) what is the acceleration of the object? (rate at which V increases – per second)
- 2) what is the initial position X_0 ? The initial velocity V_0 ?
- 3) what are the equations of motion $V(t)$ and $X(t)$?
- 4) use the tables to graph $X(t)$ and $V(t)$ (scatter plots)
- 5) fit the best fit line to $V(t)$. What is its equation ? So ?
What does the slope represent ?
- 6) fit a polynomial degree 2 to $x(t)$. What is the equation ? So ?




time	pos
0	0

time	vel
0	0

The slope of $x(t)$ is the velocity $V(t)$ at a given time t . It keeps increasing at a constant rate a .



Describe the motion
And find the instantaneous velocity




Equation of motion $X(t)$
And $V(t)$ for 2 graphs

This is a graph position vs time
What is the equation of the motion
(use physical quantities x and t)

What is the average velocity between
 $t=2$ and $t=5s$? (use equ)






What is the position at $t=3s$? (equ)


Velocity is the slope of the graph
What is V_0 ? (V at $t=0$ or slope
At $t=0$)



Use the equation to find the acceleration
And V_0

- . Which of the following velocity vs. time graphs represents an object with a negative constant acceleration?





Here are the positions of 2 trains.

1) describe the motion of train1 and train2
(constant vel ? Speeding up ? Slowing down ?)

2) What happens at t_B ?


3) Can you see when the 2 trains have the same velocity ?

- . Which one of the following quantities can be determined from the slope of a position versus time graph for an object in motion?
- a) position
 - b) velocity
 - c) acceleration
 - d) distance traveled
 - e) displacement

Which one of the following quantities can be determined from the slope of a velocity versus time graph for an object in motion?

- a) position
- b) velocity
- c) acceleration
- d) distance traveled
- e) displacement

A train car moves along a long straight track. The graph shows the position as a function of time for this train. The graph shows that the train:



1. speeds up
2. slows down all the time.
3. speeds up part of the time and slows down part of the time.
4. moves at a constant velocity.

Consider an object moving @ left (or down) and speeding up/

A) from graph 3) what is the acceleration ? $a = \underline{\hspace{2cm}}$. So There is a pull@ $\underline{\hspace{2cm}}$

B) from graph 2) what is the initial velocity ? $V_0 = \underline{\hspace{2cm}}$. Is the velocity negative or positive?

So the object is moving @ $\underline{\hspace{2cm}}$. What is the equation $V(t) =$




C) from graph 1) what is the initial position X_0 .

What is the equation of motion ?

D) Fill the tables below (using the derived equation)

E) copy and paste the table time (t) / position (x) in a spreadsheet. Make a scatter plot. Fit a polynomial Degree 2 (quadratic equation) and display equation

F) Copy and paste the table time(t)/ velocity(t) in a spreadsheet. Make a scatter plot. Fit a line and display equations.





t	x
0	
1	
2	
3	
4	
5	

t	V
0	
1	
2	
3	
4	
5	
6	

To do: virtual lab assignment slowing down

Consider the motion of an object thrown in the air with an initial velocity V_0 . It moves @ up but slows down, stops and changes direction and moves @ down but speeds up. $y(t)$ is the position of the object.
(This is not happening on Earth. But on another viable planet)



- A) from graph 3) what is the acceleration ? $a = \underline{\hspace{2cm}}$. So There is a pull @ $\underline{\hspace{2cm}}$
 B) from graph 2) what is the initial velocity ? $V_0 = \underline{\hspace{2cm}}$. Is the velocity negative or positive?
 So the object is moving @ $\underline{\hspace{2cm}}$. What is the equation $V(t) = \underline{\hspace{2cm}}$
 C) from graph 1) what is the initial position X_0 .
 What is the equation of motion ?
 D) Fill the tables below (using the derived equation)
 E) copy and paste the table time (t) / position (x) in a spreadsheet. Make a scatter plot.
 Fit a polynomial degree 2 (quadratic equation) and display equation
 F) Copy and paste the table time(t) / velocity(t) in a spreadsheet.
 Make a scatter plot. Fit a line and display equations

		t	x
t	V	0	
		1	
1		2	
2		3	
3		4	
4		5	
5		6	
6		7	
7		8	
8		9	