

# Cardano's Formula for Cubic Equations

Love Folami

January 15, 2022

## Abstract

[1]Girolamo Cardano, also known as Cardano, was an Italian doctor and mathematician best known for his work *Ars Magna*, the first Latin treatise on algebra. He described the strategies he had learned from Tartaglia for solving cubic and quartic problems.

## 1 Introduction to Cardano's Formula

Let  $P$  be the cubic equation:

$$ax^3 + bx^2 + cx + d = 0, a \neq 0$$

Then  $P$  has solutions:

$$x_1 = S + T - \frac{b}{3a}$$

$$x_2 = -\frac{S+T}{2} - \frac{b}{3a} + \frac{i\sqrt{3}}{2}(S - T)$$

$$x_3 = -\frac{S+T}{2} - \frac{b}{3a} - \frac{i\sqrt{3}}{2}(S - T)$$

where:

$$S = \sqrt[3]{R + \sqrt{Q^3 + R^2}}$$

$$T = \sqrt[3]{R - \sqrt{Q^3 + R^2}}$$

where:

$$Q = \frac{3ac-b^2}{9a^2}$$

$$R = \frac{9abc - 27a^2d - 2b^3}{54a^3}$$

The expression  $D = Q^3 + R^2$  is called the discriminant of the equation.

Let  $a, b, c, d, \in R$  Then:

- If  $D > 0$ , then one root is real and two are complex conjugates.
- If  $D = 0$ , then all roots are real, and at least two are equal.
- If  $D < 0$ , then all roots are real and unequal.

### 1.1 Some Examples

- $x^3 - 2x^2 - 5x + 6 = 0$
- $x^3 - 3x^2 + 4x - 2 = 0$

## References

- [1] J. O'Connor and R. E.F, "Girolamo cardano." <https://mathshistory.st-andrews.ac.uk/Biographies/Cardan/>.
- [2] "Cardano's formula."