

# 25. Numerical Interpolation, Differentiation, and Integration

PHILIP J. DAVIS<sup>1</sup> AND IVAN POLONSKY<sup>2</sup>

## Contents

Formulas	Page
25.1. Differences . . . . .	877
25.2. Interpolation . . . . .	878
25.3. Differentiation . . . . .	882
25.4. Integration . . . . .	885
25.5. Ordinary Differential Equations . . . . .	896
References . . . . .	898
Table 25.1. $n$ -Point Lagrangian Interpolation Coefficients ( $3 \leq n \leq 8$ ) . . . . .	900
$n=3, 4, p=-\left[\frac{n-1}{2}\right] (.01) \left[\frac{n}{2}\right], \text{ Exact}$	
$n=5, 6, p=-\left[\frac{n-1}{2}\right] (.01) \left[\frac{n}{2}\right], 10D$	
$n=7, 8, p=-\left[\frac{n-1}{2}\right] (.1) \left[\frac{n}{2}\right], 10D$	
Table 25.2. $n$ -Point Coefficients for $k$ -th Order Differentiation ( $1 \leq k \leq 5$ ) . . . . .	914
$k=1, n=3(1)6, \text{ Exact}$	
$k=2(1)5, n=k+1(1)6, \text{ Exact}$	
Table 25.3. $n$ -Point Lagrangian Integration Coefficients ( $3 \leq n \leq 10$ ) . . . . .	915
$n=3(1)10, \text{ Exact}$	
Table 25.4. Abscissas and Weight Factors for Gaussian Integration ( $2 \leq n \leq 96$ ) . . . . .	916
$n=2(1)10, 12, 15D$	
$n=16(4)24(8)48(16)96, 21D$	
Table 25.5. Abscissas for Equal Weight Chebyshev Integration ( $2 \leq n \leq 9$ ) . . . . .	920
$n=2(1)7, 9, 10D$	
Table 25.6. Abscissas and Weight Factors for Lobatto Integration ( $3 \leq n \leq 10$ ) . . . . .	920
$n=3(1)10, 8-10D$	
Table 25.7. Abscissas and Weight Factors for Gaussian Integration for Integrands with a Logarithmic Singularity ( $2 \leq n \leq 4$ ) . . . . .	920
$n=2(1)4, 6D$	

<sup>1</sup> National Bureau of Standards.

<sup>2</sup> National Bureau of Standards. (Presently, Bell Tel. Labs., Whippany, N.J.)

# 25. Numerical Interpolation, Differentiation, and Integration

Numerical analysts have a tendency to accumulate a multiplicity of tools each designed for highly specialized operations and each requiring special knowledge to use properly. From the vast stock of formulas available we have culled the present selection. We hope that it will be useful. As with all such compendia, the reader may miss his favorites and find others whose utility he thinks is marginal.

We would have liked to give examples to illuminate the formulas, but this has not been feasible. Numerical analysis is partially a science and partially an art, and short of writing a textbook on the subject it has been impossible to indicate where and under what circumstances the various formulas are useful or accurate, or to elucidate the numerical difficulties to which one might be led by uncritical use. The formulas are therefore issued together with a caveat against their blind application.

## Formulas

*Notation:* Abscissas:  $x_0 < x_1 < \dots$ ; functions:  $f, g, \dots$ ; values:  $f(x_i) = f_i, f'(x_i) = f'_i, f'', f^{(2)}, \dots$  indicate 1<sup>st</sup>, 2<sup>d</sup>,  $\dots$  derivatives. If abscissas are equally spaced,  $x_{i+1} - x_i = h$  and  $f_p = f(x_0 + ph)$  ( $p$  not necessarily integral).  $R, R_n$  indicate remainders.

## 25.1. Differences

### Forward Differences

#### 25.1.1

$$\Delta(f_n) = \Delta_n = \Delta_n^1 = f_{n+1} - f_n$$

$$\Delta_n^2 = \Delta_{n+1}^1 - \Delta_n^1 = f_{n+2} - 2f_{n+1} + f_n$$

$$\Delta_n^3 = \Delta_{n+1}^2 - \Delta_n^2 = f_{n+3} - 3f_{n+2} + 3f_{n+1} - f_n$$

$$\Delta_n^k = \Delta_{n+1}^{k-1} - \Delta_n^{k-1} = \sum_{j=0}^k (-1)^j \binom{k}{j} f_{n+k-j}$$

### Central Differences

#### 25.1.2

$$\delta(f_{n+\frac{1}{2}}) = \delta_{n+\frac{1}{2}} = \delta_{n+\frac{1}{2}}^1 = f_{n+1} - f_n$$

$$\delta_n^2 = \delta_{n+\frac{1}{2}}^1 - \delta_{n-\frac{1}{2}}^1 = f_{n+1} - 2f_n + f_{n-1}$$

$$\delta_{n+\frac{1}{2}}^3 = \delta_{n+1}^2 - \delta_n^2 = f_{n+2} - 3f_{n+1} + 3f_n - f_{n-1}$$

$$\delta_n^{2k} = \sum_{j=0}^{2k} (-1)^j \binom{2k}{j} f_{n+k-j}$$

$$\delta_{n+\frac{1}{2}}^{2k+1} = \sum_{j=0}^{2k+1} (-1)^j \binom{2k+1}{j} f_{n+k+1-j}$$

$$\delta_{\frac{1}{2}n}^k = \Delta_{\frac{1}{2}(n-k)}^k \text{ if } n \text{ and } k \text{ are of same parity.}$$

### Forward Differences

$$x_0 \quad f_0$$

$$x_1 \quad f_1 \quad \Delta_0$$

$$x_2 \quad f_2 \quad \Delta_1 \quad \Delta_0^2$$

$$x_3 \quad f_3 \quad \Delta_2 \quad \Delta_1^2 \quad \Delta_0^3$$

### Central Differences

$$x_{-1} \quad f_{-1}$$

$$x_0 \quad f_0 \quad \delta_{-\frac{1}{2}} \quad \delta_0^2$$

$$x_1 \quad f_1 \quad \delta_{\frac{1}{2}} \quad \delta_1^2 \quad \delta_{\frac{3}{2}}^3$$

$$x_2 \quad f_2 \quad \delta_{3/2}$$

### Mean Differences

#### 25.1.3

$$\mu(f_n) = \frac{1}{2}(f_{n+\frac{1}{2}} + f_{n-\frac{1}{2}})$$

### Divided Differences

#### 25.1.4

$$[x_0, x_1] = \frac{f_0 - f_1}{x_0 - x_1} = [x_1, x_0]$$

$$[x_0, x_1, x_2] = \frac{[x_0, x_1] - [x_1, x_2]}{x_0 - x_2}$$

$$[x_0, x_1, \dots, x_k] = \frac{[x_0, \dots, x_{k-1}] - [x_1, \dots, x_k]}{x_0 - x_k}$$

### Divided Differences in Terms of Functional Values

#### 25.1.5

$$[x_0, x_1, \dots, x_n] = \sum_{k=0}^n \frac{f_k}{\pi'_n(x_k)}$$



**25.1.6** where  $\pi_n(x) = (x-x_0)(x-x_1)\dots(x-x_n)$  and  $\pi'_n(x)$  is its derivative:

**25.1.7**

$$\pi'_n(x_k) = (x_k-x_0)\dots(x_k-x_{k-1})(x_k-x_{k+1})\dots(x_k-x_n)$$

Let  $D$  be a simply connected domain with a piecewise smooth boundary  $C$  and contain the points  $z_0, \dots, z_n$  in its interior. Let  $f(z)$  be analytic in  $D$  and continuous in  $D+C$ . Then,

$$\mathbf{25.1.8} \quad [z_0, z_1, \dots, z_n] = \frac{1}{2\pi i} \int_C \frac{f(z)}{\prod_{k=0}^n (z-z_k)} dz$$

$$\mathbf{25.1.9} \quad \Delta_0^n = h^n f^{(n)}(\xi) \quad (x_0 < \xi < x_n)$$

**25.1.10**

$$[x_0, x_1, \dots, x_n] = \frac{\Delta_0^n}{n!h^n} = \frac{f^{(n)}(\xi)}{n!} \quad (x_0 < \xi < x_n)$$

**25.1.11**

$$[x_{-n}, x_{-n+1}, \dots, x_0, \dots, x_n] = \frac{\delta_0^{2n}}{h^{2n}(2n)!}$$

### Reciprocal Differences

**25.1.12**

$$\rho(x_0, x_1) = \frac{x_0 - x_1}{f_0 - f_1}$$

$$\rho_2(x_0, x_1, x_2) = \frac{x_0 - x_2}{\rho(x_0, x_1) - \rho(x_1, x_2)} + f_1$$

$$\rho_3(x_0, x_1, x_2, x_3) = \frac{x_0 - x_3}{\rho_2(x_0, x_1, x_2) - \rho_2(x_1, x_2, x_3)} + \rho(x_1, x_2)$$

⋮  
⋮  
⋮

$$\rho_n(x_0, x_1, \dots, x_n) = \frac{x_0 - x_n}{\rho_{n-1}(x_0, \dots, x_{n-1}) - \rho_{n-1}(x_1, \dots, x_n)} + \rho_{n-2}(x_1, \dots, x_{n-1})$$

## 25.2. Interpolation

### Lagrange Interpolation Formulas

$$\mathbf{25.2.1} \quad f(x) = \sum_{i=0}^n l_i(x) f_i + R_n(x)$$

**25.2.2**

$$l_i(x) = \frac{\pi_n(x)}{(x-x_i)\pi'_n(x_i)} = \frac{(x-x_0)\dots(x-x_{i-1})(x-x_{i+1})\dots(x-x_n)}{(x_i-x_0)\dots(x_i-x_{i-1})(x_i-x_{i+1})\dots(x_i-x_n)}$$

### Remainder in Lagrange Interpolation Formula

**25.2.3**

$$R_n(x) = \pi_n(x) \cdot [x_0, x_1, \dots, x_n, x] = \pi_n(x) \cdot \frac{f^{(n+1)}(\xi)}{(n+1)!} \quad (x_0 < \xi < x_n)$$

**25.2.4**

$$|R_n(x)| \leq \frac{(x_n - x_0)^{n+1}}{(n+1)!} \max_{x_0 \leq x \leq x_n} |f^{(n+1)}(x)|$$

**25.2.5**

$$R_n(z) = \frac{\pi_n(z)}{2\pi i} \int_C \frac{f(t)}{(t-z)(t-z_0)\dots(t-z_n)} dt$$

The conditions of 25.1.8 are assumed here.

### Lagrange Interpolation, Equally Spaced Abscissas

#### $n$ Point Formula

$$\mathbf{25.2.6} \quad f(x_0 + ph) = \sum_k A_k^n(p) f_k + R_{n-1}$$

$$\text{For } n \text{ even,} \quad \left(-\frac{1}{2}(n-2) \leq k \leq \frac{1}{2}n\right).$$

$$\text{For } n \text{ odd,} \quad \left(-\frac{1}{2}(n-1) \leq k \leq \frac{1}{2}(n-1)\right).$$

**25.2.7**

$$A_k^n(p) = \frac{(-1)^{\frac{1}{2}n+k}}{\left(\frac{n-2}{2}+k\right)!\left(\frac{1}{2}n-k\right)!(p-k)} \prod_{t=1}^n (p+\frac{1}{2}n-t) \quad n \text{ even.}$$

$$A_k^n(p) = \frac{(-1)^{\frac{1}{2}(n-1)+k}}{\left(\frac{n-1}{2}+k\right)!\left(\frac{n-1}{2}-k\right)!(p-k)} \prod_{t=0}^{n-1} \left(p+\frac{n-1}{2}-t\right), \quad n \text{ odd.}$$

**25.2.8**

$$R_{n-1} = \frac{1}{n!} \prod_k (p-k) h^n f^{(n)}(\xi) \approx \frac{1}{n!} \prod_k (p-k) \Delta_0^n \quad (x_0 < \xi < x_n)$$

$k$  has the same range as in 25.2.6.

### Lagrange Two Point Interpolation Formula (Linear Interpolation)

$$\mathbf{25.2.9} \quad f(x_0 + ph) = (1-p)f_0 + pf_1 + R_1$$

$$\mathbf{25.2.10} \quad R_1(p) \approx .125h^2 f^{(2)}(\xi) \approx .125\Delta^2$$

**Lagrange Three Point Interpolation Formula**

25.2.11

$$f(x_0 + ph) = A_{-1}f_{-1} + A_0f_0 + A_1f_1 + R_2$$

$$\approx \frac{p(p-1)}{2}f_{-1} + (1-p^2)f_0 + \frac{p(p+1)}{2}f_1$$

25.2.12

$$R_2(p) \approx .065h^3f^{(3)}(\xi) \approx .065\Delta^3 \quad (|p| \leq 1)$$

**Lagrange Four Point Interpolation Formula**

25.2.13

$$f(x_0 + ph) = A_{-1}f_{-1} + A_0f_0 + A_1f_1 + A_2f_2 + R_3$$

$$\approx \frac{-p(p-1)(p-2)}{6}f_{-1} + \frac{(p^2-1)(p-2)}{2}f_0$$

$$- \frac{p(p+1)(p-2)}{2}f_1 + \frac{p(p^2-1)}{6}f_2$$

25.2.14

 $R_3(p) \approx$ 

$$.024h^4f^{(4)}(\xi) \approx .024\Delta^4 \quad (0 < p < 1)$$

$$.042h^4f^{(4)}(\xi) \approx .042\Delta^4 \quad (-1 < p < 0, 1 < p < 2)$$

$$(x_{-1} < \xi < x_2)$$

**Lagrange Five Point Interpolation Formula**

25.2.15

$$f(x_0 + ph) = \sum_{i=-2}^2 A_i f_i + R_4$$

$$\approx \frac{(p^2-1)p(p-2)}{24}f_{-2} - \frac{(p-1)p(p^2-4)}{6}f_{-1}$$

$$+ \frac{(p^2-1)(p^2-4)}{4}f_0 - \frac{(p+1)p(p^2-4)}{6}f_1$$

$$+ \frac{(p^2-1)p(p+2)}{24}f_2$$

25.2.16

 $R_4(p) \approx$ 

$$.012h^5f^{(5)}(\xi) \approx .012\Delta^5 \quad (|p| < 1)$$

$$.031h^5f^{(5)}(\xi) \approx .031\Delta^5 \quad (1 < |p| < 2) \quad (x_{-2} < \xi < x_2)$$

**Lagrange Six Point Interpolation Formula**

25.2.17

$$f(x_0 + ph) = \sum_{i=-3}^3 A_i f_i + R_5$$

$$\approx \frac{-p(p^2-1)(p-2)(p-3)}{120}f_{-3}$$

$$+ \frac{p(p-1)(p^2-4)(p-3)}{24}f_{-2}$$

$$- \frac{(p^2-1)(p^2-4)(p-3)}{12}f_{-1}$$

$$+ \frac{p(p+1)(p^2-4)(p-3)}{12}f_1 - \frac{p(p^2-1)(p+2)(p-3)}{24}f_2$$

$$+ \frac{p(p^2-1)(p^2-4)}{120}f_3$$

25.2.18

 $R_5(p) \approx$ 

$$.0049h^6f^{(6)}(\xi) \approx .0049\Delta^6 \quad (0 < p < 1)$$

$$.0071h^6f^{(6)}(\xi) \approx .0071\Delta^6 \quad (-1 < p < 0, 1 < p < 2)$$

$$.024h^6f^{(6)}(\xi) \approx .024\Delta^6 \quad (-2 < p < -1, 2 < p < 3)$$

$$(x_{-2} < \xi < x_3)$$

**Lagrange Seven Point Interpolation Formula**

$$25.2.19 \quad f(x_0 + ph) = \sum_{i=-3}^3 A_i f_i + R_6$$

25.2.20

$$R_6(p) \approx \begin{cases} .0025h^7f^{(7)}(\xi) \approx .0025\Delta^7 & (|p| < 1) \\ .0046h^7f^{(7)}(\xi) \approx .0046\Delta^7 & (1 < |p| < 2) \\ .019h^7f^{(7)}(\xi) \approx .019\Delta^7 & (2 < |p| < 3) \end{cases}$$

$$(x_{-3} < \xi < x_3)$$

**Lagrange Eight Point Interpolation Formula**

$$25.2.21 \quad f(x_0 + ph) = \sum_{i=-4}^4 A_i f_i + R_7$$

25.2.22

$$R_7(p) \approx \begin{cases} .0011h^8f^{(8)}(\xi) \approx .0011\Delta^8 & (0 < p < 1) \\ .0014h^8f^{(8)}(\xi) \approx .0014\Delta^8 & (-1 < p < 0) \\ & (1 < p < 2) \\ .0033h^8f^{(8)}(\xi) \approx .0033\Delta^8 & (-2 < p < -1) \\ & (2 < p < 3) \\ .016h^8f^{(8)}(\xi) \approx .016\Delta^8 & (-3 < p < -2) \\ & (3 < p < 4) \end{cases}$$

$$(x_{-4} < \xi < x_4)$$

**Aitken's Iteration Method**

Let  $f(x|x_0, x_1, \dots, x_k)$  denote the unique polynomial of  $k^{\text{th}}$  degree which coincides in value with  $f(x)$  at  $x_0, \dots, x_k$ .

25.2.23

$$f(x|x_0, x_1) = \frac{1}{x_1 - x_0} \begin{vmatrix} f_0 & x_0 - x \\ f_1 & x_1 - x \end{vmatrix}$$

$$f(x|x_0, x_2) = \frac{1}{x_2 - x_0} \begin{vmatrix} f_0 & x_0 - x \\ f_2 & x_2 - x \end{vmatrix}$$

$$f(x|x_0, x_1, x_2) = \frac{1}{x_2 - x_1} \begin{vmatrix} f(x|x_0, x_1) & x_1 - x \\ f(x|x_0, x_2) & x_2 - x \end{vmatrix}$$

$$f(x|x_0, x_1, x_2, x_3) = \frac{1}{x_3 - x_2} \begin{vmatrix} f(x|x_0, x_1, x_2) & x_2 - x \\ f(x|x_0, x_1, x_3) & x_3 - x \end{vmatrix}$$



## Taylor Expansion

25.2.24

$$f(x) = f_0 + (x - x_0)f'_0 + \frac{(x - x_0)^2}{2!}f''_0 + \dots + \frac{(x - x_0)^n}{n!}f^{(n)}_0 + R_n$$

$$25.2.25 \quad R_n = \int_{x_0}^x f^{(n+1)}(t) \frac{(x-t)^n}{n!} dt = \frac{(x - x_0)^{n+1}}{(n+1)!} f^{(n+1)}(\xi) \quad (x_0 < \xi < x)$$

## Newton's Divided Difference Interpolation Formula

25.2.26

$$f(x) = f_0 + \sum_{k=1}^n \pi_{k-1}(x) [x_0, x_1, \dots, x_k] + R_n$$

$x_0$	$f_0$			
		$[x_0, x_1]$		
$x_1$	$f_1$		$[x_0, x_1, x_2]$	
		$[x_1, x_2]$		$[x_0, x_1, x_2, x_3]$
$x_2$	$f_2$		$[x_1, x_2, x_3]$	
		$[x_2, x_3]$		
$x_3$	$f_3$			

25.2.27

$$R_n(x) = \pi_n(x) [x_0, \dots, x_n, x] = \pi_n(x) \frac{f^{(n+1)}(\xi)}{(n+1)!} \quad (x_0 < \xi < x_n)$$

(For  $\pi_n$  see 25.1.6.)

## Newton's Forward Difference Formula

25.2.28

$$f(x_0 + ph) = f_0 + p\Delta_0 + \binom{p}{2}\Delta_0^2 + \dots + \binom{p}{n}\Delta_0^n + R_n$$

$x_0$	$f_0$			
		$\Delta_0$		
$x_1$	$f_1$		$\Delta_0^2$	
		$\Delta_1$		$\Delta_0^3$
$x_2$	$f_2$		$\Delta_1^2$	
		$\Delta_2$		
$x_3$	$f_3$			

25.2.29

$$R_n = h^{n+1} \binom{p}{n+1} f^{(n+1)}(\xi) \approx \binom{p}{n+1} \Delta_0^{n+1} \quad (x_0 < \xi < x_n)$$

## Relation Between Newton and Lagrange Coefficients

25.2.30

$$\binom{p}{2} = A_{-1}^3(p) \quad \binom{p}{3} = -A_{-1}^4(p) \quad \binom{p}{4} = A_2^5(1-p) \quad \binom{p}{5} = A_3^6(2-p)$$

## Everett's Formula

25.2.31

$$f(x_0 + ph) = (1-p)f_0 + pf_1 - \frac{p(p-1)(p-2)}{3!}\delta_0^3 + \frac{(p+1)p(p-1)}{3!}\delta_1^3 + \dots - \binom{p+n-1}{2n+1}\delta_0^{2n} + \binom{p+n}{2n+1}\delta_1^{2n} + R_{2n}$$

$$= (1-p)f_0 + pf_1 + E_2\delta_0^2 + F_2\delta_1^2 + E_4\delta_0^4 + F_4\delta_1^4 + \dots + R_{2n}$$

$x_0$	$f_0$	$\delta_0^2$	$\delta_0^4$
		$\delta_1^2$	$\delta_1^4$
$x_1$	$f_1$	$\delta_1^2$	$\delta_1^4$

25.2.32

$$R_{2n} = h^{2n+2} \binom{p+n}{2n+2} f^{(2n+2)}(\xi) \approx \binom{p+n}{2n+2} \left[ \frac{\Delta_{-n-1}^{2n+2} + \Delta_{-n}^{2n+2}}{2} \right] \quad (x_{-n} < \xi < x_{n+1})$$

## Relation Between Everett and Lagrange Coefficients

25.2.33

$$E_2 = A_{-1}^4 \quad E_4 = A_{-2}^6 \quad E_6 = A_{-3}^8$$

$$F_2 = A_2^4 \quad F_4 = A_3^6 \quad F_6 = A_4^8$$

## Everett's Formula With Throwback (Modified Central Difference)

25.2.34

$$f(x_0 + ph) = (1-p)f_0 + pf_1 + E_2\delta_{m,0}^2 + F_2\delta_{m,1}^2 + R$$

25.2.35

$$\delta_m^2 = \delta^2 - .184\delta^4$$

25.2.36

$$R \approx .00045|\mu\delta_1^4| + .00061|\delta_1^5|$$

25.2.37

$$f(x_0 + ph) = (1-p)f_0 + pf_1 + E_2\delta_0^2 + F_2\delta_1^2 + E_4\delta_{m,0}^4 + F_4\delta_{m,1}^4 + R$$

25.2.38

$$\delta_m^4 = \delta^4 - .207\delta^6 + \dots$$

25.2.39

$$R \approx .000032|\mu\delta_1^6| + .000052|\delta_1^7|$$

25.2.40

$$f(x_0 + ph) = (1-p)f_0 + pf_1 + E_2\delta_0^2 + F_2\delta_1^2 + E_4\delta_0^4 + F_4\delta_1^4 + E_6\delta_{m,0}^6 + F_6\delta_{m,1}^6 + R$$

25.2.41

$$\delta_m^6 = \delta^6 - .218\delta^8 + .049\delta^{10} + \dots$$

25.2.42

$$R \approx .0000037|\mu\delta_1^8| + \dots$$

## Simultaneous Throwback

25.2.43

$$f(x_0 + ph) = (1-p)f_0 + pf_1 + E_2\delta_{m,0}^2 + F_2\delta_{m,1}^2 \\ + E_4\delta_{m,0}^4 + F_4\delta_{m,1}^4 + R$$

$$25.2.44 \quad \delta_m^2 = \delta^2 - .01312\delta^6 + .0043\delta^8 - .001\delta^{10}$$

$$25.2.45 \quad \delta_m^4 = \delta^4 - .27827\delta^6 + .0685\delta^8 - .016\delta^{10}$$

$$25.2.46 \quad R \approx .00000083|\mu\delta_3^6| + .0000094\delta^7$$

## Bessel's Formula With Throwback

25.2.47

$$f(x_0 + ph) = (1-p)f_0 + pf_1 + B_2(\delta_{m,0}^2 + \delta_{m,1}^2) \\ + B_3\delta_{\frac{1}{2}}^3 + R, \quad B_2 = \frac{p(p-1)}{4}, \quad B_3 = \frac{p(p-1)(p-\frac{1}{2})}{6}$$

$$25.2.48 \quad \delta_m^2 = \delta^2 - .184\delta^4$$

$$25.2.49 \quad R \approx .00045|\mu\delta_{\frac{1}{2}}^4| + .00087|\delta_{\frac{1}{2}}^5|$$

## Thiele's Interpolation Formula

25.2.50

$$f(x) = f(x_1) + \frac{x - x_1}{\rho(x_1, x_2) + x - x_2} \frac{\rho_2(x_1, x_2, x_3) - f(x_1) + x - x_3}{\left( \frac{\rho_3(x_1, x_2, x_3, x_4)}{-\rho(x_1, x_2) + \dots} \right)}$$

(For reciprocal differences,  $\rho$ , see 25.1.12.)

## Trigonometric Interpolation

## Gauss' Formula

$$25.2.51 \quad f(x) \approx \sum_{k=0}^{2n} f_k \zeta_k(x) = t_n(x)$$

25.2.52

$$\zeta_k(x) = \frac{\sin \frac{1}{2}(x - x_0) \dots \sin \frac{1}{2}(x - x_{k-1})}{\sin \frac{1}{2}(x_k - x_0) \dots \sin \frac{1}{2}(x_k - x_{k-1})} \\ \frac{\sin \frac{1}{2}(x - x_{k+1}) \dots \sin \frac{1}{2}(x - x_{2n})}{\sin \frac{1}{2}(x_k - x_{k+1}) \dots \sin \frac{1}{2}(x_k - x_{2n})}$$

$t_n(x)$  is a trigonometric polynomial of degree  $n$  such that  $t_n(x_k) = f_k$  ( $k=0, 1, \dots, 2n$ )

## Harmonic Analysis

Equally spaced abscissas

$$x_0 = 0, \quad x_1, \dots, x_{m-1}, x_m = 2\pi$$

25.2.53

$$f(x) \approx \frac{1}{2}a_0 + \sum_{k=1}^n (a_k \cos kx + b_k \sin kx)$$

25.2.54

$$m = 2n + 1$$

$$a_k = \frac{2}{2n+1} \sum_{r=0}^{2n} f_r \cos kx_r; \quad b_k = \frac{2}{2n+1} \sum_{r=0}^{2n} f_r \sin kx_r \\ (k=0, 1, \dots, n)$$

25.2.55

$$m = 2n$$

$$a_k = \frac{1}{n} \sum_{r=0}^{2n-1} f_r \cos kx_r; \quad b_k = \frac{1}{n} \sum_{r=0}^{2n-1} f_r \sin kx_r \\ (k=0, 1, \dots, n) \quad (k=0, 1, \dots, n-1)$$

 $b_n$  is arbitrary.

## Subtabulation

Let  $f(x)$  be tabulated initially in intervals of width  $h$ . It is desired to subtabulate  $f(x)$  in intervals of width  $h/m$ . Let  $\Delta$  and  $\bar{\Delta}$  designate differences with respect to the original and the final intervals respectively. Thus  $\bar{\Delta}_0 = f\left(x_0 + \frac{h}{m}\right) - f(x_0)$ . Assuming that the original 5<sup>th</sup> order differences are zero,

25.2.56

$$\bar{\Delta}_0 = \frac{1}{m} \Delta_0 + \frac{1-m}{2m^2} \Delta_0^2 + \frac{(1-m)(1-2m)}{6m^3} \Delta_0^3 \\ + \frac{(1-m)(1-2m)(1-3m)}{24m^4} \Delta_0^4$$

$$\bar{\Delta}_0^2 = \frac{1}{m^2} \Delta_0^2 + \frac{1-m}{m^3} \Delta_0^3 + \frac{(1-m)(7-11m)}{12m^4} \Delta_0^4$$

$$\bar{\Delta}_0^3 = \frac{1}{m^3} \Delta_0^3 + \frac{3(1-m)}{2m^4} \Delta_0^4$$

$$\bar{\Delta}_0^4 = \frac{1}{m^4} \Delta_0^4$$

From this information we may construct the final tabulation by addition. For  $m=10$ ,

25.2.57

$$\bar{\Delta}_0 = .1\Delta_0 - .045\Delta_0^2 + .0285\Delta_0^3 - .02066\Delta_0^4$$

$$\bar{\Delta}_0^2 = .01\Delta_0^2 - .009\Delta_0^3 + .007725\Delta_0^4$$

$$\bar{\Delta}_0^3 = .001\Delta_0^3 - .00135\Delta_0^4$$

$$\bar{\Delta}_0^4 = .0001\Delta_0^4$$

## Linear Inverse Interpolation

Find  $p$ , given  $f_p (= f(x_0 + ph))$ .

## Linear

25.2.58

$$p \approx \frac{f_p - f_0}{f_1 - f_0}$$



**Quadratic Inverse Interpolation****25.2.59**

$$(f_1 - 2f_0 + f_{-1})p^2 + (f_1 - f_{-1})p + 2(f_0 - f_p) \approx 0$$

**Inverse Interpolation by Reversion of Series****25.2.60** Given  $f(x_0 + ph) = f_p = \sum_{k=0}^{\infty} a_k p^k$ **25.2.61**

$$p = \lambda + c_2 \lambda^2 + c_3 \lambda^3 + \dots, \lambda = (f_p - a_0)/a_1$$

**25.2.62**

$$c_2 = -a_2/a_1$$

$$c_3 = \frac{-a_3}{a_1} + 2 \left( \frac{a_2}{a_1} \right)^2$$

$$c_4 = \frac{-a_4}{a_1} + \frac{5a_2 a_3}{a_1^2} - \frac{5a_2^3}{a_1^3}$$

$$c_5 = \frac{-a_5}{a_1} + \frac{6a_2 a_4}{a_1^2} + \frac{3a_3^2}{a_1^2} - \frac{21a_2^2 a_3}{a_1^3} + \frac{14a_2^4}{a_1^4}$$

**Inversion of Newton's Forward Difference Formula****25.2.63**

$$a_0 = f_0$$

$$a_1 = \Delta_0 - \frac{\Delta_0^2}{2} + \frac{\Delta_0^3}{3} - \frac{\Delta_0^4}{4} + \dots$$

$$a_2 = \frac{\Delta_0^2}{2} - \frac{\Delta_0^3}{2} + \frac{11\Delta_0^4}{24} + \dots$$

$$a_3 = \frac{\Delta_0^3}{6} - \frac{\Delta_0^4}{4} + \dots$$

$$a_4 = \frac{\Delta_0^4}{24} + \dots$$

(Used in conjunction with 25.2.62.)

**Inversion of Everett's Formula****25.2.64**

$$a_0 = f_0$$

$$a_1 = \delta_1 - \frac{\delta_0^2}{3} - \frac{\delta_1^2}{6} + \frac{\delta_0^4}{20} + \frac{\delta_1^4}{30} + \dots$$


$$a_2 = \frac{\delta_0^2}{2} - \frac{\delta_0^4}{24} + \dots$$

$$a_3 = \frac{-\delta_0^3 + \delta_1^3}{6} - \frac{\delta_0^4 + \delta_1^4}{24} + \dots$$


$$a_4 = \frac{\delta_0^4}{24} + \dots$$

$$a_5 = \frac{-\delta_0^4 + \delta_1^4}{120} + \dots$$


(Used in conjunction with 25.2.62.)

**Bivariate Interpolation****Three Point Formula (Linear)****25.2.65**

$$f(x_0 + ph, y_0 + qk) = (1-p-q)f_{0,0} + pf_{1,0} + qf_{0,1} + O(h^2)$$

**Four Point Formula****25.2.66**

$$f(x_0 + ph, y_0 + qk) = (1-p)(1-q)f_{0,0} + p(1-q)f_{1,0} + q(1-p)f_{0,1} + pqf_{1,1} + O(h^2)$$

**Six Point Formula****25.2.67**

$$f(x_0 + ph, y_0 + qk) = \frac{q(q-1)}{2} f_{0,-1} + \frac{p(p-1)}{2} f_{-1,0} + (1+pq-p^2-q^2)f_{0,0} + \frac{p(p-2q+1)}{2} f_{1,0} + \frac{q(q-2p+1)}{2} f_{0,1} + pqf_{1,1} + O(h^3)$$

**25.3. Differentiation****Lagrange's Formula**

$$25.3.1 \quad f'(x) = \sum_{k=0}^n l'_k(x) f_k + R'_n(x)$$

(See 25.2.1.)

$$25.3.2 \quad l'_k(x) = \sum_{j=0, j \neq k}^n \frac{\pi_n(x)}{(x-x_k)(x-x_j)\pi'_n(x_k)}$$

## 25.3.3

$$R'_n(x) = \frac{f^{(n+1)}}{(n+1)!}(\xi)\pi'_n(x) + \frac{\pi_n(x)}{(n+1)!} \frac{d}{dx} f^{(n+1)}(\xi)$$

$$\xi = \xi(x) \quad (x_0 < \xi < x_n)$$

## Equally Spaced Abscissas

## Three Points

## 25.3.4

$$f'_p = f'(x_0 + ph)$$

$$= \frac{1}{h} \left\{ (p - \frac{1}{2})f_{-1} - 2pf_0 + (p + \frac{1}{2})f_1 \right\} + R'_2$$

## Four Points

## 25.3.5

$$f'_p = f'(x_0 + ph) = \frac{1}{h} \left\{ -\frac{3p^2 - 6p + 2}{6} f_{-1} \right.$$

$$+ \frac{3p^2 - 4p - 1}{2} f_0 - \frac{3p^2 - 2p - 2}{2} f_1$$

$$\left. + \frac{3p^2 - 1}{6} f_2 \right\} + R'_3$$

## Five Points

## 25.3.6

$$f'_p = f'(x_0 + ph) = \frac{1}{h} \left\{ \frac{2p^3 - 3p^2 - p + 1}{12} f_{-2} \right.$$

$$- \frac{4p^3 - 3p^2 - 8p + 4}{6} f_{-1} + \frac{2p^3 - 5p}{2} f_0$$

$$- \frac{4p^3 + 3p^2 - 8p - 4}{6} f_1$$

$$\left. + \frac{2p^3 + 3p^2 - p - 1}{12} f_2 \right\} + R'_4$$

For numerical values of differentiation coefficients see **Table 25.2**.

## Markoff's Formulas

(Newton's Forward Difference Formula Differentiated)

## 25.3.7

$$f'(a_0 + ph) = \frac{1}{h} \left[ \Delta_0 + \frac{2p-1}{2} \Delta_0^2 \right.$$

$$\left. + \frac{3p^2 - 6p + 2}{6} \Delta_0^3 + \dots + \frac{d}{dp} \binom{p}{n} \Delta_0^n \right] + R'_n$$

## 25.3.8

$$R'_n = h^n f^{(n+1)}(\xi) \frac{d}{dp} \binom{p}{n+1} + h^{n+1} \binom{p}{n+1} \frac{d}{dx} f^{(n+1)}(\xi)$$

$$(a_0 < \xi < a_n)$$

$$25.3.9 \quad hf'_0 = \Delta_0 - \frac{1}{2} \Delta_0^2 + \frac{1}{3} \Delta_0^3 - \frac{1}{4} \Delta_0^4 + \dots$$

$$25.3.10 \quad h^2 f_0^{(2)} = \Delta_0^2 - \Delta_0^3 + \frac{11}{12} \Delta_0^4 - \frac{5}{6} \Delta_0^5 + \dots$$

## 25.3.11

$$h^3 f_0^{(3)} = \Delta_0^3 - \frac{3}{2} \Delta_0^4 + \frac{7}{4} \Delta_0^5 - \frac{15}{8} \Delta_0^6 + \dots$$

## 25.3.12

$$h^4 f_0^{(4)} = \Delta_0^4 - 2\Delta_0^5 + \frac{17}{6} \Delta_0^6 - \frac{7}{2} \Delta_0^7 + \dots$$

## 25.3.13

$$h^5 f_0^{(5)} = \Delta_0^5 - \frac{5}{2} \Delta_0^6 + \frac{25}{6} \Delta_0^7 - \frac{35}{6} \Delta_0^8 + \dots$$

## Everett's Formula

## 25.3.14

$$hf'(x_0 + ph) \approx -f_0 + f_1 - \frac{3p^2 - 6p + 2}{6} \delta_0^2 + \frac{3p^2 - 1}{6} \delta_1^2$$

$$- \frac{5p^4 - 20p^3 + 15p^2 + 10p - 6}{120} \delta_0^4 + \frac{5p^4 - 15p^2 + 4}{120} \delta_1^4$$

$$+ \dots - \left[ \binom{p+n-1}{2n+1} \right]' \delta_0^{2n} + \left[ \binom{p+n}{2n+1} \right]' \delta_1^{2n}$$

## 25.3.15

$$hf'_0 \approx -f_0 + f_1 - \frac{1}{3} \delta_0^2 - \frac{1}{6} \delta_1^2 + \frac{1}{20} \delta_0^4 + \frac{1}{30} \delta_1^4$$

## Differences in Terms of Derivatives

## 25.3.16

$$\Delta_0 \approx hf'_0 + \frac{h^2}{2!} f_0^{(2)} + \frac{h^3}{3!} f_0^{(3)} + \frac{h^4}{4!} f_0^{(4)} + \frac{h^5}{5!} f_0^{(5)}$$

## 25.3.17

$$\Delta_0^2 \approx h^2 f_0^{(2)} + h^3 f_0^{(3)} + \frac{7}{12} h^4 f_0^{(4)} + \frac{1}{4} h^5 f_0^{(5)}$$

## 25.3.18

$$\Delta_0^3 \approx h^3 f_0^{(3)} + \frac{3}{2} h^4 f_0^{(4)} + \frac{5}{4} h^5 f_0^{(5)}$$

## 25.3.19


$$\Delta_0^4 \approx h^4 f_0^{(4)} + 2h^5 f_0^{(5)}$$

## 25.3.20

$$\Delta_0^5 \approx h^5 f_0^{(5)}$$

## Partial Derivatives


## 25.3.21



$$\frac{\partial f_{0,0}}{\partial x} = \frac{1}{2h} (f_{1,0} - f_{-1,0}) + O(h^2)$$




25.3.22




$$\frac{\partial f_{0,0}}{\partial x} = \frac{1}{4h} (f_{1,1} - f_{-1,1} + f_{1,-1} - f_{-1,-1}) + O(h^2)$$

25.3.23




$$\frac{\partial^2 f_{0,0}}{\partial x^2} = \frac{1}{h^2} (f_{1,0} - 2f_{0,0} + f_{-1,0}) + O(h^2)$$

25.3.24




$$\frac{\partial^2 f_{0,0}}{\partial x^2} = \frac{1}{12h^2} (-f_{2,0} + 16f_{1,0} - 30f_{0,0} + 16f_{-1,0} - f_{-2,0}) + O(h^4)$$

25.3.25




$$\frac{\partial^2 f_{0,0}}{\partial x^2} = \frac{1}{3h^2} (f_{1,1} - 2f_{0,1} + f_{-1,1} + f_{1,0} - 2f_{0,0} + f_{-1,0} + f_{1,-1} - 2f_{0,-1} + f_{-1,-1}) + O(h^2)$$

25.3.26




$$\frac{\partial^2 f_{0,0}}{\partial x \partial y} = \frac{1}{4h^2} (f_{1,1} - f_{1,-1} - f_{-1,1} + f_{-1,-1}) + O(h^2)$$

25.3.27




$$\frac{\partial^2 f_{0,0}}{\partial x \partial y} = \frac{-1}{2h^2} (f_{1,0} + f_{-1,0} + f_{0,1} + f_{0,-1} - 2f_{0,0} - f_{1,1} - f_{-1,-1}) + O(h^2)$$

25.3.28



$$\frac{\partial^4 f_{0,0}}{\partial x^4} = \frac{1}{h^4} (f_{2,0} - 4f_{1,0} + 6f_{0,0} - 4f_{-1,0} + f_{-2,0}) + O(h^2)$$


25.3.29



$$\frac{\partial^4 f_{0,0}}{\partial x^2 \partial y^2} = \frac{1}{h^4} (f_{1,1} + f_{-1,1} + f_{1,-1} + f_{-1,-1} - 2f_{1,0} - 2f_{-1,0} - 2f_{0,1} - 2f_{0,-1} + 4f_{0,0}) + O(h^2)$$


## Laplacian

25.3.30



$$\begin{aligned}\nabla^2 u_{0,0} &= \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} \right)_{0,0} \\ &= \frac{1}{h^2} (u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1} - 4u_{0,0}) + O(h^2)\end{aligned}$$


25.3.31



$$\begin{aligned}\nabla^4 u_{0,0} &= \frac{1}{12h^2} [-60u_{0,0} + 16(u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1}) \\ &\quad - (u_{2,0} + u_{0,2} + u_{-2,0} + u_{0,-2})] + O(h^4)\end{aligned}$$


## Bi-harmonic Operator

25.3.32



$$\begin{aligned}\nabla^4 u_{0,0} &= \left( \frac{\partial^4 u}{\partial x^4} + 2 \frac{\partial^4 u}{\partial x^2 \partial y^2} + \frac{\partial^4 u}{\partial y^4} \right)_{0,0} \\ &= \frac{1}{h^4} [20u_{0,0} - 8(u_{1,0} + u_{0,1} + u_{-1,0} + u_{0,-1}) \\ &\quad + 2(u_{1,1} + u_{1,-1} + u_{-1,1} + u_{-1,-1}) \\ &\quad + (u_{0,2} + u_{2,0} + u_{-2,0} + u_{0,-2})] + O(h^2)\end{aligned}$$

25.3.33



$$\begin{aligned}\nabla^4 u_{0,0} &= \frac{1}{6h^4} [-(u_{0,3} + u_{0,-3} + u_{3,0} + u_{-3,0}) \\ &\quad + 14(u_{0,2} + u_{0,-2} + u_{2,0} + u_{-2,0}) \\ &\quad - 77(u_{0,1} + u_{0,-1} + u_{1,0} + u_{-1,0}) \\ &\quad + 184u_{0,0} + 20(u_{1,1} + u_{1,-1} + u_{-1,1} + u_{-1,-1}) \\ &\quad - (u_{1,2} + u_{2,1} + u_{1,-2} + u_{2,-1} + u_{-1,2} + u_{-2,1} \\ &\quad + u_{-1,-2} + u_{-2,-1})] + O(h^4)\end{aligned}$$

## 25.4. Integration

## Trapezoidal Rule

25.4.1

$$\begin{aligned}\int_{x_0}^{x_1} f(x) dx &= \frac{h}{2} (f_0 + f_1) - \frac{1}{2} \int_{x_0}^{x_1} (t - x_0)(x_1 - t) f''(t) dt \\ &= \frac{h}{2} (f_0 + f_1) - \frac{h^3}{12} f''(\xi) \quad (x_0 < \xi < x_1)\end{aligned}$$

## Extended Trapezoidal Rule

25.4.2

$$\begin{aligned}\int_{x_0}^{x_m} f(x) dx &= h \left[ \frac{f_0}{2} + f_1 + \dots + f_{m-1} + \frac{f_m}{2} \right] \\ &\quad - \frac{mh^3}{12} f''(\xi)\end{aligned}$$

## Error Term in Trapezoidal Formula for Periodic Functions

If  $f(x)$  is periodic and has a continuous  $k^{\text{th}}$  derivative, and if the integral is taken over a period, then

$$25.4.3 \quad |\text{Error}| \leq \frac{\text{constant}}{m^k}$$

## Modified Trapezoidal Rule

25.4.4

$$\begin{aligned}\int_{x_0}^{x_m} f(x) dx &= h \left[ \frac{f_0}{2} + f_1 + \dots + f_{m-1} + \frac{f_m}{2} \right] \\ &\quad + \frac{h}{24} [-f_{-1} + f_1 + f_{m-1} - f_{m+1}] + \frac{11m}{720} h^5 f^{(4)}(\xi)\end{aligned}$$



**Simpson's Rule****25.4.5**

$$\begin{aligned}\int_{x_0}^{x_2} f(x) dx &= \frac{h}{3} [f_0 + 4f_1 + f_2] \\ &\quad + \frac{1}{6} \int_{x_0}^{x_1} (x_0 - t)^2 (x_1 - t) f^{(3)}(t) dt \\ &\quad + \frac{1}{6} \int_{x_1}^{x_2} (x_2 - t)^2 (x_1 - t) f^{(3)}(t) dt \\ &= \frac{h}{3} [f_0 + 4f_1 + f_2] - \frac{h^5}{90} f^{(4)}(\xi)\end{aligned}$$

**Extended Simpson's Rule****25.4.6**

$$\begin{aligned}\int_{x_0}^{x_{2n}} f(x) dx &= \frac{h}{3} [f_0 + 4(f_1 + f_3 + \dots + f_{2n-1}) \\ &\quad + 2(f_2 + f_4 + \dots + f_{2n-2}) + f_{2n}] - \frac{nh^5}{90} f^{(4)}(\xi)\end{aligned}$$

**Euler-Maclaurin Summation Formula****25.4.7**

$$\begin{aligned}\int_{x_0}^{x_n} f(x) dx &= h \left[ \frac{f_0}{2} + f_1 + f_2 + \dots + f_{n-1} + \frac{f_n}{2} \right] \\ &\quad - \frac{B_2}{2!} h^2 (f'_n - f'_0) - \dots - \frac{B_{2k} h^{2k}}{(2k)!} [f_n^{(2k-1)} - f_0^{(2k-1)}] + R_{2k} \\ R_{2k} &= \frac{\theta n B_{2k+2} h^{2k+3}}{(2k+2)!} \max_{x_0 \leq x \leq x_n} |f^{(2k+2)}(x)|, \quad (-1 \leq \theta \leq 1)\end{aligned}$$

(For  $B_{2k}$ , Bernoulli numbers, see chapter 23.)

If  $f^{(2k+2)}(x)$  and  $f^{(2k+4)}(x)$  do not change sign for  $x_0 < x < x_n$  then  $|R_{2k}|$  is less than the first neglected term. If  $f^{(2k+2)}(x)$  does not change sign for  $x_0 < x < x_n$ ,  $|R_{2k}|$  is less than twice the first neglected term.

**Lagrange Formula****25.4.8**

$$\int_a^b f(x) dx = \sum_{i=0}^n (L_i^{(n)}(b) - L_i^{(n)}(a)) f_i + R_n$$

(See 25.2.1.)

**25.4.9**

$$L_i^{(n)}(x) = \frac{1}{\pi_n'(x_i)} \int_{x_0}^x \frac{\pi_n(t)}{t - x_i} dt = \int_{x_0}^x l_i(t) dt$$

$$25.4.10 \quad R_n = \frac{1}{(n+1)!} \int_a^b \pi_n(x) f^{(n+1)}(\xi(x)) dx$$

**Equally Spaced Abscissas****25.4.11**

$$\int_{x_0}^{x_k} f(x) dx = \frac{1}{h^n} \sum_{i=0}^n f_i \frac{(-1)^{n-i}}{i!(n-i)!} \int_{x_0}^{x_k} \frac{\pi_n(x)}{x - x_i} dx + R_n$$

$$25.4.12 \quad \int_{x_m}^{x_{m+1}} f(x) dx = h \sum_{i=-\lfloor \frac{n-1}{2} \rfloor}^{\lfloor \frac{n}{2} \rfloor} A_i(m) f_i + R_n \quad *$$

(See Table 25.3 for  $A_i(m)$ .)**Newton-Cotes Formulas (Closed Type)**

(For Trapezoidal and Simpson's Rules see 25.4.1-25.4.6.)

**25.4.13** (Simpson's  $\frac{3}{8}$  rule)

$$\int_{x_0}^{x_3} f(x) dx = \frac{3h}{8} (f_0 + 3f_1 + 3f_2 + f_3) - \frac{3f^{(4)}(\xi)h^5}{80}$$

**25.4.14** (Bode's rule)

$$\begin{aligned}\int_{x_0}^{x_4} f(x) dx &= \frac{2h}{45} (7f_0 + 32f_1 + 12f_2 \\ &\quad + 32f_3 + 7f_4) - \frac{8f^{(6)}(\xi)h^7}{945}\end{aligned}$$

**25.4.15**

$$\begin{aligned}\int_{x_0}^{x_5} f(x) dx &= \frac{5h}{288} (19f_0 + 75f_1 + 50f_2 + 50f_3 \\ &\quad + 75f_4 + 19f_5) - \frac{275f^{(6)}(\xi)h^7}{12096}\end{aligned}$$

**25.4.16**

$$\begin{aligned}\int_{x_0}^{x_6} f(x) dx &= \frac{h}{140} (41f_0 + 216f_1 + 27f_2 + 272f_3 \\ &\quad + 27f_4 + 216f_5 + 41f_6) - \frac{9f^{(8)}(\xi)h^9}{1400}\end{aligned}$$

**25.4.17**

$$\begin{aligned}\int_{x_0}^{x_7} f(x) dx &= \frac{7h}{17280} (751f_0 + 3577f_1 + 1323f_2 \\ &\quad + 2989f_3 + 2989f_4 + 1323f_5 + 3577f_6 \\ &\quad + 751f_7) - \frac{8183f^{(8)}(\xi)h^9}{518400}\end{aligned}$$

**25.4.18**

$$\begin{aligned}\int_{x_0}^{x_8} f(x) dx &= \frac{4h}{14175} (989f_0 + 5888f_1 - 928f_2 \\ &\quad + 10496f_3 - 4540f_4 + 10496f_5 - 928f_6 + 5888f_7 \\ &\quad + 989f_8) - \frac{2368}{467775} f^{(10)}(\xi)h^{11}\end{aligned}$$

**25.4.19**

$$\begin{aligned}\int_{x_0}^{x_9} f(x) dx &= \frac{9h}{89600} \{ 2857(f_0 + f_9) \\ &\quad + 15741(f_1 + f_8) + 1080(f_2 + f_7) + 19344(f_3 + f_6) \\ &\quad + 5778(f_4 + f_5) \} - \frac{173}{14620} f^{(10)}(\xi)h^{11}\end{aligned}$$

\*See page 11.

**25.4.20**

$$\int_{x_0}^{x_{10}} f(x) dx = \frac{5h}{299376} \{16067(f_0 + f_{10}) + 106300(f_1 + f_9) - 48525(f_2 + f_8) + 272400(f_3 + f_7) - 260550(f_4 + f_6) + 427368f_5\} - \frac{1346350}{326918592} f^{(12)}(\xi) h^{13}$$

**Newton-Cotes Formulas (Open Type)****25.4.21**

$$\int_{x_0}^{x_3} f(x) dx = \frac{3h}{2} (f_1 + f_2) + \frac{f^{(2)}(\xi) h^3}{4}$$

**25.4.22**

$$\int_{x_0}^{x_4} f(x) dx = \frac{4h}{3} (2f_1 - f_2 + 2f_3) + \frac{28f^{(4)}(\xi) h^5}{90}$$

**25.4.23**

$$\int_{x_0}^{x_5} f(x) dx = \frac{5h}{24} (11f_1 + f_2 + f_3 + 11f_4) + \frac{95f^{(4)}(\xi) h^5}{144}$$

**25.4.24**


$$\int_{x_0}^{x_6} f(x) dx = \frac{6h}{20} (11f_1 - 14f_2 + 26f_3 - 14f_4 + 11f_5) + \frac{41f^{(6)}(\xi) h^7}{140}$$

**25.4.25**

$$\int_{x_0}^{x_7} f(x) dx = \frac{7h}{1440} (611f_1 - 453f_2 + 562f_3 + 562f_4 - 453f_5 + 611f_6) + \frac{5257}{8640} f^{(6)}(\xi) h^7$$

**25.4.26**

$$\int_{x_0}^{x_8} f(x) dx = \frac{8h}{945} (460f_1 - 954f_2 + 2196f_3 - 2459f_4 + 2196f_5 - 954f_6 + 460f_7) + \frac{3956}{14175} f^{(8)}(\xi) h^9$$

**Five Point Rule for Analytic Functions****25.4.27**

$$\int_{z_0-h}^{z_0+h} f(z) dz = \frac{h}{15} \{24f(z_0) + 4[f(z_0+h) + f(z_0-h)] - [f(z_0+ih) + f(z_0-ih)]\} + R$$

$|R| \leq \frac{|h|^7}{1890} \max_{z \in S} |f^{(6)}(z)|$ ,  $S$  designates the square with vertices  $z_0 + i^k h$  ( $k=0, 1, 2, 3$ );  $h$  can be complex.

**Chebyshev's Equal Weight Integration Formula**

$$25.4.28 \quad \int_{-1}^1 f(x) dx = \frac{2}{n} \sum_{i=1}^n f(x_i) + R_n$$

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of the polynomial part of

$$x^n \exp \left[ \frac{-n}{2 \cdot 3x^2} - \frac{n}{4 \cdot 5x^3} - \frac{n}{6 \cdot 7x^4} - \dots \right]$$

(See **Table 25.5** for  $x_i$ .)

For  $n=8$  and  $n \geq 10$  some of the zeros are complex.

Remainder:

$$R_n = \int_{-1}^{+1} \frac{x^{n+1}}{(n+1)!} f^{(n+1)}(\xi) dx - \frac{2}{n(n+1)!} \sum_{i=1}^n x_i^{n+1} f^{(n+1)}(\xi_i)$$

where  $\xi = \xi(x)$  satisfies  $0 \leq \xi \leq x$  and  $0 \leq \xi_i \leq x_i$

$$(i=1, \dots, n)$$

**Integration Formulas of Gaussian Type**

(For Orthogonal Polynomials see chapter 22)

**Gauss' Formula**

$$25.4.29 \quad \int_{-1}^1 f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Legendre polynomials  $P_n(x)$ ,  $P_n(1)=1$

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of  $P_n(x)$

Weights:  $w_i = 2/(1-x_i^2) [P'_n(x_i)]^2$   
(See **Table 25.4** for  $x_i$  and  $w_i$ .)

$$R_n = \frac{2^{2n+1}(n!)^4}{(2n+1)[(2n)!]^3} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

**Gauss' Formula, Arbitrary Interval**

$$25.4.30 \quad \int_a^b f(y) dy = \frac{b-a}{2} \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \left( \frac{b-a}{2} \right) x_i + \left( \frac{b+a}{2} \right)$$



Related orthogonal polynomials:  $P_n(x)$ ,  $P_n(1)=1$

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of  $P_n(x)$

\* Weights:  $w_i = 2/(1-x_i^2) [P'_n(x_i)]^2$

$$* R_n = \frac{(b-a)^{2n+1} (n!)^4}{(2n+1) [(2n)!]^3} f^{(2n)}(\xi)$$

#### Radau's Integration Formula

##### 25.4.31

$$\int_{-1}^1 f(x) dx = \frac{2}{n^2} f_{-1} + \sum_{i=1}^{n-1} w_i f(x_i) + R_n$$

Related polynomials:

$$\frac{P_{n-1}(x) + P_n(x)}{x+1}$$

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of

$$\frac{P_{n-1}(x) + P_n(x)}{x+1}$$

Weights:

$$w_i = \frac{1}{n^2} \frac{1-x_i}{[P_{n-1}(x_i)]^2} = \frac{1}{1-x_i} \frac{1}{[P'_{n-1}(x_i)]^2}$$

Remainder:

$$R_n = \frac{2^{2n-1} \cdot n}{[(2n-1)!]^3} [(n-1)!]^4 f^{(2n-1)}(\xi) \quad (-1 < \xi < 1)$$

#### Lobatto's Integration Formula

##### 25.4.32

$$\int_{-1}^1 f(x) dx = \frac{2}{n(n-1)} [f(1) + f(-1)] + \sum_{i=2}^{n-1} w_i f(x_i) + R_n$$

Related polynomials:  $P'_{n-1}(x)$

Abscissas:  $x_i$  is the  $(i-1)^{\text{st}}$  zero of  $P'_{n-1}(x)$

Weights:

$$w_i = \frac{2}{n(n-1) [P'_{n-1}(x_i)]^2} \quad (x_i \neq \pm 1)$$

(See Table 25.6 for  $x_i$  and  $w_i$ .)

Remainder:

$$R_n = \frac{-n(n-1)^3 2^{2n-1} [(n-2)!]^4}{(2n-1) [(2n-2)!]^3} f^{(2n-2)}(\xi) \quad (-1 < \xi < 1)$$

$$25.4.33 \quad \int_0^1 x^k f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$q_n(x) = \sqrt{k+2n+1} P_n^{(k,0)}(1-2x)$$

(For the Jacobi polynomials  $P_n^{(k,0)}$  see chapter 22.)

Abscissas:

$x_i$  is the  $i^{\text{th}}$  zero of  $q_n(x)$

Weights:

$$w_i = \left\{ \sum_{j=0}^{n-1} [q_j(x_i)]^2 \right\}^{-1} *$$

(See Table 25.8 for  $x_i$  and  $w_i$ .)

Remainder:

$$R_n = \frac{f^{(2n)}(\xi)}{(k+2n+1)(2n)!} \left[ \frac{n!(k+n)!}{(k+2n)!} \right]^2 \quad (0 < \xi < 1)$$

##### 25.4.34

$$\int_0^1 f(x) \sqrt{1-x} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{1-x}} P_{2n+1}(\sqrt{1-x}), P_{2n+1}(1)=1$$

Abscissas:  $x_i = 1 - \xi_i^2$  where  $\xi_i$  is the  $i^{\text{th}}$  positive zero of  $P_{2n+1}(x)$ .

Weights:  $w_i = 2\xi_i^2 w_i^{(2n+1)}$  where  $w_i^{(2n+1)}$  are the Gaussian weights of order  $2n+1$ .

Remainder:

$$R_n = \frac{2^{4n+3} [(2n+1)!]^4}{(2n)!(4n+3)[(4n+2)!]^2} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

##### 25.4.35

$$\int_a^b f(y) \sqrt{b-y} dy = (b-a)^{3/2} \sum_{i=1}^n w_i f(y_i)$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{1-x}} P_{2n+1}(\sqrt{1-x}), P_{2n+1}(1)=1$$

Abscissas:  $x_i = 1 - \xi_i^2$  where  $\xi_i$  is the  $i^{\text{th}}$  positive zero of  $P_{2n+1}(x)$ .

Weights:  $w_i = 2\xi_i^2 w_i^{(2n+1)}$  where  $w_i^{(2n+1)}$  are the Gaussian weights of order  $2n+1$ .

$$25.4.36 \quad \int_0^1 \frac{f(x)}{\sqrt{1-x}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$P_{2n}(\sqrt{1-x}), P_{2n}(1)=1$$

Abscissas:  $x_i = 1 - \xi_i^2$  where  $\xi_i$  is the  $i^{\text{th}}$  positive zero of  $P_{2n}(x)$ .

Weights:  $w_i = 2w_i^{(2n)}$ ,  $w_i^{(2n)}$  are the Gaussian weights of order  $2n$ .

Remainder:

$$R_n = \frac{2^{4n+1}}{4n+1} \frac{[(2n)!]^3}{[(4n)!]^2} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

$$25.4.37 \quad \int_a^b \frac{f(y)}{\sqrt{b-y}} dy = \sqrt{b-a} \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$P_{2n}(\sqrt{1-x}), P_{2n}(1)=1$$

Abscissas:

$x_i = 1 - \xi_i^2$  where  $\xi_i$  is the  $i^{\text{th}}$  positive zero of  $P_{2n}(x)$ .

Weights:  $w_i = 2w_i^{(2n)}$ ,  $w_i^{(2n)}$  are the Gaussian weights of order  $2n$ .

$$25.4.38 \quad \int_{-1}^{+1} \frac{f(x)}{\sqrt{1-x^2}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Chebyshev Polynomials of First Kind

$$T_n(x), T_n(1) = \frac{1}{2^{n-1}}$$

Abscissas:

$$x_i = \cos \frac{(2i-1)\pi}{2n}$$

Weights:

$$w_i = \frac{\pi}{n}$$

Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{2n-1}} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

25.4.39

$$\int_a^b \frac{f(y) dy}{\sqrt{(y-a)(b-y)}} = \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \frac{b+a}{2} + \frac{b-a}{2} x_i$$

Related orthogonal polynomials:

$$T_n(x), T_n(1) = \frac{1}{2^{n-1}}$$

Abscissas:

$$x_i = \cos \frac{(2i-1)\pi}{2n}$$

Weights:

$$w_i = \frac{\pi}{n}$$

25.4.40

$$\int_{-1}^{+1} f(x) \sqrt{1-x^2} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Chebyshev Polynomials of Second Kind

$$U_n(x) = \frac{\sin [(n+1) \arccos x]}{\sin (\arccos x)} \quad *$$

Abscissas:

$$x_i = \cos \frac{i}{n+1} \pi \quad *$$

Weights:

$$w_i = \frac{\pi}{n+1} \sin^2 \frac{i}{n+1} \pi \quad *$$

Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{2n+1}} f^{(2n)}(\xi) \quad (-1 < \xi < 1)$$

25.4.41

$$\int_a^b \sqrt{(y-a)(b-y)} f(y) dy = \left(\frac{b-a}{2}\right)^2 \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = \frac{b+a}{2} + \frac{b-a}{2} x_i$$

Related orthogonal polynomials:

$$U_n(x) = \frac{\sin [(n+1) \arccos x]}{\sin (\arccos x)} \quad *$$

Abscissas:

$$x_i = \cos \frac{i}{n+1} \pi \quad *$$

Weights:

$$w_i = \frac{\pi}{n+1} \sin^2 \frac{i}{n+1} \pi \quad *$$

$$25.4.42 \quad \int_0^1 f(x) \sqrt{\frac{x}{1-x}} dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{x}} T_{2n+1}(\sqrt{x})$$

Abscissas:

$$x_i = \cos^2 \frac{2i-1}{2n+1} \cdot \frac{\pi}{2}$$

Weights:

$$w_i = \frac{2\pi}{2n+1} x_i$$



Remainder:

$$R_n = \frac{\pi}{(2n)! 2^{4n+1}} f^{(2n)}(\xi) \quad (0 < \xi < 1)$$

**25.4.43**

$$\int_a^b f(x) \sqrt{\frac{x-a}{b-x}} dx = (b-a) \sum_{i=1}^n w_i f(y_i) + R_n$$

$$y_i = a + (b-a)x_i$$

Related orthogonal polynomials:

$$\frac{1}{\sqrt{x}} T_{2n+1}(\sqrt{x})$$

Abscissas:

$$x_i = \cos^2 \frac{2i-1}{2n+1} \cdot \frac{\pi}{2}$$

Weights:

$$w_i = \frac{2\pi}{2n+1} x_i$$

**25.4.44** 
$$\int_0^1 \ln x f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: polynomials orthogonal with respect to the weight function  $-\ln x$

Abscissas: See **Table 25.7**

Weights: See **Table 25.7**

**25.4.45**

$$\int_0^\infty e^{-x} f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Laguerre polynomials  $L_n(x)$ .

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of  $L_n(x)$

Weights:

$$* \quad w_i = \frac{x_i}{(n+1)^2 [L_{n+1}(x_i)]^2}$$

(See **Table 25.9** for  $x_i$  and  $w_i$ .)

Remainder:

$$R_n = \frac{(n!)^2}{(2n)!} f^{(2n)}(\xi) \quad (0 < \xi < \infty)$$

**25.4.46**

$$\int_{-\infty}^\infty e^{-x^2} f(x) dx = \sum_{i=1}^n w_i f(x_i) + R_n$$

Related orthogonal polynomials: Hermite polynomials  $H_n(x)$ .

Abscissas:  $x_i$  is the  $i^{\text{th}}$  zero of  $H_n(x)$

Weights:

$$\frac{2^{n-1} n! \sqrt{\pi}}{n^2 [H_{n-1}(x_i)]^2}$$

(See **Table 25.10** for  $x_i$  and  $w_i$ .)

\*See page II.

Remainder:

$$R_n = \frac{n! \sqrt{\pi}}{2^n (2n)!} f^{(2n)}(\xi) \quad (-\infty < \xi < \infty)$$

**Filon's Integration Formula**<sup>3</sup>

**25.4.47**

$$\int_{x_0}^{x_n} f(x) \cos tx dx = h \left[ \alpha(th) (f_{2n} \sin tx_{2n} - f_0 \sin tx_0) + \beta(th) \cdot C_{2n} + \gamma(th) \cdot C_{2n-1} + \frac{2}{45} th^4 S'_{2n-1} \right] - R_n$$

**25.4.48**

$$C_{2n} = \sum_{i=0}^n f_{2i} \cos(tx_{2i}) - \frac{1}{2} [f_{2n} \cos tx_{2n} + f_0 \cos tx_0]$$

**25.4.49**

$$C_{2n-1} = \sum_{i=1}^n f_{2i-1} \cos tx_{2i-1}$$

**25.4.50**

$$S'_{2n-1} = \sum_{i=1}^n f_{2i-1}^{(3)} \sin tx_{2i-1}$$

**25.4.51**

$$R_n = \frac{1}{90} nh^5 f^{(4)}(\xi) + O(th^7)$$

**25.4.52**

$$\alpha(\theta) = \frac{1}{\theta} + \frac{\sin 2\theta}{2\theta^2} - \frac{2 \sin^2 \theta}{\theta^3}$$

$$\beta(\theta) = 2 \left( \frac{1 + \cos^2 \theta}{\theta^2} - \frac{\sin 2\theta}{\theta^3} \right)$$

$$\gamma(\theta) = 4 \left( \frac{\sin \theta}{\theta^3} - \frac{\cos \theta}{\theta^2} \right)$$

For small  $\theta$  we have

**25.4.53**

$$\alpha = \frac{2\theta^3}{45} - \frac{2\theta^5}{315} + \frac{2\theta^7}{4725} - \dots$$

$$\beta = \frac{2}{3} + \frac{2\theta^2}{15} - \frac{4\theta^4}{105} + \frac{2\theta^6}{567} - \dots$$

$$\gamma = \frac{4}{3} - \frac{2\theta^2}{15} + \frac{\theta^4}{210} - \frac{\theta^6}{11340} + \dots$$

**25.4.54**

$$\int_{x_0}^{x_{2n}} f(x) \sin tx dx = h \left[ \alpha(th) (f_0 \cos tx_0 - f_{2n} \cos tx_{2n}) + \beta S_{2n} + \gamma S_{2n-1} + \frac{2}{45} th^4 C'_{2n-1} \right] - R_n$$

**25.4.55**

$$S_{2n} = \sum_{i=0}^n f_{2i} \sin(tx_{2i}) - \frac{1}{2} [f_{2n} \sin(tx_{2n}) + f_0 \sin(tx_0)]$$

<sup>3</sup> For certain difficulties associated with this formula, see the article by J. W. Tukey, p. 400, "On Numerical Approximation," Ed. R. E. Langer, Madison, 1959.

$$25.4.56 \quad S_{2n-1} = \sum_{i=1}^n f_{2i-1} \sin(tx_{2i-1})$$

$$25.4.57 \quad C'_{2n-1} = \sum_{i=1}^n f_{2i-1}^{(3)} \cos(tx_{2i-1})$$

(See Table 25.11 for  $\alpha, \beta, \gamma$ .)

#### Iterated Integrals

25.4.58

$$\begin{aligned} \int_0^x dt_n \int_0^{t_n} dt_{n-1} \dots \int_0^{t_3} dt_2 \int_0^{t_2} f(t_1) dt_1 \\ = \frac{1}{(n-1)!} \int_0^x (x-t)^{n-1} f(t) dt \end{aligned}$$


25.4.59

$$\begin{aligned} \int_a^x dt_n \int_a^{t_n} dt_{n-1} \dots \int_a^{t_3} dt_2 \int_a^{t_2} f(t_1) dt_1 \\ = \frac{(x-a)^n}{(n-1)!} \int_0^1 t^{n-1} f(x-(x-a)t) dt \end{aligned}$$

#### Multidimensional Integration

Circumference of Circle  $\Gamma$ :  $x^2 + y^2 = h^2$ .


25.4.60



$$\begin{aligned} \frac{1}{2\pi h} \int_{\Gamma} f(x, y) ds = \frac{1}{2m} \sum_{n=1}^{2m} f\left(h \cos \frac{\pi n}{m}, h \sin \frac{\pi n}{m}\right) \\ + O(h^{2m-2}) \end{aligned}$$


Circle  $C$ :  $x^2 + y^2 \leq h^2$ .

25.4.61




$$\frac{1}{\pi h^2} \iint_C f(x, y) dx dy = \sum_{i=1}^n w_i f(x_i, y_i) + R$$


$(x_i, y_i)$	$w_i$	
$(0, 0)$	$1/2$	$R = O(h^4)$
$(\pm h, 0), (0, \pm h)$	$1/8$	



$(x_i, y_i)$	$w_i$	
$(\pm \frac{h}{2}, \pm \frac{h}{2})$	$1/4$	$R = O(h^4)$




$(x_i, y_i)$	$w_i$	
$(0, 0)$	$1/2$	
$(\pm h, 0)$	$1/12$	$R = O(h^4)$
$(\pm \frac{h}{2}, \pm \frac{h}{2} \sqrt{3})$	$1/12$	






$(x_i, y_i)$	$w_i$
$(0, 0)$	$1/6$
$(\pm h, 0)$	$1/24$
$(0, \pm h)$	$1/24$
$(\pm \frac{h}{2}, \pm \frac{h}{2})$	$1/6$



$(x_i, y_i)$	$w_i$
$(0, 0)$	$1/4$
$(\pm \sqrt{\frac{2}{3}} h, 0)$	$1/8$
$(\pm \sqrt{\frac{1}{6}} h, \pm \frac{h}{2} \sqrt{2})$	$1/8$




$(x_i, y_i)$	$w_i$
$(0, 0)$	$1/9$
$(\sqrt{\frac{6-\sqrt{6}}{10}} h \cos \frac{2\pi k}{10}, \sqrt{\frac{6-\sqrt{6}}{10}} h \sin \frac{2\pi k}{10})$	$\frac{16+\sqrt{6}}{360}$
$(k=1, \dots, 10)$	
$(\sqrt{\frac{6+\sqrt{6}}{10}} h \cos \frac{2\pi k}{10}, \sqrt{\frac{6+\sqrt{6}}{10}} h \sin \frac{2\pi k}{10})$	$\frac{16-\sqrt{6}}{360}$
$R=O(h^{10})$	


Square<sup>4</sup>  $S: |x| \leq h, |y| \leq h$

25.4.62

$$\frac{1}{4h^2} \iint_S f(x, y) dx dy = \sum_{i=1}^n w_i f(x_i, y_i) + R$$




$(x_i, y_i)$	$w_i$
$(0, 0)$	$4/9$
$(\pm h, \pm h)$	$1/36$
$(\pm h, 0)$	$1/9$
$(0, \pm h)$	$1/9$



$(x_i, y_i)$	$w_i$
$(\pm h \sqrt{\frac{1}{3}}, \pm h \sqrt{\frac{1}{3}})$	$1/4$

$R=O(h^4)$



$(x_i, y_i)$	$w_i$
$(0, 0)$	$16/81$

<sup>4</sup> For regions, such as the square, cube, cylinder, etc., which are the Cartesian products of lower dimensional regions, one may always develop integration rules by "multiplying together" the lower dimensional rules. Thus if

$$\int_0^1 f(x) dx \approx \sum_{i=1}^n w_i f(x_i)$$

is a one dimensional rule, then

$$\int_0^1 \int_0^1 f(x, y) dx dy \approx \sum_{i,j=1}^n w_i w_j f(x_i, y_j)$$

becomes a two dimensional rule. Such rules are not necessarily the most "economical".


$$\begin{aligned} \left(\pm\sqrt{\frac{3}{5}}h, \pm\sqrt{\frac{3}{5}}h\right) & 25/324 \\ & R=O(h^6) \\ \left(0, \pm\sqrt{\frac{3}{5}}h\right) & 10/81 \\ \left(\pm\sqrt{\frac{3}{5}}h, 0\right) & 10/81 \end{aligned}$$

**Equilateral Triangle T**


Radius of Circumscribed Circle= $h$

**25.4.63**


$$\frac{1}{\frac{3}{4}\sqrt{3}h^2} \iint_T f(x,y) dx dy = \sum_{i=1}^n w_i f(x_i, y_i) + R$$



$(x_i, y_i)$	$w_i$	
$(0,0)$	$3/4$	$R=O(h^3)$
$(h,0)$	$1/12$	
$\left(-\frac{h}{2}, \pm\frac{h}{2}\sqrt{3}\right)$	$1/12$	



$(x_i, y_i)$	$w_i$	
$(0,0)$	$27/60$	$R=O(h^4)$
$(h,0)$	$3/60$	
$\left(-\frac{h}{2}, \pm\frac{h}{2}\sqrt{3}\right)$	$3/60$	
$\left(-\frac{h}{2}, 0\right)$	$8/60$	
$\left(\frac{h}{4}, \pm\frac{h}{4}\sqrt{3}\right)$	$8/60$	




$(x_i, y_i)$	$w_i$	
$(0,0)$	$270/1200$	$R=O(h^6)$
$\left(\left(\frac{\sqrt{15}+1}{7}\right)h, 0\right)$	$\frac{155-\sqrt{15}}{1200}$	
$\left(\left(\frac{-\sqrt{15}+1}{14}\right)h, \pm\left(\frac{\sqrt{15}+1}{14}\right)\sqrt{3}h\right)$		
$\left(\left(\frac{-\sqrt{15}-1}{7}\right)h, 0\right)$	$\frac{155+\sqrt{15}}{1200}$	
$\left(\left(\frac{\sqrt{15}-1}{14}\right)h, \pm\left(\frac{\sqrt{15}-1}{14}\right)\sqrt{3}h\right)$		

**Regular Hexagon H**

Radius of Circumscribed Circle= $h$


**25.4.64**

$$\frac{1}{\frac{3}{2}\sqrt{3}h^2} \iint_H f(x,y) dx dy = \sum_{i=1}^n w_i f(x_i, y_i) + R$$



$(x_i, y_i)$	$w_i$	
$(0,0)$	$21/36$	$R=O(h^4)$
$\left(\pm\frac{h}{2}, \pm\frac{h}{2}\sqrt{3}\right)$	$5/72$	
$(\pm h, 0)$	$5/72$	






$(x_i, y_i)$	$w_i$	
$(0, 0)$	258/1008	
$(\pm \frac{h}{10} \sqrt{14}, \pm \frac{h}{10} \sqrt{42})$	125/1008	$R = O(h^6)$
$(\pm h \frac{\sqrt{14}}{5}, 0)$	125/1008	


Surface of Sphere  $\Sigma$ :  $x^2 + y^2 + z^2 = h^2$

#### 25.4.65

$$\frac{1}{4\pi h^2} \int_{\Sigma} f(x, y, z) d\sigma = \sum_{i=1}^n w_i f(x_i, y_i, z_i) + R$$



$(x_i, y_i, z_i)$	$w_i$	
$(\pm h, 0, 0)$	1/6	$R = O(h^4)$
$(0, \pm h, 0)$	1/6	
$(0, 0, \pm h)$	1/6	




$(x_i, y_i, z_i)$	$w_i$
$(\pm \sqrt{\frac{1}{2}} h, \pm \sqrt{\frac{1}{2}} h, 0)$	1/15
$(\pm \sqrt{\frac{1}{2}} h, 0, \pm \sqrt{\frac{1}{2}} h)$	
$(0, \pm \sqrt{\frac{1}{2}} h, \pm \sqrt{\frac{1}{2}} h)$	
$(\pm h, 0, 0)$	$R = O(h^6)$
$(0, \pm h, 0)$	
$(0, 0, \pm h)$	

$(x_i, y_i, z_i)$	$w_i$
$(\pm \sqrt{\frac{1}{3}} h, \pm \sqrt{\frac{1}{3}} h, \pm \sqrt{\frac{1}{3}} h)$	27/840
$(\pm \sqrt{\frac{1}{2}} h, \pm \sqrt{\frac{1}{2}} h, 0)$	32/840 $R = O(h^8)$
$(\pm \sqrt{\frac{1}{2}} h, 0, \pm \sqrt{\frac{1}{2}} h)$	
$(0, \pm \sqrt{\frac{1}{2}} h, \pm \sqrt{\frac{1}{2}} h)$	
$(\pm h, 0, 0)$	40/840
$(0, \pm h, 0)$	
$(0, 0, \pm h)$	

Sphere  $S$ :  $x^2 + y^2 + z^2 \leq h^2$

#### 25.4.66


$$\frac{1}{\frac{4}{3}\pi h^3} \iiint_S f(x, y, z) dx dy dz = \sum_{i=1}^n w_i f(x_i, y_i, z_i) + R$$



$(x_i, y_i, z_i)$	$w_i$	
$(0, 0, 0)$	$2/5$	
$(\pm h, 0, 0)$	$1/10$	
		$R = O(h^4)$
$(0, \pm h, 0)$	$1/10$	
$(0, 0, \pm h)$	$1/10$	
Cube <sup>5</sup> $C$ : $ x  \leq h$		
		$ y  \leq h$
		$ z  \leq h$

25.4.67

$$\frac{1}{8h^3} \iiint_C f(x, y, z) dx dy dz = \sum_{i=1}^n w_i f(x_i, y_i, z_i) + R$$



$(x_i, y_i, z_i)$	$w_i$	
$(\pm h, 0, 0)$	$1/6$	
		$R = O(h^4)$
$(0, \pm h, 0)$	$1/6$	
$(0, 0, \pm h)$	$1/6$	

25.4.68

$$\begin{aligned} \frac{1}{8h^3} \iiint_C f(x, y, z) dx dy dz \\ = \frac{1}{360} [-496f_m + 128\sum f_r + 8\sum f_r + 5\sum f_e] + O(h^6) \end{aligned}$$

25.4.69

$$= \frac{1}{450} [91\sum f_r - 40\sum f_e + 16\sum f_d] + O(h^6)$$

 where  $f_m = f(0, 0, 0)$ .


<sup>5</sup> See footnote to 25.4.62.

 $\sum f_r$  = sum of values of  $f$  at the 6 points midway from the center of  $C$  to the 6 faces.

 $\sum f_r$  = sum of values of  $f$  at the 6 centers of the faces of  $C$ .

 $\sum f_v$  = sum of values of  $f$  at the 8 vertices of  $C$ .

 $\sum f_e$  = sum of values of  $f$  at the 12 midpoints of edges of  $C$ .

 $\sum f_d$  = sum of values of  $f$  at the 4 points on the diagonals of each face at a distance of  $\frac{1}{2}\sqrt{5}h$  from the center of the face.

 Tetrahedron:  $\mathcal{T}$ 

25.4.70

$$\begin{aligned} \frac{1}{V} \iiint_{\mathcal{T}} f(x, y, z) dx dy dz &= \frac{1}{40} \sum f_v + \frac{9}{40} \sum f_r \\ &\quad + \text{terms of 4th order} \\ &= \frac{32}{60} f_m + \frac{1}{60} \sum f_v + \frac{4}{60} \sum f_e \\ &\quad + \text{terms of 4th order} \end{aligned}$$

where

 $V$ : Volume of  $\mathcal{T}$ 
 $\sum f_v$ : Sum of values of the function at the vertices of  $\mathcal{T}$ .

 $\sum f_e$ : Sum of values of the function at midpoints of the edges of  $\mathcal{T}$ .

 $\sum f_r$ : Sum of values of the function at the center of gravity of the faces of  $\mathcal{T}$ .

 $f_m$ : Value of function at center of gravity of  $\mathcal{T}$ .



25.5. Ordinary Differential Equations<sup>6</sup>First Order:  $y' = f(x, y)$ 

Point Slope Formula

25.5.1 
$$y_{n+1} = y_n + hy'_n + O(h^2)$$

25.5.2 
$$y_{n+1} = y_{n-1} + 2hy'_n + O(h^3)$$

Trapezoidal Formula

25.5.3 
$$y_{n+1} = y_n + \frac{h}{2} (y'_{n+1} + y'_n) + O(h^3)$$

Adams' Extrapolation Formula

25.5.4 
$$y_{n+1} = y_n + \frac{h}{24} (55y'_n - 59y'_{n-1} + 37y'_{n-2} - 9y'_{n-3}) + O(h^5)$$

Adams' Interpolation Formula

25.5.5 
$$y_{n+1} = y_n + \frac{h}{24} (9y'_{n+1} + 19y'_n - 5y'_{n-1} + y'_{n-2}) + O(h^5)$$

## Runge-Kutta Methods

Second Order

25.5.6 
$$y_{n+1} = y_n + \frac{1}{2} (k_1 + k_2) + O(h^3)$$

$$k_1 = hf(x_n, y_n), k_2 = hf(x_n + h, y_n + k_1)$$

25.5.7 
$$y_{n+1} = y_n + k_2 + O(h^3)$$

$$k_1 = hf(x_n, y_n), k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1\right)$$

Third Order

25.5.8 
$$y_{n+1} = y_n + \frac{1}{6}k_1 + \frac{2}{3}k_2 + \frac{1}{6}k_3 + O(h^4)$$

$$k_1 = hf(x_n, y_n), k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1\right)$$

$$k_3 = hf(x_n + h, y_n - k_1 + 2k_2)$$

## 25.5.9

$$y_{n+1} = y_n + \frac{1}{4}k_1 + \frac{3}{4}k_3 + O(h^4)$$

$$k_1 = hf(x_n, y_n), k_2 = hf\left(x_n + \frac{1}{3}h, y_n + \frac{1}{3}k_1\right)$$

$$k_3 = hf\left(x_n + \frac{2}{3}h, y_n + \frac{2}{3}k_2\right)$$

Fourth Order

25.5.10 
$$y_{n+1} = y_n + \frac{1}{6}k_1 + \frac{1}{3}k_2 + \frac{1}{3}k_3 + \frac{1}{6}k_4 + O(h^5)$$

$$k_1 = hf(x_n, y_n), k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1\right)$$

$$k_3 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_2\right), k_4 = hf(x_n + h, y_n + k_3)$$

25.5.11 
$$y_{n+1} = y_n + \frac{1}{8}k_1 + \frac{3}{8}k_2 + \frac{3}{8}k_3 + \frac{1}{8}k_4 + O(h^5)$$

$$k_1 = hf(x_n, y_n), k_2 = hf\left(x_n + \frac{1}{3}h, y_n + \frac{1}{3}k_1\right)$$

$$k_3 = hf\left(x_n + \frac{2}{3}h, y_n - \frac{1}{3}k_1 + k_2\right),$$

$$k_4 = hf(x_n + h, y_n + k_1 - k_2 + k_3)$$

Gill's Method

25.5.12 
$$y_{n+1} = y_n + \frac{1}{6}\left(k_1 + 2\left(1 - \sqrt{\frac{1}{2}}\right)k_2\right.$$

$$\left. + 2\left(1 + \sqrt{\frac{1}{2}}\right)k_3 + k_4\right) + O(h^5)$$

$$k_1 = hf(x_n, y_n)$$

$$k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1\right)$$

$$k_3 = hf\left(x_n + \frac{1}{2}h, y_n + \left(-\frac{1}{2} + \sqrt{\frac{1}{2}}\right)k_1\right.$$

$$\left. + \left(1 - \sqrt{\frac{1}{2}}\right)k_2\right)$$

$$k_4 = hf\left(x_n + h, y_n - \sqrt{\frac{1}{2}}k_2 + \left(1 + \sqrt{\frac{1}{2}}\right)k_3\right)$$

## Predictor-Corrector Methods

Milne's Methods

25.5.13 
$$P: y_{n+1} = y_{n-3} + \frac{4h}{3}(2y'_n - y'_{n-1} + 2y'_{n-2}) + O(h^5)$$

$$C: y_{n+1} = y_{n-1} + \frac{h}{3}(y'_{n-1} + 4y'_n + y'_{n+1}) + O(h^5)$$

<sup>6</sup> The reader is cautioned against possible instabilities especially in formulas 25.5.2 and 25.5.13. See, e.g. [25.11], [25.12].

## 25.5.14

$$P: y_{n+1} = y_{n-5} + \frac{3h}{10} (11y'_n - 14y'_{n-1} + 26y'_{n-2} - 14y'_{n-3} + 11y'_{n-4}) + O(h^7)$$

$$C: y_{n+1} = y_{n-3} + \frac{2h}{45} (7y'_{n+1} + 32y'_n + 12y'_{n-1} + 32y'_{n-2} + 7y'_{n-3}) + O(h^7)$$

## Formulas Using Higher Derivatives

## 25.5.15

$$P: y_{n+1} = y_{n-2} + 3(y_n - y_{n-1}) + h^2(y''_n - y''_{n-1}) + O(h^5)$$

$$C: y_{n+1} = y_n + \frac{h}{2} (y'_{n+1} + y'_n) - \frac{h^2}{12} (y''_{n+1} - y''_n) + O(h^5)$$

## 25.5.16

$$P: y_{n+1} = y_{n-2} + 3(y_n - y_{n-1}) + \frac{h^3}{2} (y'''_n + y'''_{n-1}) + O(h^7)$$

$$C: y_{n+1} = y_n + \frac{h}{2} (y'_{n+1} + y'_n) - \frac{h^2}{10} (y''_{n+1} - y''_n) + \frac{h^3}{120} (y'''_{n+1} + y'''_n) + O(h^7)$$

## Systems of Differential Equations

First Order:  $y' = f(x, y, z)$ ,  $z' = g(x, y, z)$ .

## Second Order Runge-Kutta

## 25.5.17

$$y_{n+1} = y_n + \frac{1}{2} (k_1 + k_2) + O(h^3),$$

$$z_{n+1} = z_n + \frac{1}{2} (l_1 + l_2) + O(h^3)$$

$$k_1 = hf(x_n, y_n, z_n), \quad l_1 = hg(x_n, y_n, z_n)$$

$$k_2 = hf(x_n + h, y_n + k_1, z_n + l_1),$$

$$l_2 = hg(x_n + h, y_n + k_1, z_n + l_1)$$

## Fourth Order Runge-Kutta

## 25.5.18

$$y_{n+1} = y_n + \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4) + O(h^5),$$

$$z_{n+1} = z_n + \frac{1}{6} (l_1 + 2l_2 + 2l_3 + l_4) + O(h^5)$$

$$k_1 = hf(x_n, y_n, z_n) \quad l_1 = hg(x_n, y_n, z_n)$$

$$k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1, z_n + \frac{1}{2}l_1\right)$$

$$l_2 = hg\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_1, z_n + \frac{1}{2}l_1\right)$$

$$k_3 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_2, z_n + \frac{1}{2}l_2\right)$$

$$l_3 = hg\left(x_n + \frac{1}{2}h, y_n + \frac{1}{2}k_2, z_n + \frac{1}{2}l_2\right)$$

$$k_4 = hf(x_n + h, y_n + k_3, z_n + l_3)$$

$$l_4 = hg(x_n + h, y_n + k_3, z_n + l_3)$$

Second Order:  $y'' = f(x, y, y')$

## Milne's Method

## 25.5.19

$$P: y'_{n+1} = y'_{n-3} + \frac{4h}{3} (2y''_{n-2} - y''_{n-1} + 2y''_n) + O(h^5)$$

$$C: y'_{n+1} = y'_{n-1} + \frac{h}{3} (y''_{n-1} + 4y''_n + y''_{n+1}) + O(h^5)$$

## Runge-Kutta Method

## 25.5.20

$$y_{n+1} = y_n + h \left[ y'_n + \frac{1}{6} (k_1 + k_2 + k_3) \right] + O(h^5)$$

$$y'_{n+1} = y'_n + \frac{1}{6} (k_1 + 2k_2 + 2k_3 + k_4)$$

$$k_1 = hf(x_n, y_n, y'_n)$$

$$k_2 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{h}{2}y'_n + \frac{h}{8}k_1, y'_n + \frac{k_1}{2}\right)$$

$$k_3 = hf\left(x_n + \frac{1}{2}h, y_n + \frac{h}{2}y'_n + \frac{h}{8}k_1, y'_n + \frac{k_2}{2}\right)$$

$$k_4 = hf\left(x_n + h, y_n + hy'_n + \frac{h}{2}k_3, y'_n + k_3\right)$$

Second Order:  $y'' = f(x, y)$

## Milne's Method

## 25.5.21

$$P: y_{n+1} = y_n + y_{n-2} - y_{n-3}$$

$$+ \frac{h^2}{4} (5y''_n + 2y''_{n-1} + 5y''_{n-2}) + O(h^6)$$

$$C: y_n = 2y_{n-1} - y_{n-2} + \frac{h^2}{12} (y''_n + 10y''_{n-1} + y''_{n-2}) + O(h^6)$$

## Runge-Kutta Method

$$25.5.22 \quad y_{n+1} = y_n + h \left( y'_n + \frac{1}{6} (k_1 + 2k_2) \right) + O(h^4)$$

$$y'_{n+1} = y'_n + \frac{1}{6} k_1 + \frac{2}{3} k_2 + \frac{1}{6} k_3$$

$$k_1 = hf(x_n, y_n)$$

$$k_2 = hf\left(x_n + \frac{h}{2}, y_n + \frac{h}{2}y'_n + \frac{h}{8}k_1\right)$$

$$k_3 = hf\left(x_n + h, y_n + hy'_n + \frac{h}{2}k_2\right).$$



## References

## Texts

(For textbooks on numerical analysis, see texts in chapter 3)

- [25.1] J. Balbrecht and L. Collatz, Zur numerischen Auswertung mehrdimensionaler Integrale, *Z. Angew. Math. Mech.* **38**, 1–15 (1958).
- [25.2] Berthod-Zaborowski, Le calcul des intégrales de la forme:  $\int_0^1 f(x) \log x \, dx$ .  
H. Mineur, *Techniques de calcul numérique*, pp. 555–556 (Librairie Polytechnique Ch. Béranger, Paris, France, 1952).
- [25.3] W. G. Bickley, Formulae for numerical integration, *Math. Gaz.* **23**, 352 (1939).
- [25.4] W. G. Bickley, Formulae for numerical differentiation, *Math. Gaz.* **25**, 19–27 (1941).
- [25.5] W. G. Bickley, Finite difference formulae for the square lattice, *Quart. J. Mech. Appl. Math.*, **1**, 35–42 (1948).
- [25.6] G. Birkhoff and D. Young, Numerical quadrature of analytic and harmonic functions, *J. Math. Phys.* **29**, 217–221 (1950).
- [25.7] L. Fox, The use and construction of mathematical tables, *Mathematical Tables vol. I*, National Physical Laboratory (Her Majesty's Stationery Office, London, England, 1956).
- [25.8] S. Gill, Process for the step-by-step integration of differential equations in an automatic digital computing machine, *Proc. Cambridge Philos. Soc.* **47**, 96–108 (1951).
- [25.9] P. C. Hammer and A. H. Stroud, Numerical evaluation of multiple integrals II, *Math. Tables Aids Comp.* **12**, 272–280 (1958).
- [25.10] P. C. Hammer and A. W. Wymore, Numerical evaluation of multiple integrals I, *Math. Tables Aids Comp.* **11**, 59–67 (1957).
- [25.11] P. Henrici, *Discrete variable methods in ordinary differential equations* (John Wiley & Sons, Inc., New York, N. Y., 1961).
- [25.12] F. B. Hildebrand, *Introduction to numerical analysis* (McGraw-Hill Book Co., Inc., New York, N. Y., 1956).
- [25.13] Z. Kopal, *Numerical analysis* (John Wiley & Sons, Inc., New York, N. Y., 1955).
- [25.14] A. A. Markoff, *Differenzenrechnung* (B. G. Teubner, Leipzig, Germany, 1896).
- [25.15] S. E. Mikeladze, Quadrature formulas for a regular function, *Soobšč. Akad. Nauk Gruz. SSR.* **17**, 289–296 (1956).
- [25.16] W. E. Milne, A note on the numerical integration of differential equations, *J. Research NBS* **43**, 537–542 (1949) RP2046.
- [25.17] D. J. Panov, *Formelsammlung zur numerischen Behandlung partieller Differentialgleichungen nach dem Differenzenverfahren* (Akad. Verlag, Berlin, Germany, 1955).
- [25.18] E. Radau, Études sur les formules d'approximation qui servent à calculer la valeur d'une intégrale définie, *J. Math. Pures Appl.* (3) **6**, 283–336 (1880).

- [25.19] R. D. Richtmeyer, *Difference methods for initial-value problems* (Interscience Publishers, New York, N. Y., 1957).
- [25.20] M. Sadowsky, A formula for approximate computation of a triple integral, *Amer. Math. Monthly* **47**, 539–543 (1940).
- [25.21] H. E. Salzer, A new formula for inverse interpolation, *Bull. Amer. Math. Soc.* **50**, 513–516 (1944).
- [25.22] H. E. Salzer, Formulas for complex Cartesian interpolation of higher degree, *J. Math. Phys.* **28**, 200–203 (1949).
- [25.23] H. E. Salzer, Formulas for numerical integration of first and second order differential equations in the complex plane, *J. Math. Phys.* **29**, 207–216 (1950).
- [25.24] H. E. Salzer, Formulas for numerical differentiation in the complex plane, *J. Math. Phys.* **31**, 155–169 (1952).
- [25.25] A. Sard, Integral representations of remainders, *Duke Math. J.* **15**, 333–345 (1948).
- [25.26] A. Sard, Remainders: functions of several variables, *Acta Math.* **84**, 319–346 (1951).
- [25.27] G. Schulz, *Formelsammlung zur praktischen Mathematik* (DeGruyter and Co., Berlin, Germany, 1945).
- [25.28] A. H. Stroud, A bibliography on approximate integration, *Math. Comp.* **15**, 52–80 (1961).
- [25.29] G. J. Tranter, *Integral transforms in mathematical physics* (John Wiley & Sons, Inc., New York, N. Y., 1951).
- [25.30] G. W. Tyler, Numerical integration with several variables, *Canad. J. Math.* **5**, 393–412 (1953).

## Tables

- [25.31] L. J. Comrie, *Chambers' six-figure mathematical tables*, vol. 2 (W. R. Chambers, Ltd., London, England, 1949).
- [25.32] P. Davis and P. Rabinowitz, Abscissas and weights for Gaussian quadratures of high order, *J. Research NBS* **56**, 35–37 (1956) RP2645.
- [25.33] P. Davis and P. Rabinowitz, Additional abscissas and weights for Gaussian quadratures of high order: Values for  $n=64, 80$ , and  $96$ , *J. Research NBS* **60**, 613–614 (1958) RP2875.
- [25.34] E. W. Dijkstra and A. van Wijngaarden, *Table of Everett's interpolation coefficients* (Elcelsior's Photo-offset, The Hague, Holland, 1955).
- [25.35] H. Fishman, Numerical integration constants, *Math. Tables Aids Comp.* **11**, 1–9 (1957).
- [25.36] H. J. Gawlik, Zeros of Legendre polynomials of orders 2–64 and weight coefficients of Gauss quadrature formulae, *A.R.D.E. Memo (B) 77/58*, Fort Halstead, Kent, England (1958).
- [25.37] Gt. Britain H.M. Nautical Almanac Office, *Interpolation and allied tables* (Her Majesty's Stationery Office, London, England, 1956).



- [25.38] I. M. Longman, Tables for the rapid and accurate numerical evaluation of certain infinite integrals involving Bessel functions, *Math. Tables Aids Comp.* **11**, 166–180 (1957).
- [25.39] A. N. Lowan, N. Davids, and A. Levenson, Table of the zeros of the Legendre polynomials of order 1–16 and the weight coefficients for Gauss' mechanical quadrature formula, *Bull. Amer. Math. Soc.* **48**, 739–743 (1942).
- [25.40] National Bureau of Standards, Tables of Lagrangian interpolation coefficients (Columbia Univ. Press, New York, N.Y., 1944).
- [25.41] National Bureau of Standards, Collected Short Tables of the Computation Laboratory, Tables of functions and of zeros of functions, *Applied Math. Series 37* (U.S. Government Printing Office, Washington, D.C., 1954).
- [25.42] P. Rabinowitz, Abscissas and weights for Lobatto quadrature of high order, *Math. Tables Aids Comp.* **69**, 47–52 (1960).
- [25.43] P. Rabinowitz and G. Weiss, Tables of abscissas and weights for numerical evaluation of integrals of the form

$$\int_0^{\infty} e^{-x} x^n f(x) dx,$$

*Math. Tables Aids Comp.* **68**, 285–294 (1959).

- [25.44] H. E. Salzer, Tables for facilitating the use of Chebyshev's quadrature formula, *J. Math. Phys.* **26**, 191–194 (1947).
- [25.45] H. E. Salzer and R. Zucker, Table of the zeros and weight factors of the first fifteen Laguerre polynomials, *Bull. Amer. Math. Soc.* **55**, 1004–1012 (1949).
- [25.46] H. E. Salzer, R. Zucker, and R. Capuano, Table of the zeros and weight factors of the first twenty Hermite polynomials, *J. Research NBS* **48**, 111–116 (1952) RP2294.
- [25.47] H. E. Salzer, Table of coefficients for obtaining the first derivative without differences, *NBS Applied Math. Series 2* (U.S. Government Printing Office, Washington, D.C., 1948).
- [25.48] H. E. Salzer, Coefficients for facilitating trigonometric interpolation, *J. Math. Phys.* **27**, 274–278 (1949).
- [25.49] H. E. Salzer and P. T. Roberson, Table of coefficients for obtaining the second derivative without differences, *Convair-Astronautics*, San Diego, Calif. (1957).
- [25.50] H. E. Salzer, Tables of osculatory interpolation coefficients, *NBS Applied Math. Series 56* (U.S. Government Printing Office, Washington, D.C., 1958).