

ATLAS Muon Barrel Level-1 trigger performance in 2017 data


Introduction

The ATLAS experiment utilises the Resistive Plate Chambers detector (RPC) for the first level muon trigger system in the barrel region of the detector. This poster presents measurements of RPC detector and trigger performance using proton-proton collisions at a centre-of-mass energy of 13 TeV collected in 2017, showing results in terms of the trigger timing and efficiency.

The ATLAS RPC Detector and Trigger System


The present ATLAS muon trigger in the barrel region based on

- 3 concentric RPC layers
- 16 physical sectors
- each physical sector is segmented in 2 trigger sectors
- totally 64 trigger sectors in side A and side C
- each trigger sector is segmented along η in towers




The Level-1 (L1) trigger algorithm is based on hit coincidence of 3 concentric RPC stations

- Low p_T trigger: coincidence between the innermost two RPC stations
- High p_T trigger: additional confirmation on the third external station




Trigger efficiency vs. offline muon η




Efficiency of L1 MU10 and MU11 trigger in 2017 including (in green) or excluding (yellow) the newly commissioned trigger chambers in the “feet” region of the ATLAS Muon Spectrometer as a function of η of the offline muon candidates in the barrel detector region, for sector 12 and 14 of the “feet” region.

Trigger efficiency vs. offline muon ϕ




Efficiency of L1 MU10 and MU11 trigger in 2017 and comparison with 2016 trigger efficiency. The efficiency is plotted as a function of ϕ of offline muon candidates in the barrel detector region.

Trigger timing performance



Fraction of RPC trigger hits associated to the correct BC for each of the barrel muon trigger towers for one example run in 2017.




Fraction for the whole RPC trigger system as a function of time. Each point corresponds to a different ATLAS run recorded in 2017. The fraction of high p_T muons associated to the correct BC is 99.6%.

Trigger efficiency vs. offline muon p_T


L1 muon barrel trigger efficiency for reconstructed muons as a function of transverse momentum is shown for 6 L1 thresholds.

- efficiencies for the low p_T trigger thresholds (MU4, MU6, MU10) reach a plateau of about 80%
- efficiencies for high p_T trigger thresholds (MU11, MU20, MU21) rise to around 70%




Plateau value of the trigger efficiency as a function of time.

- each point corresponds to a different ATLAS run recorded in 2017
- only runs with integrated luminosity greater than 50 pb⁻¹ and at least 1000 muons collected are used



Ratio of trigger efficiencies (2017/2016)



η - ϕ map of the ratio between the Level-1 Barrel muon trigger efficiency in 2017 and 2016 for the trigger threshold MU10 and MU11 are shown.

- slight decrease in trigger efficiency in 2017 mainly due to gas leaks \rightarrow HV disconnected.
- efficiency gained in the feet sectors credited with re-cabling of the high p_T feet trigger performed in the 2016-2017 shutdown
- improvement in sector 13 where the BME chambers work now