Centre No.			Pape	er Refer	rence			Surname		Initia	l(s)
Candidate No.		6 6	7	7	/	0	1	Signature		•	
	Paper Reference(s) 6677/01				•		•	,	Exan	niner's use	e only
			. ~								
	Edex	cel G	C	\mathbb{E}					Team I	Leader's u	se only
	Mechai	nics M	1								
				.	C	ha:	di.				
	Advance									Question Number	Leave Blank
	Tuesday		•			Mo	rnın	ıg		1	
	Time: 1 h	our 30 1	min	utes						2	
										3	
										4	
	Materials requirements Mathematical Fo			- Ito		cluded	l with	question papers		5	
		`					1			6	
		lifications. C	alculat	ors m	ust not	have	the fa	cility for symbolic		7	
	algebra manipu mathematical fo				integr	ation,	or ha	ve retrievable		,	
I	to Condidates										
	to Candidates ove, write your centre n	umber, cand	idate r	numbe	r, youi	surna	ame, i	nitials and signatu	re.		
Check that you Answer ALL to	have the correct questi	on paper.									
You must writ	e your answer for each of										
	ore space to complete your merical value of g is re					e addi	itional	sheets.			
	ator is used, the answer					ate de	gree (of accuracy.			
				-	-			-			
	for Candidates										
A booklet 'Ma	thematical Formulae and	d Statistical	[ables]	is pro	ovided						

This publication may be reproduced only in accordance with Edexcel Limited copyright policy. ©2009 Edexcel Limited.

Full marks may be obtained for answers to ALL questions.

Answers without working may not gain full credit.

You must ensure that your answers to parts of questions are clearly labelled. You must show sufficient working to make your methods clear to the Examiner.



Advice to Candidates



The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2). There are 7 questions in this question paper. The total mark for this paper is 75. There are 24 pages in this question paper. Any blank pages are indicated.



Turn over

Total

Find the value of u .	
Find the value of u .	(5)
	(-)

Question 1 continued	Leave blank
	_
	-
	-
	_
	_
	-
	-
	_
	_
	-
	-
	_
	_
	-
	-
	-
	_
	-
	-
	_
	_
	-
	-
	_
	_
	-
	Q1
(Total 5 marks	
(2002 0 1100210	,

2.	A small ball is projected vertically upwards from ground level with speed u m s ⁻¹ . The takes 4 s to return to ground level.	ball
	(a) Draw, in the space below, a velocity-time graph to represent the motion of the during the first 4 s.	ball (2)
	(b) The maximum height of the ball above the ground during the first 4 s is 19.6 m. F	
	the value of u .	(3)
		_
		_
		_
		_
		_
		_

Question 2 continued	Leave blank
	02
	Q2
(Total 5 marks)	

3.	Two particles A and B are moving on a smooth horizontal plane. The mass of A is km , where $2 < k < 3$, and the mass of B is m . The particles are moving along the same straight line, but in opposite directions, and they collide directly. Immediately before they collide the speed of A is $2u$ and the speed of B is $4u$. As a result of the collision the speed of A is halved and its direction of motion is reversed.
	(a) Find, in terms of k and u , the speed of B immediately after the collision. (3)
	(b) State whether the direction of motion of B changes as a result of the collision, explaining your answer. (3)
	Given that $k = \frac{7}{3}$,
	(c) find, in terms of m and u , the magnitude of the impulse that A exerts on B in the collision.
	(3)

Question 3 continued	Leave blank
	Q3
(Total 9 marks)	

Leave blank

4.




Figure 1

A bench consists of a plank which is resting in a horizontal position on two thin vertical legs. The plank is modelled as a uniform rod PS of length 2.4 m and mass 20 kg. The legs at Q and R are 0.4 m from each end of the plank, as shown in Figure 1.

Two pupils, Arthur and Beatrice, sit on the plank. Arthur has mass 60 kg and sits at the middle of the plank and Beatrice has mass 40 kg and sits at the end *P*. The plank remains horizontal and in equilibrium. By modelling the pupils as particles, find

(a) the magnitude of the normal reaction between the plank and the leg at Q and the magnitude of the normal reaction between the plank and the leg at R.

(7)

Beatrice stays sitting at P but Arthur now moves and sits on the plank at the point X. Given that the plank remains horizontal and in equilibrium, and that the magnitude of the normal reaction between the plank and the leg at Q is now twice the magnitude of the normal reaction between the plank and the leg at R,

(b) find the distance QX.

(6)

	Leave blank
Question 4 continued	Diank

Question 4 continued	b

Question 4 continued	Leave
	Q4
(Total 13	marks)

5.

Leave blank




Figure 2

A small package of mass 1.1 kg is held in equilibrium on a rough plane by a horizontal force. The plane is inclined at an angle α to the horizontal, where $\tan \alpha = \frac{3}{4}$. The force sets in a vertical plane containing a line of greatest slope of the plane and has magnitude

acts in a vertical plane containing a line of greatest slope of the plane and has magnitude P newtons, as shown in Figure 2.

The coefficient of friction between the package and the plane is 0.5 and the package is modelled as a particle. The package is in equilibrium and on the point of slipping down the plane.

- (a) Draw, on Figure 2, all the forces acting on the package, showing their directions clearly. (2)
- (b) (i) Find the magnitude of the normal reaction between the package and the plane.

(ii) Find	l the	value	of P .
-----------	-------	-------	----------

(11)



	Leave blank
Question 5 continued	

uestion 5 continued	 	

Question 5 continued		Lea bla
		Q
	(Total 13 marks)	

_
Leave
hlank

6.	Two forces, $(4\mathbf{i} - 5\mathbf{j})$ N and $(p\mathbf{i} + q\mathbf{j})$ N, act on a particle P of mass m kg. The result of the two forces is R . Given that R acts in a direction which is parallel to vector $(\mathbf{i} - 2\mathbf{j})$,	tant the
	(a) find the angle between \mathbf{R} and the vector \mathbf{j} ,	(2)
		(3)
	(b) show that $2p + q + 3 = 0$.	
		(4)
	Given also that $q = 1$ and that P moves with an acceleration of magnitude $8\sqrt{5}$ m s ⁻² ,	
	(c) find the value of m .	(-)
		(7)

Question 6 continued	Leave blank

Question 6 continued	

Question 6 continued		Leav blank
		Q
	(Total 14 marks)	

7.

Leave blank




Figure 3

One end of a light inextensible string is attached to a block P of mass 5 kg. The block P is held at rest on a smooth fixed plane which is inclined to the horizontal at an angle α ,

where $\sin \alpha = \frac{3}{5}$. The string lies along a line of greatest slope of the plane and passes over

a smooth light pulley which is fixed at the top of the plane. The other end of the string is attached to a light scale pan which carries two blocks Q and R, with block Q on top of block R, as shown in Figure 3. The mass of block Q is 5 kg and the mass of block R is 10 kg. The scale pan hangs at rest and the system is released from rest. By modelling the blocks as particles, ignoring air resistance and assuming the motion is uninterrupted, find

- (a) (i) the acceleration of the scale pan,
 - (ii) the tension in the string,

(8)

(b) the magnitude of the force exerted on block Q by block R,

(3)

(c) the magnitude of the force exerted on the pulley by the string.

(5)



Question 7 continued	Leave



END	TOTAL FUN FAFEN; /5 MIARNS	
	(Total 16 marks) TOTAL FOR PAPER: 75 MARKS	
		Q



