

Edexcel GCE

# Mathematics

Statistics S2 6684

Summer 2005

advancing learning, changing lives

Final Mark Scheme

Edexcel GCE

# Mathematics

## General Instructions


1. The total number of marks for the paper is 75.
2. Method (M) marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
3. Accuracy (A) marks can only be awarded if the relevant method (M) marks have been earned.
4. (B) marks are independent of method marks.
5. Method marks should not be subdivided.
6. For misreading which does not alter the character of a question or materially simplify it, deduct two from any A or B marks gained, in that part of the question affected. Indicate this action by 'MR' in the body of the script (but see also note 10).
7. If a candidate makes more than one attempt at any question:
  - (a) If all but one attempt is crossed out, mark the attempt which is NOT crossed out.
  - (b) If either all attempts are crossed out or none are crossed out, mark all the attempts and score the highest single attempt.
8. Marks for each question, or part of a question, must appear in the right-hand margin and, in addition, total marks for each question, even where zero, must be ringed and appear in the right-hand margin and on the grid on the front of the answer book. It is important that a check is made to ensure that the totals in the right-hand margin of the ringed marks and of the unringed marks are equal. The total mark for the paper must be put on the top right-hand corner of the front cover of the answer book.
9. For methods of solution not in the mark scheme, allocate the available M and A marks in as closely equivalent a way as possible, and indicate this by the letters 'OS' (outside scheme) put alongside in the body of the script.
10. All A marks are 'correct answer only' (c.a.o.) unless shown, for example, as A1 f.t. to indicate that previous wrong working is to be followed through. In the body of the script the symbol  $\checkmark$  should be used for correct f.t. and  $\nabla$  for incorrect f.t. After a misread, however, the subsequent A marks affected are treated as A f.t., but manifestly absurd answers should never be awarded A marks.
11. Ignore wrong working or incorrect statements following a correct answer.

June 2005  
6684 Statistics S2  
Mark Scheme

Question Number	Scheme	Marks
1(a)	$X \sim B(n, 0.04)$ $E(X) = np$ $5 = 0.04n$ $n = 125$	Implied B1 Use of $np = 5$ M1 125 A1 (3)
(b)	$E(X) = 3$ $np = 3$ $sd = \sqrt{npq} = \sqrt{3(1-0.04)}$ $= \sqrt{2.88}$ $= 1.70$	$np = 3$ B1 Use of $npq$ M1 $\sqrt{3(1-0.04)}$ A1 awrt 1.70 A1 (4)
		<b>Total 7</b>
2(a)	$f(x) = \frac{1}{4}, 2 \leq x \leq 6$ $= 0$ , otherwise	$\frac{1}{4}$ and range B1 $0$ and range B1 (2)
(b)	$E(X) = 4$ by symmetry or formula	4 B1 (1)
(c)	$\text{Var}(X) = \frac{(6-2)^2}{12}$ $= \frac{4}{3}$	Use of formula M1 $1.\dot{3}$ or $1\frac{1}{3}$ or $\frac{4}{3}$ or 1.33 A1 (2)
(d)	$F(x) = \int_2^x \frac{1}{4} dt = \left[ \frac{1}{4}t \right]_2^x$ $= \frac{1}{4}(x-2)$ $F(x) = \frac{1}{4}(x-2), 2 \leq x \leq 6$ $= 1, x > 6$ $= 0, x < 2$	Use of $\int f(x) dx$ M1 $\frac{1}{4}(x-2)$ or equiv. A1 $\frac{1}{4}(x-2)$ and range B1ft ends and ranges B1 (4)
(e)	$P(2.3 < X < 3.4) = \frac{1}{4}(3.4 - 2.3)$ $= 0.275$	Use of area or $F(x)$ M1 $0.275$ or $\frac{11}{40}$ A1 (2)
		<b>Total 11</b>

Question Number	Scheme	Marks
3(a)	Misprints are random / independent, occur singly in space and at a constant rate	Context, any 2 B1, B1 (2)
(b)	$P(X = 0) = e^{-2.5}$ $= 0.08208..... = 0.0821$	Po (2.5) 0.0821 M1 A1 (2)
(c)	$Y \sim \text{Po}(5)$ for 2 pages $P(Y > 7) = 1 - P(X \leq 7)$ $= 1 - 0.8666 = 0.1334$	Implied Use of 1 – and correct inequality 0.1334 B1 M1 A1 (3)
(d)	For 20 pages, $Y \sim P_o(50)$ $Y \sim N(50, 50)$ approx $P(Y < 40) = P(Y \leq 39.5)$ $= P\left(Z \leq \frac{39.5 - 50}{\sqrt{50}}\right)$ $= P(Z \leq -1.4849)$ $= 1 - 0.93 = 0.07$	$P_o(50)$ $N(50, 50)$ cc $\pm 0.5$ standardise above all correct awrt – 1.48 0.07 B1 B1 M1 M1 A1 A1 A1 (7)
		<b>Total 14</b>
4(a)	Individual member or element of the population or sampling frame	B1 (1)
(b)	A <u>list</u> of <u>all</u> sampling units or <u>all</u> the population	B1 (1)
(c)	<u>All</u> possible <u>samples</u> are chosen from a population; the <u>values</u> of a <u>statistic</u> and the associated <u>probabilities</u> is a sampling distribution	B1 B1 (2)
		<b>Total 4</b>

Question Number	Scheme	Marks
5(a)	$X \sim B(200, 0.02)$ <u>n large, P small</u> so $X \sim \text{Po}(np) = \text{Po}(4)$ $P(X = 5) = \frac{e^{-4} 4^5}{5!} = 0.1563$	Implied conditions, $P_0(4)$ $P(X \leq 5) - P(X \leq 4)$ 0.1563 M1 A1 (5)
(b)	$P(X < 5) = P(X \leq 4)$ $= 0.6288$	$P(X \leq 4)$ 0.6288 M1 A1 (2) <b>Total 7</b>
6(a)	$\int_0^2 k(4x - x^3) dx = 1$ $k \left[ 2x^2 - \frac{1}{4}x^4 \right]_0^2 = 1$ $k(8 - 4) = 1$ $k = \frac{1}{4}$	$\int f(x) dx = 1$ , all correct M1 A1 [*] A1 cso A1 (4)
(b)	$E(X) = \int_0^2 x \cdot \frac{1}{4}(4x - x^3) dx$ $= \left[ \frac{1}{3}x^3 - \frac{1}{20}x^5 \right]_0^2$ $= \frac{16}{15}$	$\int xf(x) dx$ M1 [*] A1 $1.07$ or $1\frac{1}{15}$ or $\frac{16}{15}$ or $1.0\dot{6}$ A1 (3)
(c)	At mode, $f'(x) = 0$ $4 - 3x^2 = 0$ $x = \frac{2}{\sqrt{3}}$	Implied Attempt to differentiate $\sqrt{\frac{4}{3}}$ or 1.15 or $\frac{2}{\sqrt{3}}$ or $\frac{2\sqrt{3}}{3}$ M1 M1 A1 (3)
(d)	At median, $\int_0^x \frac{1}{4}(4t - t^3) dt = \frac{1}{2}$ $\frac{1}{4} \left( 2x^2 - \frac{1}{4}x^4 \right) = \frac{1}{2}$ $x^4 - 8x^2 + 8 = 0$ $x^2 = 4 \pm 2\sqrt{2}$ $x = 1.08$	$F(x) = \frac{1}{2}$ or $\int f(x) dx = \frac{1}{2}$ M1 Attempt to integrate M1 Attempt to solve quadratic Awrt 1.08 M1 A1 (4)

(e)	mean (1.07) < median (1.08) < mode (1.15) ⇒ negative skew	any pair cao	M1 A1 (2)
(f)		lines $x < 0$ and $x > 2$ , labels, 0 and 2  negative skew between 0 and 2	B1  B1 (2)  <b>Total 18</b>
7 (a)	$X \sim B(10, p)$	Binomial (10, 0.75)	B1, B1 (2)
(b)	$P(X = 6) = 0.9219 - 0.7759$ $= 0.1460$	$P(X \leq 6) - P(X \leq 5)$ 0.1460	M1 A1 (2)
(c)	$H_0: p = 0.75$ (or $p = 0.25$ ) $H_1: p < 0.75$ (or $p > 0.25$ ) Under $H_0$ , $X \sim B(20, 0.75)$ (or $Y \sim B(20, 0.25)$ )	Correct $H_0$ One tailed $H_1$ Implied	B1 B1 B1
	$P(X \leq 13) = 1 - 0.7858 = 0.2142$ (or $P(Y \geq 7)$ ) Insufficient evidence to reject $H_0$ as $0.2412 > 0.05$ Doctor's belief is not supported by the sample	$P(X \leq 13)$ and $1 -$ , 0.2142	M1, A1  Context A1  (6)
	<i>(OR CR <math>P(X \leq 12) = 1 - 0.8982 = 0.1018</math>            (or <math>P(Y \geq 8)</math>)  <math>P(X \leq 11) = 1 - 0.9591 = 0.0409</math>            (or <math>P(Y \geq 9)</math>)            13 outside critical region            (or 7))</i>	<i>either</i>	<i>(M1 A1)</i>
(d)	$P(X \leq c) \leq 0.01$ for $p=0.75$ (or $P(Y \geq 20-c) \leq 0.01$ for $p=0.25$ ) $P(X \leq 9) = 1 - 0.9961 = 0.0039$ (or $P(Y \geq 11)$ ) $P(X \leq 10) = 1 - 0.9861 = 0.0139$ (or $P(Y \geq 10)$ ) C. R. is [0,9], so greatest no. of patients is 9.	0.9961 or 0.9981 9	M1 A1 B1 B1 (4) <b>Total 14</b>