

Mark Scheme (Results) January 2011

GCE

GCE Mechanics M1 (6677) Paper 1



Edexcel is one of the leading examining and awarding bodies in the UK and throughout the world. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers.

Through a network of UK and overseas offices, Edexcel's centres receive the support they need to help them deliver their education and training programmes to learners.

For further information, please call our GCE line on 0844 576 0025, our GCSE team on 0844 576 0027, or visit our website at www.edexcel.com.

If you have any subject specific questions about the content of this Mark Scheme that require the help of a subject specialist, you may find our Ask The Expert email service helpful.

Ask The Expert can be accessed online at the following link:

http://www.edexcel.com/Aboutus/contact-us/

January 2011
Publications Code UA026577
All the material in this publication is copyright
© Edexcel Ltd 2011

General Instructions for Marking

- 1. The total number of marks for the paper is 75.
- 2. The Edexcel Mathematics mark schemes use the following types of marks:
 - M marks: method marks are awarded for 'knowing a method and attempting to apply it', unless otherwise indicated.
 - A marks: Accuracy marks can only be awarded if the relevant method (M) marks have been earned.
 - B marks are unconditional accuracy marks (independent of M marks)
 - Marks should not be subdivided.

3. Abbreviations

These are some of the traditional marking abbreviations that will appear in the mark schemes.

- bod benefit of doubt
- ft follow through
- the symbol √will be used for correct ft
- cao correct answer only
- cso correct solution only. There must be no errors in this part of the question to obtain this mark
- isw ignore subsequent working
- awrt answers which round to
- SC: special case
- oe or equivalent (and appropriate)
- · dep dependent
- indep independent
- dp decimal places
- sf significant figures
- * The answer is printed on the paper
- The second mark is dependent on gaining the first mark

January 2011 Mechanics M1 6677 Mark Scheme

| Question Number | Scheme | Marks | |
|--------------------|---|-------------|----|
| 1. (a) | Conservation of momentum: 4m-6=m+9 m=5 | M1 A1 A1 | 3) |
| (b) | Impulse = change in momentum = $3 \times 3 - (3 \times -2) = 15$ | M1 A1 (2 | |

| Question Number | Scheme | Marks | |
|--------------------|---|-------------|--------------------|
| 2. (a) | $-6.45 = u - 9.8 \times 0.75$ $0.9 = u **$ | M1 A1 A1 | (3) |
| (b) | $0 = 0.81 - 2 \times 9.8 \times s$ s = 0.041 or 0.0413 | M1 A1 | (2) |
| (c) | $h = -0.9 \times 0.75 + 4.9 \times 0.75^{2}$ $h = 2.1 \text{ or } 2.08$ | M1 A1 | |
| | | | (3) [8] |

| Question Number | Scheme | Marks |
|--------------------|--|-------------------|
| 3. (a) | $ \begin{array}{cccccccccccccccccccccccccccccccccccc$ | M1.0.1 |
| | Taking moments about B: $5 \times R_C = 20g \times 3$ $R_C = 12g \text{ or } 60g/5 \text{ or } 118 \text{ or } 120$ | M1A1 A1 |
| | Resolving vertically: $R_C + R_B = 20g$ $R_B = 8g \text{ or } 78.4 \text{ or } 78$ | M1 A1 (5) |
| (b) | $ \begin{array}{cccccccccccccccccccccccccccccccccccc$ | |
| | Resolving vertically: $50g = R + R$ | B1 |
| | Taking moments about B: $5 \times 25g = 3 \times 20g + (6-x) \times 30g$ | M1 A1 A1 |
| | 30x = 115 x = 3.8 or better or 23/6 oe | A1 (5) [10] |

| Question Number | Scheme | Marks |
|--------------------|---|----------|
| 4. | <u></u> | |
| (a) | speed = $\sqrt{2^2 + (-5)^2}$ | M1 |
| | $=\sqrt{29}=5.4$ or better | A1 |
| | | (2) |
| (b) | ((7i+10j)-(2i-5j))/5 | M1 A1 |
| | $ \frac{\left(\left(7\mathbf{i} + 10\mathbf{j}\right) - \left(2\mathbf{i} - 5\mathbf{j}\right)\right)/5}{=\left(5\mathbf{i} + 15\mathbf{j}\right)/5 = \mathbf{i} + 3\mathbf{j}} $ | A1 |
| | $\mathbf{F} = m\mathbf{a} = 2(\mathbf{i} + 3\mathbf{j}) = 2\mathbf{i} + 6\mathbf{j}$ | DM1 A1ft |
| | | (5) |
| | | |
| (c) | $\mathbf{v} = \mathbf{u} + \mathbf{a}t = (2\mathbf{i} - 5\mathbf{j}) + (\mathbf{i} + 3\mathbf{j})t$ | M1 |
| | $\left(-5+3t\right)$ j | A1 |
| | | |
| | Parallel to $i \Rightarrow -5 + 3t = 0$ | M1 |
| | t = 5/3 | A1 |
| | | (4) |
| | | [11] |

| Question Number | Scheme | Marks |
|--------------------|--|--------------------------|
| 5. (a) (i) | 1st section correct 2nd & 3rd sections correct Numbers and v marked correctly on the axes. | B1 B1 DB1 |
| (ii) | 1 st section correct 2 nd section correct 3 rd section correct and no "extras" on the sketch | B1 B1 B1 (6) |
| (b) | $\frac{70+40}{2} \times v = 880$ $v = 880 \times \frac{2}{110} = 16$ | M1 A1 DM1 A1 (4) [10] |

| Question Number | Scheme | Marks |
|--------------------|---|--|
| 6. (a) | 30 N F 120 N | |
| | Resolving perpendicular to the plane: $S = 120\cos\alpha + 30\sin\alpha$ = 114 * | M1 A1 A1 A1 (4) |
| (b) | R P F $120 N$ | |
| | Resolving perpendicular to the plane: $R = 120 \cos \alpha$ $= 96$ $F_{\text{max}} = \frac{1}{2}R$ Resolving parallel to the plane: In equilibrium: $P_{\text{max}} = F_{\text{max}} + 120 \sin \alpha$ $= 48 + 72 = 120$ | M1 A1 A1 M1 M1 A(2,1,0) A1 |
| (c) | $30+F=120\sin\alpha$ OR $30-F=120\sin\alpha$ So $F=42N$ acting up the plane. | (8) M1 A1 A1 (3) [15] |

| Question Number | Scheme | Marks |
|--------------------|---|--------------------------------------|
| 7. (a) | $ \begin{array}{cccc} P & T & B & \tan \theta = \frac{5}{12} \\ \hline A & 7 & \text{kg} & \cos \theta = \frac{12}{13} \end{array} $ | |
| | For A: $7g - T = 7a$ For B: parallel to plane $T - F - 3g \sin \theta = 3a$ perpendicular to plane $R = 3g \cos \theta$ $F = \mu R = 3g \cos \theta = 2g \cos \theta$ Eliminating T , $7g - F - 3g \sin \theta = 10a$ Equation in g and a: $7g - 2g \times \frac{12}{13} - 3g \frac{5}{13} = 7g - \frac{39}{13}g = 4g = 10a$ | M1 A1 M1 A1 M1 A1 M1 DM1 |
| | $a = \frac{2g}{5}$ oe or 3.9 or 3.92 | A1 (10) |
| (b) | After 1 m, $v^2 = u^2 + 2as$, $v^2 = 0 + 2 \times \frac{2g}{5} \times 1$ v = 2.8 | M1 A1 (2) |
| (c) | $-(F+3g \sin \theta) = 3a$ $\frac{2}{3} \times 3g \times \frac{12}{13} + 3g \times \frac{5}{13} = 3g = -3a, \ a = -g$ $v = u + at, \ 0 = 2.8 - 9.8t,$ $t = \frac{2}{7} \text{ oe, } 0.29. \ 0.286$ | M1 A1 DM1 A1 (4) [16] |

Further copies of this publication are available from Edexcel Publications, Adamsway, Mansfield, Notts, NG18 4FN

Telephone 01623 467467 Fax 01623 450481 Email <u>publications@linneydirect.com</u> Order Code UA026577 January 2011

For more information on Edexcel qualifications, please visit www.edexcel.com/quals

Edexcel Limited. Registered in England and Wales no.4496750 Registered Office: One90 High Holborn, London, WC1V 7BH