

Question number	Scheme	Marks
1.		
(a)	Take a (simple) random sample from (mutually exclusive) groups of the population Sample sizes within strata in strict proportion to numbers in each strata in the population <b>Advantage:</b> More accurate estimate of variance of population mean Individual estimates for strata available <b>Disadvantage:</b> Difficult if strata are large Definition of strata problematic (may overlap)	1g/1h B1 B1 Any one B1 Any one B1
(b)	Non-random sampling from groups of the population <b>Advantage:</b> Representative sample can be achieved with small sample size Cheap (costs kept to a minimum) Administration relatively easy <b>Disadvantage:</b> Not possible to estimate sampling errors due to lack of randomness Judgment of interviewer can affect choice of sample – bias OK Non-response not recorded Difficulties of defining controls e.g. social class	B1 B1dep Any one (not quick) B1 Any one B1
		(4) (4)
		8
2.		
(a)	$X \sim N(124, 20^2)$ or $\bar{X} \sim (124, \frac{20^2}{30})$ or assume $\sigma^2$ estimated by $s^2$ or CLT, vals. $\bar{x} \pm 2.5758 \frac{\sigma}{\sqrt{n}} = 124 \pm 2.5758 \frac{20}{\sqrt{30}}$ 2.5758, formula + attempt, all correct & 2.58, 2.576 $= 124 \pm 9.405$ $= (115, 133)$	B1,B1 B1M1A1 3 sf A1
(b)	140 is not in confidence interval Underweight apples chosen or Sample may not be representative/may be biased	M1 Any one A1J
		(6) (2)
		8

Question number	Scheme	Marks																
3.																		
(a)	$E(X-Y)=20-10=10$	Require minus, 10 <b>M1A1</b>																
(b)	$Var(X-Y)=5+4=9$	Require plus, 9 <b>M1A1</b>																
(c)	$X-Y \sim N(10,9)$ $P(13 < X-Y < 16)=P(X-Y < 16)-P(X-Y < 13)$ $= P(Z < \frac{16-10}{3}) - P(Z < \frac{13-10}{3})$ $= P(Z < 2) - P(Z < 1)$ $= 0.9772 - 0.8413 = 0.1359$	Implied <b>B1f</b> Subtract <b>M1</b> Standardise <b>M1</b> 2&1 <b>A1</b> 0.1359 <b>A1</b>																
		(2) (2) (5)																
		9																
4.	$H_0$ : Taking drug and catching a cold are independent (not associated) $H_1$ : Taking drug and catching a cold are not independent ( associated) (not ditto)	<b>B1</b> both <b>B1</b>																
	<table><tr><td></td><td>Cold</td><td>Not Cold</td><td></td></tr><tr><td>Drug</td><td>34 (39.5)</td><td>66 (60.5)</td><td>100</td></tr><tr><td>Dummy</td><td>45 (39.5)</td><td>55 (60.5)</td><td>100</td></tr><tr><td></td><td>79</td><td>121</td><td>200</td></tr></table>		Cold	Not Cold		Drug	34 (39.5)	66 (60.5)	100	Dummy	45 (39.5)	55 (60.5)	100		79	121	200	All totals <b>B1</b>
	Cold	Not Cold																
Drug	34 (39.5)	66 (60.5)	100															
Dummy	45 (39.5)	55 (60.5)	100															
	79	121	200															
	<table><tr><td><math>O</math></td><td><math>E</math></td><td><math>\frac{(O - E)^2}{E}</math></td></tr><tr><td>34</td><td>39.5</td><td>0.766</td></tr><tr><td>66</td><td>60.5</td><td>0.5</td></tr><tr><td>45</td><td>39.5</td><td>0.765</td></tr><tr><td>55</td><td>60.5</td><td>0.5</td></tr></table>	$O$	$E$	$\frac{(O - E)^2}{E}$	34	39.5	0.766	66	60.5	0.5	45	39.5	0.765	55	60.5	0.5	$E = \frac{RT \times CT}{GT}$ <b>M1A1A1</b>	
$O$	$E$	$\frac{(O - E)^2}{E}$																
34	39.5	0.766																
66	60.5	0.5																
45	39.5	0.765																
55	60.5	0.5																
	$\sum \frac{(O - E)^2}{E} = 2.53$ (NB with Yates 2.09)	attempt & add, awrt 0.766 & 0.5 twice, awrt 2.53																
	<b>M1A1A1</b>																	
	$\nu = 1, \chi^2(5\%) = 3.841 > 2.53$	1, 3.841 <b>B1,B1</b>																
	No reason to believe that the chance of catching a cold is affected by taking the new drug	<b>A1f</b>																
		11																

Question number	Scheme	Marks																																								
5	<p><math>\mu_a</math> and <math>\mu_b</math> are mean weight of population after and before closure respectively.</p> <p><math>H_0 : \mu_b = \mu_a</math></p> <p><math>H_1 : \mu_b &gt; \mu_a</math></p> <p><math display="block">z = \frac{10-8}{\sqrt{\frac{2.64^2}{100} + \frac{1.94^2}{120}}}</math></p> <p>Fraction, denom Ok alone</p> <p><math display="block">z = \frac{2}{\sqrt{0.1011}} = 6.29</math></p> <p>awrt 6.29</p> <p>Critical region is <math>z \geq 1.6449</math>, <math>6.29 &gt; 1.6449</math> or in critical region or Reject <math>H_0</math></p> <p>(or <math>P(Z \geq 6.29) = 0, 0 &lt; 0.05</math> or z in critical region or Reject <math>H_0</math> B1M1)</p> <p>There is evidence that closing the factory has reduced mean river pollution</p>	<p><b>B1</b></p> <p><b>B1B1</b></p> <p><b>M1A1</b> <b>M1A1</b></p> <p><b>A1</b></p> <p><b>B1, M1</b></p> <p><b>A1f</b></p> <p><b>(11)</b></p> <div>11</div>																																								
6 (a)	<table><tr><td>A</td><td>2</td><td>5</td><td>3</td><td>7</td><td>8</td><td>1</td><td>4</td><td>6</td><td></td></tr><tr><td>B</td><td>3</td><td>2</td><td>6</td><td>5</td><td>7</td><td>4</td><td>1</td><td>8</td><td></td></tr><tr><td> d </td><td>1</td><td>3</td><td>3</td><td>2</td><td>1</td><td>3</td><td>3</td><td>2</td><td></td></tr><tr><td>d<sup>2</sup></td><td>1</td><td>9</td><td>9</td><td>4</td><td>1</td><td>9</td><td>9</td><td>4</td><td>46</td></tr></table> <p><math display="block">r_s = 1 - \frac{6 \times 46}{8 \times 63}</math></p> <p><math>r_s = 0.452</math></p>	A	2	5	3	7	8	1	4	6		B	3	2	6	5	7	4	1	8		d	1	3	3	2	1	3	3	2		d <sup>2</sup>	1	9	9	4	1	9	9	4	46	<p>d M1</p> <p><math>\sum d^2</math> M1A1</p> <p><b>M1A1f</b></p> <p>0.452 <b>A1</b></p> <p><b>(6)</b></p>
A	2	5	3	7	8	1	4	6																																		
B	3	2	6	5	7	4	1	8																																		
d	1	3	3	2	1	3	3	2																																		
d <sup>2</sup>	1	9	9	4	1	9	9	4	46																																	
(b)	<p><math>H_0 : \rho = 0</math>, <math>H_1 : \rho \neq 0 (\rho &gt; 0)</math></p> <p>critical values are <math>\pm 0.7381</math> (0.6429)</p> <p><math>0.452 &lt; 0.7381</math> (<math>0.452 &lt; 0.6429</math>) or not sig or Insufficient evidence to reject <math>H_0</math></p> <p>No agreement between the two judges.</p>	<p><b>B1B1</b></p> <p>0.7381(0.6429) <b>B1</b></p> <p><b>M1</b></p> <p>Context <b>A1f</b></p> <p><b>(5)</b></p> <div>11</div>																																								

Question number	Scheme	Marks																											
7																													
(a)	$\mu = 0.3 \times 50 + 0.2 \times 10 + 0.5 \times 2 = 18$ $\sigma^2 = (0.3 \times 50^2 + 0.2 \times 10^2 + 0.5 \times 2^2) - 18^2 = 448$	M1A1 M1A1 (4)																											
(b)	<table><tr><td>(50,50)</td><td>or</td><td>(50,50) without ordered pairs</td></tr><tr><td>(10,2)</td><td></td><td>(10,2)</td></tr><tr><td>(2,10)</td><td></td><td>(10,10)</td></tr><tr><td>(10,10)</td><td></td><td>(50,10)</td></tr><tr><td>(50,10)</td><td></td><td>(2,2)</td></tr><tr><td>(10,50)</td><td></td><td>(50,2)</td></tr><tr><td>(2,2)</td><td></td><td></td></tr><tr><td>(50,2)</td><td></td><td></td></tr><tr><td>(2,50)</td><td></td><td></td></tr></table> <p>either, -1 each missing pair</p>	(50,50)	or	(50,50) without ordered pairs	(10,2)		(10,2)	(2,10)		(10,10)	(10,10)		(50,10)	(50,10)		(2,2)	(10,50)		(50,2)	(2,2)			(50,2)			(2,50)			B2 (2)
(50,50)	or	(50,50) without ordered pairs																											
(10,2)		(10,2)																											
(2,10)		(10,10)																											
(10,10)		(50,10)																											
(50,10)		(2,2)																											
(10,50)		(50,2)																											
(2,2)																													
(50,2)																													
(2,50)																													
(c)	<table><tr><td><math>\bar{x}</math></td><td>2</td><td>6</td><td>10</td><td>26</td><td>30</td><td>50</td></tr><tr><td><math>P(\bar{X} = \bar{x})</math></td><td>0.25</td><td>0.2</td><td>0.04</td><td>0.3</td><td>0.12</td><td>0.09</td></tr></table> <p>All means, probabs multiplied, -1 each error</p>	$\bar{x}$	2	6	10	26	30	50	$P(\bar{X} = \bar{x})$	0.25	0.2	0.04	0.3	0.12	0.09	B1 M1 A2 (4)													
$\bar{x}$	2	6	10	26	30	50																							
$P(\bar{X} = \bar{x})$	0.25	0.2	0.04	0.3	0.12	0.09																							
(d)	$P(2 \leq \bar{X} < 7) = 0.25 + 0.2 = 0.45$	Probabilities of 2 and 6 added, 0.45 M1 A1J (2)																											
(e)	$E(\bar{X}) = 2 \times 0.25 + 6 \times 0.2 + \dots = 18$ $\text{Var}(\bar{X}) = 2^2 \times 0.25 + 6^2 \times 0.2 + \dots - 18^2 = 224$ So $E(\bar{X}) = 18 = \mu$ and $\text{Var}(\bar{X}) = 224 = \frac{1}{2} \sigma^2$ as required.	$\sum xP(X = x)$ from table, 18 M1 A1 M1A1 A1 (5)																											

17