Centre No.					Pape	er Refer	ence			Surname	Initial(s)
Candidate No.			6	6	7	8	/	0	1	Signature	

6678/01

Edexcel GCE

Mechanics M2

Advanced/Advanced Subsidiary

Wednesday 21 May 2008 - Afternoon

Time: 1 hour 30 minutes

|--|

mathematical formulae stored in them.

Mathematical Formulae (Green)

Items included with question papers

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

If you need more space to complete your answers to any question, use additional sheets.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 7 questions in this question paper. The total mark for this paper is 75.

There are 24 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You must show sufficient working to make your methods clear to the examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

H29498A





Examiner's use only

Team Leader's use only

7

Turn over


W850/R6678/57570 3/3/3/3/2/

Question 1 continued	L t
destion 1 continued	
	-
	-
	-
	_
	_
	_
	_
	_
	_
	_
	-
	-
	-
	-
	_
	_
	_
	_
	-
	-
	-
	-
	-
	_
	_
	_
	-
	-
	-
	Q

2.	A particle A of mass $4m$ is moving with speed $3u$ in a straight line on a smooth horizontal table. The particle A collides directly with a particle B of mass $3m$ moving with speed $2u$ in the same direction as A . The coefficient of restitution between A and B is e . Immediately after the collision the speed of B is $4eu$.						
	(a) Show that $e = \frac{3}{4}$.						
	(a) Show that $e = \frac{1}{4}$.	5)					
	(b) Find the total kinetic energy lost in the collision.						
	(4) I ma the term innerte energy rose in the terms.	4)					
		_					
		-					
		-					
		_					
		_					
		_					
		_					
		_					
		_					
		_					
		-					
		-					
		_					
		_					
		_					
		_					
		_					
		_					
		_					
		_					
		_					
		-					
		-					
		-					
		-					

4

Question 2 continued	Leav blanl
	_
	_
	-
	_
	_
	_
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	Q2



Leave blank

Figure 1

A package of mass 3.5 kg is sliding down a ramp. The package is modelled as a particle and the ramp as a rough plane inclined at an angle of 20° to the horizontal. The package slides down a line of greatest slope of the plane from a point A to a point B, where AB = 14 m. At A the package has speed 12 m s⁻¹ and at B the package has speed 8 m s⁻¹, as shown in Figure 1. Find

(a) the total energy lost by the package in travelling from A to B, (5)

(b) the coefficient of friction between the package and the ramp. (5)

Question 3 continued	Leave blank
	Q3
(Total 10 marks)	

Leave
blank

1.	A particle <i>P</i> of mass 0.5 kg is moving under the action of a single force F newtons. At time
	t seconds, $\mathbf{F} = (6t - 5) \mathbf{i} + (t^2 - 2t) \mathbf{j}$.
	The velocity of P at time t seconds is \mathbf{v} m s ⁻¹ . When $t = 0$, $\mathbf{v} = \mathbf{i} - 4\mathbf{j}$.
	(a) Find v at time t seconds. (6)
	When $t = 3$, the particle P receives an impulse $(-5\mathbf{i} + 12\mathbf{j}) \mathrm{N}\mathrm{s}$.
	(b) Find the speed of P immediately after it receives the impulse. (6)
_	

	Leave blank
Question 4 continued	Diank

Question 4 continued	

Question 4 continued	Lea blar
	_
	_
	_
	_
	_
	_
	_
	-
	-
	-
	_
	_
	_
	-
	-
	-
	_
	_
	_
	-
	-
	_
	_
	_
	_
	_
	-
	-
	_ Q




Figure 2

A plank rests in equilibrium against a fixed horizontal pole. The plank is modelled as a uniform rod AB and the pole as a smooth horizontal peg perpendicular to the vertical plane containing AB. The rod has length 3a and weight W and rests on the peg at C, where AC = 2a. The end A of the rod rests on rough horizontal ground and AB makes an angle α with the ground, as shown in Figure 2.

(a) Show that the normal reaction on the rod at A is $\frac{1}{4}(4-3\cos^2\alpha)W$.

Given that the rod is in limiting equilibrium and that $\cos \alpha = \frac{2}{3}$,

 \boldsymbol{A}

(b) find the coefficient of friction between the rod and the ground.

(5)

Leave blank

Question 5 continued	Leave blank

Question 5 continued	b

Question 5 continued	Le
destion 5 continued	
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_






Figure 3

Figure 3 shows a rectangular lamina OABC. The coordinates of O, A, B and C are (0, 0), (8, 0), (8, 5) and (0, 5) respectively. Particles of mass km, 5m and 3m are attached to the lamina at A, B and C respectively.

The *x*-coordinate of the centre of mass of the three particles *without the lamina* is 6.4.

(a) Show that k = 7.

(4)

The lamina OABC is uniform and has mass 12m.

(b) Find the coordinates of the centre of mass of the combined system consisting of the three particles and the lamina.

(6)

The combined system is freely suspended from O and hangs at rest.

(c) Find the angle between OC and the horizontal.

(3)

Question 6 continued	Leave blank

Question 6 continued	b

Question 6 continued	Le bl




Figure 4

A ball is thrown from a point A at a target, which is on horizontal ground. The point A is 12 m above the point O on the ground. The ball is thrown from A with speed 25 m s⁻¹ at an angle of 30° below the horizontal. The ball is modelled as a particle and the target as a point T. The distance OT is 15 m. The ball misses the target and hits the ground at the point B, where OTB is a straight line, as shown in Figure 4. Find

(a) the time taken by the ball to travel from A to B,

(5)

Leave blank

(b) the distance TB.

(4)

The point X is on the path of the ball vertically above T.

(c) Find the speed of the ball at X.

(5)

Question 7 continued	Leave blank

Question 7 continued	

Question 7 continued		Le
	(Total 14 marks)	(
	TOTAL FOR PAPER: 75 MARKS	

