Centre No.			Paper Reference Surname						Initial(s)			
Candidate No.			6	6	7	8	/	0	1	Signature		
		r Reference									Examiı	ner's use only

Edexcel GCE

Mechanics M2

Advanced/Advanced Subsidiary

Friday 22 May 2009 - Morning

Time: 1 hour 30 minutes

Materials required for examination
Mathematical Formulae (Orange or

Items included with question papers

Nil

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions to Candidates

In the boxes above, write your centre number, candidate number, your surname, initials and signature. Check that you have the correct question paper.

Answer ALL the questions.

You must write your answer to each question in the space following the question.

If you need more space to complete your answers to any question, use additional sheets.

Whenever a numerical value of g is required, take $g = 9.8 \text{ m s}^{-2}$.

When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information for Candidates

A booklet 'Mathematical Formulae and Statistical Tables' is provided.

Full marks may be obtained for answers to ALL questions.

Green)

The marks for individual questions and the parts of questions are shown in round brackets: e.g. (2).

There are 8 questions in this question paper. The total mark for this paper is 75.

There are 28 pages in this question paper. Any blank pages are indicated.

Advice to Candidates

You must ensure that your answers to parts of questions are clearly labelled. You must show sufficient working to make your methods clear to the examiner. Answers without working may not gain full credit.

This publication may be reproduced only in accordance with Edexcel Limited copyright policy.

N34274A W850/R6678/57570 3/5/4/3





Team Leader's use only

Find the speed of the particle immediately after the impulse.					
	(5)				

Question 1 continued	Leav
	21

2. At time $t = 0$ a particle P leaves the origin O and moves along the x-axis. At the velocity of P is v m s ⁻¹ , where	time t seconds
$v=8t-t^2.$	
(a) Find the maximum value of v .	(4)
	(4)
(b) Find the time taken for <i>P</i> to return to <i>O</i> .	(5)

Question 2 continued	Leave blank
	Q2
(Total 9 marks)	

3.	A truck of mass of 300 kg moves along a straight horizontal road with a constant speed of 10 m s^{-1} . The resistance to motion of the truck has magnitude 120 N .
	(a) Find the rate at which the engine of the truck is working. (2)
	On another occasion the truck moves at a constant speed up a hill inclined at $ heta$ to the
	horizontal, where $\sin \theta = \frac{1}{14}$. The resistance to motion of the truck from non-gravitational
	forces remains of magnitude 120 N. The rate at which the engine works is the same as in part (a).
	(b) Find the speed of the truck.
	(4)

	Leave blank
Question 3 continued	
	Q3
(Total 6 marks)	




Figure 1

A uniform rod AB, of length 1.5 m and mass 3 kg, is smoothly hinged to a vertical wall at A. The rod is held in equilibrium in a horizontal position by a light strut CD as shown in Figure 1. The rod and the strut lie in the same vertical plane, which is perpendicular to the wall. The end C of the strut is freely jointed to the wall at a point 0.5 m vertically below A. The end D is freely joined to the rod so that AD is 0.5 m.

(a) Find the thrust in CD.

(4)

Leave blank

(b) Find the magnitude and direction of the force exerted on the rod AB at A.

(7)

Question 4 continued	Leave blank

Question 4 continued	

Question 4 continued		Leave blank
		Q4
	Гotal 11 marks)	




Figure 2

A shop sign ABCDEFG is modelled as a uniform lamina, as illustrated in Figure 2. ABCD is a rectangle with BC = 120 cm and DC = 90 cm. The shape EFG is an isosceles triangle with EG = 60 cm and height 60 cm. The mid-point of AD and the mid-point of EG coincide.

(a) Find the distance of the centre of mass of the sign from the side AD.

(5)

Leave blank

The sign is freely suspended from A and hangs at rest.

(b) Find the size of the angle between AB and the vertical.

(4)

Question 5 continued	Leave blank

Question 5 continued	b]

Question 5 continued	Leave blank
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	Q5
(Total 9 mark	as)



Leave blank




Figure 3

A child playing cricket on horizontal ground hits the ball towards a fence 10 m away. The ball moves in a vertical plane which is perpendicular to the fence. The ball just passes over the top of the fence, which is 2 m above the ground, as shown in Figure 3.

The ball is modelled as a particle projected with initial speed u m s⁻¹ from point O on the ground at an angle α to the ground.

(a) By writing down expressions for the horizontal and vertical distances, from O of the ball t seconds after it was hit, show that

$$2 = 10 \tan \alpha - \frac{50g}{u^2 \cos^2 \alpha}.$$

(6)

Given that $\alpha = 45^{\circ}$,

(b) find the speed of the ball as it passes over the fence.

(6)



Question 6 continued	Leave blank

Question 6 continued	

uestion 6 continued		L b
	(Total 12 marks)	




Figure 4

A particle P of mass 2 kg is projected up a rough plane with initial speed 14 m s⁻¹, from a point X on the plane, as shown in Figure 4. The particle moves up the plane along the line of greatest slope through X and comes to instantaneous rest at the point Y. The plane is

inclined at an angle α to the horizontal, where $\tan \alpha = \frac{7}{24}$. The coefficient of friction between the particle and the plane is $\frac{1}{8}$.

(a) Use the work-energy principle to show that XY = 25 m.

(7)

Leave blank

After reaching *Y*, the particle *P* slides back down the plane.

(b) Find the speed of P as it passes through X.

(4)

Question 7 continued		Leave blank
	Question 7 continued	

Question 7 continued	

Question 7 continued	Leave blank
	Q7
(Total 11 marks)	

3.	Particles A, B and C of masses 4m, 3m and m respectively, lie at rest in a straight line on a smooth horizontal plane with B between A and C. Particles A and B are projected towards
	each other with speeds u m s ⁻¹ and v m s ⁻¹ respectively, and collide directly.
	As a result of the collision, A is brought to rest and B rebounds with speed kv m s ⁻¹ . The
	coefficient of restitution between A and B is $\frac{3}{4}$.
	(a) Show that $u = 3v$.
	(6)
	(b) Find the value of k . (2)
	Immediately after the collision between A and B , particle C is projected with speed $2v$ m s ⁻¹ towards B so that B and C collide directly.
	(c) Show that there is no further collision between A and B.
	(4)

	Leave blank
Question 8 continued	

Question 8 continued	b

	Leave
Question 8 continued	blank

Question 8 continued		blank
		Q
	(Total 12 marks)	
	TOTAL FOR PAPER: 75 MARKS	
END		