

Serre Equations with non flat bottom:

$$\frac{\partial G}{\partial t} + \frac{\partial}{\partial x} \left( G\bar{u} + \frac{gh^2}{2} - \frac{2h^3}{3} \left( \frac{\partial \bar{u}}{\partial x} \right)^2 \right) = -\frac{\partial}{\partial x} \left( \frac{h^2}{2} \Psi \right) - \left( gh + \frac{h^2}{2} \Phi + h\Psi \right) \frac{\partial b}{\partial x}$$

where:

$$G = \bar{u}h - h^2 \frac{\partial h}{\partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^3}{3} \frac{\partial^2 \bar{u}}{\partial x^2}$$

However I'm going to assume that we do not want temporal derivative in our source term, so we want to modify G so that we can do this. The important thing here to note is that b is independent of time. By simple calculus it is true that:

$$\frac{\partial}{\partial t} \left( \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} \right) = h \frac{\partial h}{\partial t} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^2}{2} \frac{\partial^2 b}{\partial t \partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial^2 \bar{u}}{\partial t \partial x}$$

since b is independent of time, so is its derivative with respect to x and so  $\frac{\partial^2 b}{\partial t \partial x} = 0$ . Thus:

$$\frac{\partial}{\partial t} \left( \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} \right) = h \frac{\partial h}{\partial t} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial^2 \bar{u}}{\partial t \partial x}$$

Applying the equation for conservation of mass [] gives:

$$\frac{\partial}{\partial t} \left( \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} \right) = h \frac{\partial(\bar{u}h)}{\partial x} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial^2 \bar{u}}{\partial t \partial x}$$

$$\frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial^2 \bar{u}}{\partial t \partial x} = h^2 \frac{\partial \bar{u}}{\partial x} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} + h\bar{u} \frac{\partial h}{\partial x} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} - \frac{\partial}{\partial t} \left( \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x} \right)$$

So the first equation can be rearranged to give:

$$\frac{\partial G}{\partial t} + \frac{\partial}{\partial x} \left( G\bar{u} + \frac{h^2}{2} \Psi + \frac{gh^2}{2} - \frac{2h^3}{3} \left( \frac{\partial \bar{u}}{\partial x} \right)^2 \right) - \frac{h^2}{2} \frac{\partial^2 \bar{u}}{\partial x \partial t} \frac{\partial b}{\partial x} = - \left( gh + \frac{h^2}{2} \left( \frac{\partial \bar{u}}{\partial x} \right)^2 - \bar{u} \frac{\partial^2 \bar{u}}{\partial x^2} + h\Psi \right) \frac{\partial b}{\partial x}$$

$$\frac{\partial H}{\partial t} + \frac{\partial}{\partial x} \left( H\bar{u} + \frac{gh^2}{2} - \frac{2h^3}{3} \left( \frac{\partial \bar{u}}{\partial x} \right)^2 \right) = - \left( gh - \frac{h^2}{2} \left( \frac{\partial \bar{u}}{\partial x} \right)^2 - \bar{u} \frac{\partial^2 \bar{u}}{\partial x^2} + h\Psi - h\bar{u} \frac{\partial h}{\partial x} \frac{\partial \bar{u}}{\partial x} \right) \frac{\partial b}{\partial x}$$

Where:

$$H = \bar{u}h - h^2 \frac{\partial h}{\partial x} \frac{\partial \bar{u}}{\partial x} + \frac{h^3}{3} \frac{\partial^2 \bar{u}}{\partial x^2} + \frac{h^2}{2} \frac{\partial b}{\partial x} \frac{\partial \bar{u}}{\partial x}$$