
a Two routes to opposite biases
probe bias, exemplars held out
shape − texture accuracy
diet
no diet
magno, diet +0.13
shape
parvo, no diet −0.08
magno, no diet −0.21
parvo, diet −0.14
texture
untrained
trained
(150 epochs)
b Carrier × transform: probe
probe bias, exemplars held out
chromatic (L−M)
achromatic (luminance)
+0.34
no
transform
with
transform
c ...and behaviour
cue-conflict shape bias
0.88
no
transform
with
transform
d How each readout decides the images
cue-conflict images (of 1,200)
decided as shape
decided as texture
neither category
shape bias
0.90 0.42
0.65 0.42
0.40 0.36
diet no diet
magno
only
diet no diet
both
w = 0.5
diet no diet
parvo
only
e Accuracy-matched control: probe
magno-stream probe bias,
exemplars held out
gap
+0.46
closes 4.3%
of the gap
diet
0.556
no diet
0.70
no diet,
matched
0.55 (n=3)
validation accuracy at the scored checkpoint
f ...and behaviour
magno-stream shape bias
(cue-conflict)
gap
+0.48
closes 6.1%
of the gap
diet
0.556
no diet
0.70
no diet,
matched
0.55 (n=3)
validation accuracy at the scored checkpoint