
a Normalised index (as Fig. 3a)
normalised index
median of (Um − Up)/(Um + Up)
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
untrained (diet front end)
no diet (3 seeds)
diet (3 seeds)
V1v
V1d
V2v
V2d
V3v
V3d
hV4
b Raw statistic
raw statistic
median of Um − Up (units of r² / NC)
0.20
0.15
0.10
0.05
0.00
−0.05
−0.10
−0.15
V1v
V1d
V2v
V2d
V3v
V3d
hV4
c Gradient within eccentricity bands
gradient hV4 − mean(V1v, V1d)
within stratum (normalised)
labels: subjects with diet < no diet; exact Wilcoxon p
diet
no diet
0.4
0.2
0.0
−0.2
−0.4
−0.6
−0.8
−1.0
−1.2
7/7
p = 0.016
7/8
p = 0.016
8/8
p = 0.008
8/8
p = 0.008
8/8
p = 0.008
0–0.5°
0.5–1°
1–2°
2–4°
> 4°
pRF eccentricity band (NSD prf-eccrois)
d Gradient within pRF-size tertiles
gradient hV4 − mean(V1v, V1d)
within stratum (normalised)
labels: subjects with diet < no diet; exact Wilcoxon p
diet
no diet
0.4
0.2
0.0
−0.2
−0.4
−0.6
−0.8
−1.0
−1.2
6/6
p = 0.031
8/8
p = 0.008
8/8
p = 0.008
small
medium
large
pRF-size tertile (per subject, V1 + hV4 voxels)