
a Two routes to opposite biases
linear-probe bias
shape − texture accuracy
diet
no diet
1.0
0.5
0.0
−0.5
shape
texture
magno, diet +0.02
magno, no diet -0.64
parvo, no diet -0.42
parvo, diet -0.43
untrained
trained
(150 epochs)
b Carrier × transform: probe
linear-probe bias (shape − texture)
chromatic carrier (L−M)
achromatic carrier (luminance)
+0.82
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
no
transform
with
transform
c ...and behaviour
cue-conflict shape bias
0.88
1.0
0.8
0.6
0.4
no
transform
with
transform
d How each readout decides the images
cue-conflict images (of 1,200)
decided as shape
decided as texture
neither category
shape bias
0.90 0.42
0.65 0.42
0.40 0.36
1200
800
400
0
diet no diet
magno
only
diet no diet
both
w = 0.5
diet no diet
parvo
only
e Accuracy-matched control: probe
magno-stream probe bias
1.00
0.75
0.50
0.25
0.00
−0.25
gap
+1.03
closes 7.2%
of the gap
diet
0.556
no diet
0.70
no diet,
matched
0.55 (n=3)
validation accuracy at the scored checkpoint
f ...and behaviour
magno-stream shape bias
(cue-conflict)
1.0
0.9
0.8
0.7
0.6
0.5
0.4
0.3
gap
+0.48
closes 6.1%
of the gap
diet
0.556
no diet
0.70
no diet,
matched
0.55 (n=3)
validation accuracy at the scored checkpoint