
a Which stream explains which level of the ventral hierarchy
b Learning builds it or erodes it
c The diet gradient is near-perfectly monotonic
d Cue conflict, through the fitted cortex
e What the diet costs
stream preference per ROI
median of (Um − Up) / (Um + Up)
↑ magno-explained
↓ parvo-explained
trained, swap (1 seed)
trained, no diet (3 seeds)
line: mean over 8 subjects; shading: ± SEM
untrained (diet front end)
trained, diet (3 seeds)
V1v
V1d
V2v
V2d
V3v
V3d
hV4
retinotopic area, V1 → hV4 (Wang et al. 2015 atlas)
gradient hV4 − mean(V1v, V1d)
-0.07 no diet
-0.54
-0.79 diet
diet steeper in 8/8 subjects, p = 0.0078
untrained
trained
Spearman ρ, hierarchical rank vs stream preference (per subject)
shaded: within the exact null over all 630 labellings of the 7 ROIs (2.5th percentile as dashed rule)
diet: the most extreme of all 630 labellings, 3/3 seeds, p = 0.0016
-0.71
-0.58
-0.97
untrained
trained, no diet
trained, diet
predicted shape-bias index magno − parvo (per subject)
p = 0.0078
8/8 subjects
+0.11
+1.53
no diet
diet
median r² / noise ceiling (reliable voxels, nsdgeneral)
the diet predicts BOLD 13% worse
p = 0.0078, 8/8 subjects
0.464
0.405
no diet
diet