

a Predicted cortex is not texture-biased on these images
cortical shape-bias rate B
P(shape template closer than texture template)
observers 0.96
model (3 seeds)
control (3 seeds)
channel swap (3 seeds)
untrained
input floor (oriented energy)
model's readout 0.68
V1
V2
V3
hV4
ventral
lateral
parietal
b Evidence present, but matched by the input floor beyond hV4
model (per subject)
input floor
permutation null, 97.5th pct
shape margin v shape rate
% of ceiling (0.62)
cue margin v resolution rate
% of ceiling (0.73)
7/8
8/8
8/8
7/8
0/8
1/8
1/8
7/8
8/8
8/8
7/8
0/8
3/8
1/8
c Opposite evidence in the same voxels
magno > parvo: 8/8 subjects, 24/24 seed-cells,
every region; joint between them in 8/8
cortical shape-bias rate B, model
same voxels, one band of the fit
luminance band (magno)
joint fit
L−M band (parvo)
d The trajectory adds nothing
extrastriate correspondence per seed
% of ceiling (hV4, ventral, lateral mean)
model
control
swap
floor
untrained
MW p
1.00 / 1.00
MW p
0.10 / 0.10
shape
target
resolution
target
e hV4, image by image
r = 0.108 [0.050, 0.155] exemplar bootstrap
17% of ceiling; per subject 13–18%
n = 1194 images
observers' shape rate, residualised
predicted hV4 shape margin, residualised
(subject mean; Fisher z)