
Gene and TE density across three Drosophila genomes

Dmir → Dpse → Dmel, outside to inside · four tracks per species

A | TE-class composition per chromosome



Common 0–1 scales: gene density = genes per 100 kb / 140; TE density = TE-covered bases / assembled window span.

Solid segments show each class's share of TE-covered bases; segment positions are not genomic coordinates. Mixed = overlapping classes.

† Dmir Y: all 62 Y-associated scaffolds (110.50 Mb); length-ordered for display only. Ticks show cumulative span; small marks show boundaries.

Each species spans the same arc independently: angular alignment does not imply synteny. Non-Y unplaced scaffolds and mitochondria are not shown.


* Dmir XR includes A-derived material; Dpse X combines A + D. Gene/TE densities use assembled span; terminal windows use their actual length.

Dated tree: Gao et al. (2007) model; plotted range midpoints 32.5 and 4.35 Ma, ranges 30–35 and 3.76–4.93 Ma (grey bars). Ages are model-dependent.

Gene and TE density across three Drosophila genomes

Dmir → Dpse → Dmel, outside to inside · four tracks per species

B | Dominant TE class along each chromosome



Common 0–1 scales: gene density = genes per 100 kb / 140; TE density = TE-covered bases / assembled window span.

Solid colours show the largest class by covered bases in each 100 kb window; minority classes are not displayed. Mixed = overlapping classes.

† Dmir Y: all 62 Y-associated scaffolds (110.50 Mb); length-ordered for display only. Ticks show cumulative span; small marks show boundaries.

Each species spans the same arc independently: angular alignment does not imply synteny. Non-Y unplaced scaffolds and mitochondria are not shown.

* Dmir XR includes A-derived material; Dpse X combines A + D. Gene/TE densities use assembled span; terminal windows use their actual length.

Dated tree: Gao et al. (2007) model; plotted range midpoints 32.5 and 4.35 Ma, ranges 30–35 and 3.76–4.93 Ma (grey bars). Ages are model-dependent.