

MATH 646: Homework 2

1. Let $M = (0, \infty)$ equipped with the Lebesgue measure and consider the map $T : M \rightarrow M$ given by

$$Tx = \begin{cases} \frac{1}{x} & \text{if } x \in (0, 1] \\ x - 1 & \text{if } x \in (1, \infty). \end{cases}$$

What is the induced transformation T_A where $A = (0, 1]$? Show that the measure (called Gauss measure) μ given by

$$\mu(B) = \frac{1}{\log 2} \int_B \frac{1}{1+x} dx,$$

is T_A invariant.

2. Let $\{a_n\}$ be a sequence of complex numbers. Show that

$$\lim_{n \rightarrow \infty} a_n = 0,$$

implies that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} |a_j| = 0,$$

which in turn implies that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{j=0}^{n-1} a_j = 0.$$

but that the reverse implications are wrong, in general.

3. Let T be an endomorphism of (M, \mathcal{B}, μ) . Show that the following are equivalent
 - (a) $\lim_{n \rightarrow \infty} \mu(T^{-n}(A) \cap B) = \mu(A)\mu(B)$ for all $A, B \in \mathcal{B}$
 - (b) $\lim_{n \rightarrow \infty} \langle U_T^n f, g \rangle = \langle f, 1 \rangle \langle 1, g \rangle$ for all $f, g \in L^2(\mu)$.
 - (c) $\lim_{n \rightarrow \infty} \langle U_T^n f, f \rangle = \langle f, 1 \rangle \langle 1, f \rangle$ for all $f \in L^2(\mu)$.
 - (d) $\lim_{n \rightarrow \infty} \langle U_T^n f, f \rangle = \langle f, 1 \rangle \langle 1, f \rangle$ for all $f \in L^2(\mu)$ with $\langle f, 1 \rangle = 0$.
4. Let T be an endomorphism of (M, \mathcal{B}, μ) . Let ρ be a probability measure (M, \mathcal{B}) and let ρ^n be the measure define by $\rho^n(A) = \rho(T^{-n}A)$, $A \in \mathcal{B}$. Show that T is mixing if and only if $\lim_{n \rightarrow \infty} \rho^n(A) = \mu(A)$ for all $A \in \mathcal{B}$ and for all ρ which are absolutely continuous with respect to μ .
5. Let T be an endomorphism of (M, \mathcal{B}, μ) and let $\{f_j\}$ be a orthonormal basis for $L^2(\mu)$. Show that T is mixing if and only if $\lim_{n \rightarrow \infty} \langle U_T^n f_j, f_k \rangle = \langle f_j, 1 \rangle \langle 1, f_k \rangle$ for all j, k .

6. **A weak mixing transformation which is not strong mixing.** Consider the measure space $([0, 1], \mathcal{B}, m)$ where \mathcal{B} is the Borel σ -algebra and m Lebesgue measure. We define the map T by

$$Tx = \begin{cases} x + 1/2 & \text{if } x < 1/2 \\ x - 1/4 & \text{if } 1/2 < x < 3/4 \\ x - 5/8 & \text{if } 3/4 < x < 7/8 \\ \vdots & \vdots \\ x - (1 - 1/2^n - 1/2^{n+1}) & \text{if } 1 - 1/2^n < x < 1 - 1/2^{n+1} \\ \vdots & \vdots \end{cases}$$

Note that the map T is not defined for dyadic rationals $j/2^n$ which can be ignored in the sequel since they form a set of measure 0. Note further that, for any n , the map T permutes the dyadic intervals $(j/2^n, (j+1)/2^n)$, although not in accordance with the order on the intervals.

- Show that the map T is ergodic and has pure point spectrum: to do this let $K_n = (0, 1/2^n)$ and show that

$$f_n(x) = \sum_{j=0}^{2^n-1} \omega_n^{-j} \chi_{K_n}(T^j x)$$

are eigenfunctions for the eigenvalues $\omega_n = \exp(2\pi i/2^n)$.

Let

$$I_0 = (0, 1/2), I_1 = (1/2, 3/4), I_2 = (3/4, 7/8), \dots$$

We have $[0, 1] = M = \bigcup_{n=0}^{\infty} I_n \bmod 0$ and we set $A = \sum_{n=0}^{\infty} I_{2n}$. Let A' be a copy of A and set $M^A = M \cup A'$ (disjoint union) and define the map $T^A : M^A \rightarrow M^A$ by

$$\begin{aligned} T^A x' &= Tx & \text{if } x' \in A' \\ T^A x &= x' & \text{if } x \in A \\ T^A x &= Tx & \text{if } x \in M \setminus A \end{aligned}$$

This is the integral automorphism corresponding to the function $f(x) = 1 + \chi_A(x)$ as defined in Homework 1.

- Show (directly) that the map T_A is ergodic.

Consider the function

$$S_n = \chi_A + \chi_A \circ T + \dots + \chi_A \circ T^{2^n-1}$$

which is the number of visits x makes to A in the first 2^n iterations of T .

- Show that S_n assumes only two values, s_n and $s_n + 1$ and

$$m\{x : S_n(x) = s_n + 1\} = \frac{2}{3} \text{ or } \frac{1}{3},$$

depending whether n is even or odd. *Hint:* If $J_n = (1 - (1/2^n), 1)$, each of the iterations $TJ_n, \dots, T^{2^n-1}J_n$ is either contained in A or disjoint from A .

- Show that $(T^A)^{2^n+S_n(x)}(x) = T^{2^n}(x)$ for all $x \in M$ and that

$$(T^A)^{2^n+s_n+1}I_0 \subset I_0 \cup I'_0.$$

Deduce that this excludes strong mixing.

Next we show that T^A is weak-mixing.

- Let f be an eigenfunction for U_{T^A} with eigenvalues $e^{2\pi i\alpha}$ ($0 \leq \alpha < 1$ and $|f| = 1$). Show that

$$\left| \int f(T^{2^n}x) \bar{f}(x) dm \right| = \left| \int e^{2\pi i\alpha S_n(x)} dm \right| < 1$$

if $\alpha \neq 0$.

- Approximate f by linear combinations of characteristic functions of dyadic intervals and conclude that α must be 0.