

STAT 315: Probability Basics (Section 2.2–2.5)

Luc Rey-Bellet

University of Massachusetts Amherst

luc@math.umass.edu

January 30, 2025

What is probability?

- Probability describe an experiment whose outcome cannot be described with certainty. Keyword: **random** or **stochastic**
 - ▶ Roll of a dice
 - ▶ The price of GameStop stock or the price of Bitcoin tomorrow
 - ▶ The winner of Super Bowl LVIX: Chiefs or Eagles?
 - ▶ The amount of rain on Amherst due to Hurricane Ida
- **Frequentist approach**: deduce the probability by repeating the experiment N times (N very large) (**Law of Large numbers**)

$$P(\text{roll a 5}) \approx \frac{\text{number of 5 in } N \text{ rolls}}{N}$$

→ Can be simulated on a computer!

- **Subjective (Bayesian) approach**: Probability is a measures of one's belief in the occurrence of a future event
 - You can't repeat the Super Bowl but you can bet on it!

Review of set notation

- Denote sets of points by capital letters, A, B_1, B_2, S, \dots , and points by lower case, a_1, a_2, b, c, x, \dots
- If the elements in A are a_1, a_2, a_3 we write $A = \{a_1, a_2, a_3\}$
- Denote by S the sets of all elements (the **sample space**) and by \emptyset the set with no element.

Sets operations

- $A \subset B$ (A is contained in B): every element in A is also in B .
- $A \cup B$ (the union of A and B): the set elements which belong **either to A or to B** .
- $A \cap B$ (the intersection of A and B): the sets of elements which belong **both to A and B** .
- \bar{A} (the complement of A): the set of element in S which do not belong to A .
- $B \setminus A = B \cap \bar{A}$: the elements in B which are not in A .

Laws of set algebra

Distributive Laws

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

$$A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$$

DeMorgan's Law

$$\overline{A \cap B} = \overline{A} \cup \overline{B}$$

$$\overline{A \cup B} = \overline{A} \cap \overline{B}$$

Use Venn's diagrams

Probabilistic experiment: discrete case

We use the language of set theory to describe an experiment with random outcomes:

Sample space and events

- S is called the **sample space**: the list of all possible outcomes of the experiment.

$$S = \{e_1, e_2, e_3, \dots\} \quad \text{finite or countable set}$$

- A subset $A \subset S$ is called an **event**. Think of it as a question you ask about the experiment: **does the event A occur?**)

Example (Roll a pair of dice)

$$S = \{(1, 1), (1, 2), (1, 3), \dots, (6, 6)\} \quad S \text{ has 36 elements}$$

$$A = \{\text{The sum of the dice is 4}\} = \{(1, 3), (2, 2), (3, 1)\}$$

$$B = \{\text{Exactly one 6}\} = \{(1, 6), (6, 1), (2, 6), (6, 2), \dots, (5, 6), (6, 5)\}$$

Set operations in probabilistic language

Intuitive meaning of set operations

- $A \cap B = \text{"}A \text{ and } B\text{"}$ \longrightarrow both A and B occur.
- $A \cup B = \text{"}A \text{ or } B\text{"}$ \longrightarrow either A or B occur.
- $\bar{A} = \text{"not } A\text{"}$ \longrightarrow A does not occur.
- $A \cap B = \emptyset \longrightarrow A \text{ and } B \text{ are mutually exclusive, they cannot occur simultaneously.}$
- $B \setminus A = B \cap \bar{A} = \text{"}B \text{ but not } A\text{"}$, B occurs but not A

Laws of probability I

Laws of Probability

S is the sample space. To every event A in S (i.e., $A \subset S$) we assign a number $P(A)$, called the probability of A , with the following properties

- Axiom 1: $0 \leq P(A) \leq 1$.
- Axiom 2: $P(S) = 1$.
- Axiom 3: If A_1, A_2, A_3, \dots are pairwise mutually exclusive $A_i \cap A_j = \emptyset$ if $i \neq j$ then

$$P(A_1 \cup A_2 \cup A_3 \cup \dots) = P(A_1) + P(A_2) + P(A_3) + \dots$$

For the discrete case $S = \{e_1, e_2, \dots\}$ simply assign numbers $P(\{e_i\}) = p_i$ with $0 \leq p_i \leq 1$ and $\sum_i p_i = 1$. We have then


$$P(A) = \sum_{i: e_i \in A} p_i.$$

Laws of probability II

Simple consequence of the laws of probability

- ① $P(\emptyset) = 0$
- ② If $A \cap B = \emptyset$ (mutually exclusive) then $P(A \cup B) = P(A) + P(B)$
- ③ $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ (Inclusion-Exclusion)
- ④ If $A \subset B$ then $P(A) \leq P(B)$ (Monotonicity)
- ⑤ $P(\overline{A}) = 1 - P(A)$ (Complement)

The Monty Hall Problem



- There are 2 goats and \$1 million hidden behind 3 doors.
- You pick a door (without opening it), say door 1.
- The game hosts open of the other two door (on the picture, door 3) and reveals a goat behind it.
- You are given the following xchoice
 - Keep your door?
 - Switch?
- What should you do to maximize your probability of winning?

Any questions?