

Math 645: Homework 2

1. Show the following: If the Cauchy problem $x' = f(t, x)$, $x(t_0) = x_0$, where $f(t, x)$ is a continuous function, has a unique solution, then the Euler polygons converge to this solution.
2. Consider the Cauchy problem $x' = f(t, x)$, $x(0) = 0$, where

$$f(t, x) = \begin{cases} 4\operatorname{sign}(x)\sqrt{|x|} & \text{if } |x| \geq t^2 \\ 4\operatorname{sign}(x)\sqrt{|x|} + 4(t - \frac{|x|}{t}) \cos(\pi \frac{\log t}{\log 2}) & \text{if } |x| < t^2 \end{cases} \quad (1)$$

The function f is continuous on \mathbf{R}^2 . Consider the Euler polygons $x_h(t)$ with $h = 2^{-i}$, $i = 1, 2, 3, \dots$. Show that $x_h(t)$ does not converge as $h \rightarrow 0$, compute its accumulation points, and show that they are solution of the Cauchy problem. *Hint:* $\pm 4t^2$.

3. Consider the the Cauchy problem $x' = f(t, x)$, $x(0) = 0$ where f is given by

$$f(t, x) = \begin{cases} 0 & \text{if } t \leq 0, \ x \in \mathbf{R} \\ 2t & \text{if } t > 0, \ x \leq 0 \\ 2t - \frac{4x}{t} & \text{if } t > 0, \ 0 \leq x < t^2 \\ -2t & \text{if } t > 0, \ t^2 \leq x \end{cases} \quad (2)$$

- (a) Show that f is continuous. What does that imply for the Cauchy problem?
- (b) Show that f does not satisfy a Lipschitz condition in any neighborhood of the origin.
- (c) Apply Picard-Lindelöf iteration with $x_0(t) \equiv 0$. Are the accumulation points solutions?
- (d) Show that the Cauchy problem has a unique solution. What is the solution?

Note that this problem shows that existence and uniqueness of the solution does not imply that the Picard-Lindelöf iteration converges to the unique solution.

4. Consider the Cauchy problem $x' = \lambda x$, $x(0) = 1$, with $\lambda > 0$ on the interval $[0, 1]$. Compute the Euler polygons $x_h(t)$. Show that

$$\frac{\lambda}{1 + \lambda h} x_h(t) \leq \frac{d}{dt} x_h(t) \leq \lambda x_h(t). \quad (3)$$

and deduce from this the classical inequality

$$\left(1 + \frac{\lambda}{n}\right)^n \leq e^\lambda \leq \left(1 + \frac{\lambda}{n}\right)^{n+\lambda} \quad (4)$$

5. Consider the Cauchy problem for the Ricatti equation $x' = t^2 + x^2$, $x(0) = 1$. Show that the solution $x(t)$ satisfies the bounds

$$\frac{1}{1-t} \leq x(t) \leq \tan(t + \pi/4) \quad (5)$$

Hint: Consider the Cauchy problems $u' = u^2$ and $v' = 1 + v^2$ and compare $x(t)$ with $u(t)$ and $v(t)$.

6. Let $V(q)$ be a function of class \mathcal{C}^2 such that $\lim_{|q| \rightarrow \infty} V(q) = \infty$ and consider the Hamiltonian system $q' = p$, $p' = -dV(q)/dq$. Show that the solution exists for all positive as well as all negative times.
7. Let a , b , c , and d be positive constants. Consider the Predator-Prey equation $x' = x(a - by)$, $y' = y(cx - d)$ with positive initial conditions $x(t_0)$ and $y(t_0)$. Using the change of variables $p = \log(x)$ and $q = \log(y)$ express the equations as an Hamiltonian systems and deduce that the solution exist for all t .
8. Find the maximal interval of existence $I_{\max} = (\omega_-, \omega_+)$ for the following Cauchy problems.
- (a) $x' = x^3$.
 - (b) $x' = e^{-x}$.
 - (c) $x'_1 = 1/x_1$, $x'_2 = x_1$.
 - (d) $x' = 1 - x^2$.
 - (e) $x'' + x + x^3 = 0$.
 - (f) $x'' + x' + x + x^3 = 0$.
 - (g) $x' = \sin(2tx)x^3/(1 + t^2 + x^2)$.
9. Prove the generalized Gronwall Lemma. Let $h(t)$ and $g(t)$ be continuous nonnegative function and let $a > 0$ be a positive constant such that, for any $t \in [0, T]$

$$g(t) \leq a + \int_0^t h(s)g(s) ds. \quad (6)$$

Then, for any $t \in [0, T]$

$$g(t) \leq ae^{\int_0^t h(s) ds}. \quad (7)$$

10. Consider the FitzHugh-Nagumo equation

$$\begin{aligned} x'_1 &= f_1(x_1, x_2) = g(x_1) - x_2, \\ x'_2 &= f_2(x_1, x_2) = \sigma x_1 - \gamma x_2, \end{aligned} \quad (8)$$

where σ and γ are positive constants and the function g is given by $g(x) = -x(x - 1/2)(x - 1)$.

- (a) In the x_1 - x_2 plane draw the graph of the curves $f_1(x_1, x_2) = 0$ and $f_2(x_1, x_2) = 0$.
- (b) Consider the rectangles $ABCD$ whose sides are parallel to the X_1 and x_2 axis with two opposite corners located on the $f_2(x_1, x_2) = 0$. Show that if the rectangle is taken sufficiently large, a solution which start inside the rectangle stays inside the rectangle forever. Deduce from this that the equations for any initial conditions x_0 have a unique solutions for all time $t > 0$.

11. Show that the solutions of

$$\begin{aligned}x_1' &= x_1(3 - x_1 - 2x_2), \\x_2' &= x_2(4 - 2x_1 - 3x_2),\end{aligned}\tag{9}$$

have a unique solution for all $t \geq 0$, for any initial conditions x_{10} , x_{20} wich are nonnegative. *Hint:* A possibility is to use a similar procedure as in the previous exercise.